



## Abstract

This study suggested an individual radial growth model for three oak species (*Q. ariabilis*, *Q. acutissima* and *Q. mongolica*) in mixed and uneven-aged oak stands in central South Korea. The site quality dependent model showed relatively better statistical performance than the site quality independent model.

Different components, such as tree age, tree size, competition and growing site, were employed to build and evaluate the individual radial growth model. The components of age, tree size and competition were proven to have significant effects on the tree radial growth. The age and competition had negative effects on the radial growth, while the tree size had a positive effect. The age effect on the radial growth was greater for larger trees and with lower competition. The tree size was also expected to have a greater effect on younger trees and with lower competition. The competition had greater negative effects on younger and bigger trees. The classical site index, derived from the mean age and dominant height, was proven to not be applicable to the individual radial growth model. Among the topographical factors, only the aspect index was proven to be statistically significant for explaining the radial growth. These analyses of the effects of the different components on the radial growth of oak species were found to be reasonable and consistent with well-known silvicultural experiences.

*Keywords: Individual tree, radial growth model, mixed stand, uneven-aged stand, oak stands*

## 1. Introduction

The management of uneven-aged and mixed stands is of increasing importance for sound and sustainable forest management. Stand level growth models have proven to be useful for managing even-aged and pure stands, and such models are widely used, especially in Korea (Kim 1963; Kim and Lee 1970; Lee 1971; Yoo and Noh 1987; Yoo et al. 1986). However, they are of limited use in uneven-aged and mixed stands, where many different species, ages and sizes of trees exist (Lee et al. 2004). A crucial factor for managing uneven-aged and mixed forests is knowledge relating to growth at the individual tree level for different tree species (Lee et al. 2004). Analyzing the growth of mixed stands, therefore, requires a modeling approach related to the growth of individual trees. Individual tree

1 growth models need to simulate each individual tree as a basic unit (Vanclay 1994; Gadow  
2 and Hui 1999; Bartelink 2000).

3 Most oak stands, which account for approximately 26% of the total forested area in South  
4 Korea (Korea Forest Service 2008), are naturally regenerated (Lee et al. 2004). *Quercus*  
5 *variabilis*, *Quercus acutissima* and *Quercus mongolica* are dominant tree species in oak  
6 stands in South Korea, and are recognized as potential natural species requiring intensive  
7 management. Their growth rates are comparable with other tree species, particularly red  
8 pine (*Pinus densiflora*), but only few individual tree growth models have been developed  
9 for oak species, especially for mixed and uneven-aged stands in South Korea. Lee et al.  
10 (2004) developed a diameter growth model for mixed stands of red pine and oak in Korea.  
11 However, this model dealt with the growth of two different tree species, no models are  
12 currently available for different oak species in Korean forests.

13 This study aimed at developing an individual tree growth model for three oak species  
14 (*Q. variabilis*, *Q. acutissima* and *Q. mongolica*) in mixed and uneven-aged oak stands in  
15 the central region of Korea, which takes the tree age, size, competition and site condition  
16 into consideration.

## 17 18 **2. Materials**

19 A total of 40 temporary plots, covering between 0.01 and 0.08 ha, were installed in a  
20 mixed and uneven-aged oak forest zone (*Q. variabilis*, *Q. acutissima* and *Q. mongolica*),  
21 which represented the core of the natural distribution of oak in central Korea. The stands  
22 were naturally regenerated, without a history of silvicultural treatment. The plots were  
23 selected to cover a large range of tree ages and sizes, as well as competition and topographic  
24 situations. The diameter at the breast height (dbh) and the height of all the trees were  
25 measured in each sample plot. The tree coordinates were established by measuring the  
26 distance and azimuth from the primary subject tree of a sample plot to the neighboring trees.  
27 The increment cores were taken from 5 to 8 sample trees in each plot, with the tree rings  
28 measured to within 1/100 mm using a tree ring measurement system. A total of 217 sample  
29 trees, consisting of 94 *Q. variabilis*, 66 *Q. acutissima* and 57 *Q. mongolica*, were also made  
30 available for this study.

1 In order to avoid measurement errors in the annual tree ring widths, as well as the  
2 influence of short-term climatic fluctuations, the periodic annual increment for the  
3 preceding 10 years was used as the annual radial increment ( $\Delta r$ ) in the analysis. The basic  
4 statistics for the data set are given in Table 1.

5 <Table 1>

## 6 7 **3. Methods**

### 8 **3.1. Radial growth model**

#### 9 **3.1.1 Model components**

10 Individual tree growth has been modeled using either the basal area (Wykoff 1990;  
11 Quicke et al. 1994; Monserud and Sterba 1996; Jögiste 2000; Rose Jr and Lynch 2001) or  
12 diameter increment (Pukkala 1989; Lee 1996; Lee et al. 1999; Gourlet-Fleury and Houllier  
13 2000; Rautiainen et al. 2000; Lee et al. 2004). West (1979) concluded there was no reason  
14 for expressing the growth as either the diameter or basal area increment (Lee et al. 2004).  
15 In this study, the annual radial increment at the breast height ( $\Delta r$ ) was selected as the  
16 dependent variable for the radial growth model. It was also assumed that the annual radial  
17 growth of a tree depended on the tree age and size, as well as the competition and site quality  
18 at the stand level, as suggested by equation 1 in Lee et al. (2004).

$$19 \text{ growth} = f(\text{Age}, \text{Size}, \text{CI}, \text{Site}) \quad (1)$$

20 The tree age has generally been considered an important variable for radial growth, and  
21 was included as an independent variable in our model. For the tree size variable, the dbh  
22 was employed as an independent variable, because assessing the height and crown variables  
23 in the field is expensive and can be associated with a high level of measurement bias (Lee  
24 et al. 2004).

25 In even-aged forests, the effect of the site quality on tree growth is generally accounted  
26 for by the site index, which is derived from the stand age and dominant height (Schröder  
27 2000; Lee et al. 2004). The stand age and dominant height are generally incorporated to  
indirectly consider the effect of the site quality on tree growth (Lee 1996; Lee et al. 1999;

1 Lee et al. 2004). Generally, a simple and unique index for assessing the site quality in  
 2 uneven-aged and mixed stands is unavailable. Therefore, an alternative approach is to  
 3 directly include specific attributes of the growing site (Wykoff 1990; Hasenauer and  
 4 Monserud 1996; Monserud and Sterba 1996; Sterba et al. 2002; Lee et al. 2004). Wykoff  
 5 (1990) used the aspect, slope, elevation, habitat type and geographic location as indicators  
 6 of the site quality for stands composed of mixed species. Hasenauer and Monserud (1996),  
 7 Monserud and Sterba (1996), Sterba et al. (2002), and Lee et al. (2004) also incorporated  
 8 the elevation, slope and aspect into their individual tree growth models for mixed stands. In  
 9 this study, two models were built and evaluated, as follows; 1) a site quality-independent  
 10 model (*SQiD*), and 2) a site quality-dependent model (*SQD*). For the site quality dependent  
 11 model (*SQD-1*), the dominant height and age were initially included as independent  
 12 variables for representing the site index. Topographic variables; the aspect, elevation, and  
 13 slope, were also introduced to explain the site quality in the site quality dependent model  
 14 (*SQD-2*).

15 A competition component, assuming values in the interval [0, 1], was used as an  
 16 influencing factor on the radial growth. In this study, Hegyi's competition index (*HgCI*, eq  
 17 (2)) was applied, which was calculated from the *DBH* ratio and distance between the subject  
 18 tree and competitor trees (Hegyi 1974).

$$HgCI_i = \sum_{j=1}^n \frac{d_j / d_i}{Dist_{ij}} \quad (2)$$

where;

$d_i$  = diameter of the subject tree (cm) at breast height

$d_j$  = diameter of the competitor tree (cm) at breast height

$Dist_{ij}$  = distance between subject tree and competitor tree (m)

19 In this study, an expanded iterative search method was applied to select the competitor  
 20 trees (Figure 1), which consisted of two steps: in the first step, the neighbors to the subject  
 21 tree ( $S_i$ ) were selected as the primary competitor trees ( $PC_j$ ), using the iterative search  
 22 method developed by Lee and Gadaw (1997). This search method differentiates between  
 23 active competitors facing the reference tree and passive competitors positioned behind an  
 24 active competitor, when viewed from the subject tree. In the second step, the neighbors to  
 25 each primary competitor tree ( $PC_j$ ) were selected as secondary competitor trees ( $SC_{jk}$ ),

1 using the same search method.

2 <Figure 1>

3

### 4 **3.1.2 Model Structure**

5 The following radial growth model (eq. 3), suggested by Lee et al. (2004), was selected  
6 for modeling the individual radial growth of three oak species;

$$\Delta r_{t,j} = a_0 A_{t,j}^{a_1} D_{t,j}^{a_2} e^{b \cdot CI_{t,j}^{0.5}} SQ_t \quad (3)$$

7 where  $\Delta r_{t,j}$  is the annual radial increment at the breast height of tree  $j$ , at time  $t$  (cm),  $A_{t,j}$

8 the tree age (years),  $D_{t,j}$  the tree diameter at the breast height (cm),  $CI_{t,j}$  Hegyi's

9 competition index for tree  $j$  at time  $t$ ,  $SQ_t$  the stand level site quality, with  $SQ_t = 1$  for the

10 SQ-independent model (SQiD),  $SQ_t = Ao_t^{c_1} Ho_t^{c_2}$  for the SQ-dependent model with

11 dominant height and age (*SQD-1*),  $Ho$  the stand dominant height (mean height of the 3 to 5  
12 highest trees; m),  $Ao$  the stand dominant age (mean age of the 3 to 5 highest trees; in years),

13  $SQ_t = e^{c_1 AI_t^{0.5} + c_2 EL_t^{0.5} + c_3 SL_t^{0.5}}$  the SQ-dependent model with topographic factors (*SQD-2*),  $AI$  the

14 aspect index; where  $AI = 2 \times |1 - aspect/180|$ , with values ranging from 0 (south) to 2 (north),

15  $EL$  the elevation (m) and  $SL$  the slope ( $^\circ$ ).

16

### 17 **3.2. Statistical performance evaluation**

18 The model was refitted several times with the initial values estimated from the previous

19 fits to ensure stability of the parameter estimates, using the SAS NLIN procedure (SAS

20 Institute 1998). The statistical performance of the model was evaluated using the root of the

21 mean square of error ( $\sqrt{MSE}$ ), the coefficient of determination ( $R^2$ ), Akaike's Information

22 Criteria (*AIC*; Burnham and Anderson 2002) and the significance level of the estimated

23 coefficients

## 4. Results and Discussion

### 4.1. Statistical performance of the models

Table 2 shows the parameter estimates and related statistics for all the models (*SQiD*, *SQD-1* and *SQD-2*). The models showed relatively good performance for explaining the variations in radial growth, with  $R^2$  values ranging from 0.45 to 0.54 for *Q. variabilis*, 0.53 to 0.66 for *Q. acutissima* and 0.47 to 0.59 for *Q. mongolica*. The statistical evaluation of the models using  $\sqrt{MSE}$ ,  $R^2$  and *AIC* indicated that the site quality dependent models were slightly superior to the site quality independent model. Lee et al. (2004) also proved that the dbh models for red pine and oaks could be significantly improved by the addition of site quality variables to the models. Our model employed relatively simple variables for tree size and competition, which can be easily obtained from simple measurements. Therefore, the radial growth model in this study might be less accurate than those requiring information on the tree height or crown to calculate the competition index (Biging and Dobbertin 1995; Pretzsch 1995; Hasenauer and Monserud 1996; Monserud and Sterba 1996; Bachmann 1997), but with relatively simple components this model can be more widely used in situations where information on the tree height and crown are not available.

<Table 2>

### 4.2. Statistical feasibility of the parameters

#### 4.2.1. Tree age

The tree age parameters were found to be significant in all models at a significance level of 0.001 or 0.01. The negative coefficient for tree age indicated that the annual radial growth decreased with increasing tree age when the other variables remain constant, which was consistent with the general experience of other individual tree growth models (Quicke et al. 1994; Lee 1996; Lee et al. 1999; Jögiste 2000; Lee et al. 2004). Figure 2 shows the effects of tree age and size (dbh) on radial growth in the *SQiD* model. The competition index was fixed at a value of 1.5. The three dimensional graphs shown in Fig. 2 are useful for evaluating the combined effects of two independent variables (McFadden and Oliver 1988;

1 Quicke et al. 1994; Jögiste 2000; Lee et al. 2004). The negative effect of tree age on radial  
2 growth becomes increasingly distinct with younger age, bigger size (Figure 2) and lower  
3 competition (Figure 3). For *Q. acutissima*, the negative effect of tree age was noticeably  
4 lower compared to the other two oak species, suggesting *Q. acutissima* should be less  
5 sensitive to the changing age of a tree.

6 <Figure 2>

#### 8 **4.2.2. Tree size**

9 The coefficients of the tree size component, *DBH*, were significant at the relative lower  
10 0.05 or 0.01 level, except for *Q. variabilis* in SQD-1. Lee et al. (2004) also reported a relative  
11 lower significant level for the DBH parameter of *Quercus* species. The DBH coefficients  
12 were positive, with the exception of those for *Q. variabilis* in the *SQD-1* model, as  
13 confirmed in many other studies (Wykoff and Monserud 1988; Wykoff 1990; Quicke et al.  
14 1994; Hasenauer and Monserud 1996; Monserud and Sterba 1996; Lee 1996; Jögiste 2000;  
15 Sterba et al. 2002; Lee et al. 2004). The positive effect of the tree size on the radial growth  
16 can be explained by the size, which reflects the effects of previous competition and/or vigor  
17 on tree growth (Lee et al. 2004). The positive effect of size on radial growth distinctly  
18 appears at a younger tree age (Figure 2) and with a lower competition index (Figure 4). This  
19 implies that larger trees grow faster when there is less competition. The positive effect of  
20 tree size was similar for all oak species with respect to the competition effect.

21 < Figure 3 >

#### 23 **4.2.3. Competition**

24 All competition coefficients were significant at the 0.001 level. The negative coefficients  
25 indicate that the radial growth would be expected to decrease with increasing competition,  
26 as the radial increment was found to decrease with increasing competition in other studies  
27 (Wykoff 1990; Holmes and Reed 1991; Quicke et al. 1994; Biging and Dobbertin 1995;  
28 Hasenauer and Monserud 1996; Lee 1996; Monserud and Sterba 1996; Jögiste 2000; Sterba

1 et al. 2002; Lee et al. 2004). The negative effect of competition on tree growth was larger  
2 with lower competition and age (Figure 3), and bigger trees (Figure 4). This indicates that  
3 younger and bigger trees are more sensitive to competition. The negative effect of  
4 competition was similar for all oak species with respect to the size effect.

5 <Figure 4>

#### 7 **4.2.4. Site Index**

8 The dominant height parameters in the site quality dependent model (*SQD-1*) were only  
9 significant for *Q. variabilis*, while the dominant age parameters were significant for all  
10 species at the 0.001 or 0.05 level. The site index, which can be represented by the dominant  
11 age and height, has been reported as being questionable, as the age and dbh can already  
12 capture the influence of the site on individual tree growth (Lemon and Schumacher 1962;  
13 Cole and Stage 1972, Quicke et al. 1994), and due to the initial suppression of smaller trees  
14 (Peng 2000). Lee et al (2004) also reported a similar problem when using the site index, or  
15 dominant age and height, for the mixed stands of *P. densiflora* and *Q. variabilis*. Other  
16 works have also pointed out that the classical site index approach is not applicable to  
17 uneven-aged multi-species forests (Wykoff and Monserud 1988; Sterba et al. 2002).

#### 19 **4.2.5. Topography**

20 The site quality-dependent model, with a topographic index, had the advantage of  
21 accessing the growing site by employing the same criteria for different tree species (Lee et  
22 al. 2004). The topographic parameters in the site quality dependent model (*SQD-2*) showed  
23 different performances according to the tree species. For *Q. variabilis* and *Q. mongolica*  
24 only the aspect parameter was significant at the 0.001 and 0.05 levels. For *Q. acutissima*,  
25 the aspect and slope were significant at the 0.01 level. It was notable that the aspect  
26 parameters were found to be significant for all three species, which was similar to the result  
27 of Monserud and Sterba (1996), where only the aspect affected oak growth, but different  
28 from that of Lee et al. (2004), which showed a non-significant effect of the aspect on the  
29 radial growth for mixed stands of *P. densiflora* and *Q. variabilis*. It was also notable that

1 the coefficients of elevation were found be insignificant for all species, which was similar  
2 to the result of Hasenauer and Monserud (1996), who reported an insignificant influence of  
3 this site factor on the growths of stone pine and oak in Austria, but different to that of Lee  
4 et al. (2004), where the elevation parameter was significant for *Q. variabilis* at the 0.0001  
5 level. The coefficients of slope for *Q. variabilis* and *Q. mongolica* were insignificant, as  
6 reported by Hasenauer and Monserud (1996) and Lee et al. (2004).

7 The coefficients of the aspect index (*AI*), which increased from south to north, were  
8 positive for *Q. acutissima* and *Q. variabilis* and negative for *Q. mongolica*. This might imply  
9 that the northern slope favored diameter growth of *Q. acutissima* and *Q. variabilis*, but not  
10 for *Q. mongolica*. However, *Q. acutissima* and *Q. variabilis* appear on the southern slope  
11 and *Q. mongolica* on the northern slope, when the slope is identified by the main ridge  
12 stretching from the top of the mountain (Lee et al. 2001; Lee et al. 2003; Ham et al. 2004).  
13 The mean value and range of the *AI* in Table 1 also satisfactorily represent the spatial  
14 distribution of the oak species. The spatial distribution and growth of tree species can be  
15 influenced by the micro-topography which is identified by small branch ridge within the  
16 same aspect by the big main ridge (Lee et al. 2008; Kim et al. 2009). Considering the actual  
17 spatial distribution and *AI* range (Table 1) of oak species in the study area, the positive  
18 effect of *AI* on *Q. acutissima* and *Q. variabilis* can only imply that these species appear  
19 mostly on the southern slope from the main ridge, but their growth could improve with  
20 increasing *AI*, or from south to north formed by the small branch ridge on the southern slope.  
21 Similarly, the negative effect of *AI* on *Q. mongolica* only suggests that the growth of *Q.*  
22 *mongolica*, mostly appearing on the northern slope from the main ridge, can be facilitated  
23 with decreasing *AI*, or from north to south formed by the small branch ridge on the northern  
24 slope.

## 25 26 **5. Conclusion**

27 This study prepared individual radial growth models for different oak species, and  
28 showed that the statistical performance of the model can be significantly improved by  
29 incorporating variables for site quality in terms of site index and topographical factors.  
30 However, the mean age and dominant height, which together represent the site index,

1 showed low levels of significance or non-significance in explaining the radial growth of oak  
2 tree species. Similarly to the site index, the topographical factors appeared to be problematic  
3 for representing the site quality. The coefficients of elevation were not significant for all  
4 oak species, with only the aspect index proven to have a significant effect on radial growth  
5 for all oak species.

6 The individual radial growth model developed in this study employed tree age and size,  
7 competition and growing site as factors influencing radial growth. Relatively easily  
8 available variables in the field, such as tree age, dbh, and Hegey's competition index, which  
9 only uses the dbh and distance, dominant height, aspect, slope and elevation, were used for  
10 the model components. This can be attributed as the reason our radial growth model might  
11 be less accurate than those requiring crown information to represent the competition and  
12 soil information when assessing the site quality. However, this model, with relatively simple  
13 variables, can be more widely used in situations where information on the crown and soil  
14 are not available.

15 The components of age, tree size and competition have been proven to have a significant  
16 effect on the tree radial growth. Age and competition have negative effect on the radial  
17 growth, while the tree size has a positive effect. The age effect on the radial growth was  
18 greater for larger trees and with lower competition. The tree size was also expected to have  
19 a greater effect on younger trees and with lower competition. Competition was found to  
20 have a greater negative effect on younger and bigger trees. These analyses of the effects of  
21 the different components on the radial growth of oak species were found to be reasonable  
22 and consistent with well-known silvicultural experiences.

## 24 **Acknowledgement**

25 This research is supported by the Korea Ministry of Environment under the "Climate  
26 Change Correspondence Program" (Project Number: 2014001310008).

## 28 **References**

- 29 Bachmann, M. 1997. Zum Einfluss von Konkurrenz auf das Einzelbaumwachstum in  
30 Fichten/Tannen/Buchen-Bergwäldern. AFJZ. 168, 127–130  
31 Bartelink, H.H. 2000. A growth model for mixed stands. For. Ecol. Manage. 134, 29-43

- 1 Biging, G.S., Dobbertin, M. 1995. Evaluation of competition indices in individual tree  
2 growth models. *For. Sci.* 41, 360–377
- 3 Burnham, K.P., Anderson, D.R. 2002. Model selection and multimodel inference: A  
4 practical information theoretic approach. Springer
- 5 Cole, M.C., Stage, A.R. 1972. Estimating future diameters of lodgepole pine trees. USDA  
6 For. Serv. Res. Pap. INT-131, 20-25
- 7 Gadow, K.v., Hui, G.Y. 1999. Modelling forest development. Kluwer Academic Publisher.
- 8 Gourlet-Fleury, S., Houllier F. 2000. Modelling diameter increment in a lowland evergreen  
9 rain forest in French Guiana. *For. Ecol. Manage.* 131, 269-289
- 10 Ham, B.Y., Lee, W.K., Chong, J.S., Lee, J.H. 2004. Estimation of Spatial Distribution of  
11 Occurrence Probability of Oak species Using IKONOS satellite imagery and GIS.  
12 *Kor. J. For. Meas.* 7, 74-84 (in Korean with English abstract)
- 13 Hasenauer, H., Monserud, R.A. 1996. A crown ratio model for Austrian forests. *For. Ecol.*  
14 *Manage.* 84, 49-60
- 15 Hegyi, F. 1974. A simulation model for managing jack pine stands. In: Growth models for  
16 tree and stand simulation, Fries, J. (ed.). Royal Coll. Of For., Stockholm, Sweden, 74-  
17 90
- 18 Holmes, M.J. Reed, D.D. 1991. Competition indices for mixed species Northern Hardwoods.  
19 *For. Sci.* 37, 1338–1349
- 20 Jõgiste, K. 2000. A basal area increment model for Norway spruce in mixed stands in  
21 Estonia. *Scand. J. For. Res.* 15, 97-102
- 22 Kim, D.C. 1963. A study of the yield and growth of the red pine (*Pinus densiflora*) produced  
23 in Kangwon province. The research reports of Korean Agriculture Research Institute  
24 6, 71-90 (in Korean)
- 25 Kim, D.C., Lee, H.K. 1970. A study on the yield and growth of the *Quercus* spp. The  
26 research reports of Korean Forestry Research Institute 17: 9-27 (in Korean)
- 27 Kim, T.M., Lee, W.K., Son, Y., Yoo, S., Kim, S.R. 2009. Topographical Analysis for Spatial  
28 Distribution of *Pinus densiflora*. *J. Kor. For. Soc.* 98, 764-771 (in Korean with  
29 English abstract)
- 30 Korea Forest Service. 2008. Statistical Yearbook of Forestry, 2008. Korea Forest Service,  
31 Seoul, 2009, Chapter II, 30-32 (in Korean)

- 1 Lee, H.K. 1971. A study on the yield and growth of the central region pine. The research  
2 reports of Korean Forestry Research Institute 18, 9-30 (in Korean)
- 3 Lee, W.K. 1996. Estimating the competition indices and diameter growth of individual trees  
4 through position-dependent stand survey. J. Kor. For. Soc. 85, 539-551 (in Korean  
5 with English abstract)
- 6 Lee, W.K. Gadow, K.v. 1997. Iterative Bestimmung von Konkurrenzbaeume in *Pinus*  
7 *densiflora* Bestaende (Iterative selection of competitor trees in *Pinus densiflora*  
8 stands). AFJZ 168, 41-45
- 9 Lee, W.K., Seo, J.H, Hwang, J.W., Kim, Z.S. 1999. DBH-growth model by competition  
10 index of *Pinus koraiensis*. Kor. J. For. Meas. 2, 21-30 (in Korean with English abstract)
- 11 Lee, W.K., Lee, J.H., Chung, K.H., Jun, E.J. 2001. Spatial characteristics of forest type  
12 distribution on the basis of geo-morphological factors and IKONOS satellite  
13 imagery. Kor. J. For. Meas. 4, 74-82 (in Korean with English abstract)
- 14 Lee, J.H., Lee, W.K., Jun, E.J., Kim, S.W., Kwak, D.A. 2003. Regional Level Impact of  
15 Global Warming on the Distribution of Oak Stands in Central Korea. Kor. J. Quat.  
16 Res. 17, 135-138 (in Korean with English abstract)
- 17 Lee, W. K., Gadow, K.v., Chung D.J., Lee J. L. 2004. DBH growth model in *Pinus densiflora*  
18 and *Quercus variabilis* mixed stands in central Korea. Ecol. Modell. 176, 187-200
- 19 Lee, W.K., Kim, T.M., Jung, S.E., Choi, H.A. 2008. Optimal zoning for analyzing  
20 ecological and topological characteristics in the process of up-scaling from plot to  
21 regional scale. A3 Foresight Program 2008 Seoul Workshop
- 22 Lemon, P.E. Schumacher, F.X. 1962: Volume and diameter growth of ponderosa pine trees  
23 as influence by site index, density, age, and size. For. Sci. 8, 236-249
- 24 McFadden, G., Oliver, D.C. 1988. Three-dimensional forest growth model relating tree size,  
25 tree number and stand age: relation to previous growth models and to self-thinning.  
26 For. Sci. 34, 662-676
- 27 Monserud, R.A. and Sterba, H., 1996: A basal area increment model for individual trees  
28 growing in even- and uneven-aged forest stands in Australia. For. Ecol. Manage. 80:  
29 57-80.
- 30 Peng, C. 2000. Growth and yield models for uneven-aged stands: past, present and future.

- 1 For. Ecol. Manage. 132, 259-279
- 2 Pretzsch, H. 1995. Perspektiven einer modellorientierten Waldwachstumsforschung. Forstw.  
3 Cbl. 114, 188-209
- 4 Pukkala, T. 1989. Predicting diameter growth in even-aged Scots pine stands with a spatial  
5 and non-spatial model. Silva. Fenn. 23, 101-116
- 6 Quicke, H.E., Meldahl, R.S., Kush, J.S. 1994. Basal area growth of individual trees: A model  
7 derived from a regional longleaf pine growth study. For. Sci. 40, 528-542
- 8 Rautiainen, O., Pukkala, T., Miina, J., 2000, Optimizing the management of even-aged  
9 *Shorea robusta* stands in southern Nepal using individual tree growth models. For.  
10 Ecol. Manage. 12, 417-429
- 11 Rose Jr, C.E., Lynch, T.B. 2001. Estimating parameters for tree basal area growth with a  
12 system of equations and seemingly unrelated regressions. For. Ecol. Manage. 148, 51-  
13 61.
- 14 SAS Institute Inc. 1998. The SAS system for Windows, Release 6.12.
- 15 Schröder, J. 2000. Comparison of a spatial and a non-spatial model for predicting basal area  
16 increment of individual Maritime pine trees in Galicia. In: Schroeder, J. 2000. Analyse  
17 der Wuchsdynamik von *Pinus pinaster* Ait. In Nordwest-Spanien. Dissertation in  
18 Göttingen.
- 19 Sterba, H., Blab, A., Katzensteiner, K. 2002. Adapting an individual tree growth model for  
20 Norway spruce (*Picea abies* L. Karst.) in pure and mixed species stands. For. Ecol.  
21 Manage. 159, 101-110
- 22 Vanclay, J.K. 1994. Modelling forest growth and yield—applications to mixed tropical forests.  
23 CAB International, Wallingford, UK.
- 24 West, P.W. 1979. Use of diameter increment and basal area increment in tree growth studies.  
25 Can. J. For. Res. 10, 71-77
- 26 Wykoff, W.R. 1990. A basal area increment model for individual conifers in northern Rocky  
27 Mountains. For. Sci. 36, 1077-1104
- 28 Wykoff, W.R., Monserud, R.A. 1988. Representing site quality in increment models: a  
29 comparison of methods. In: Ek. A.R., Shifley, S.R., Burk, T.E. (Eds.): Forest Growth  
30 and Modelling and Prediction, vol. 1. USDA For. Serv. Gen. Tech. Rep. NC-120, 184-  
31 191

- 1 Yoo, J.W., Shim, D.S., Noh, K.H., Park, C.W., Lee, H.K., Kim, S.I. 1986. Studies on the  
2 yield and growth of Korean white pine (*Pinus koraiensis*) and Sawtooth oak (*Quercus*  
3 *mongolica*) stand. The research reports of Korean Forestry Research Institute 33, 13-  
4 34
- 5 Yoo, J.W., Noh, K.H. 1987. A study on the yield and growth of Mongolian oak (*Quercus*  
6 *mongolica*) stand. The research reports of Korean Forestry Research Institute 34, 1-  
7 11
- 8

1 <Table list>

2 Table 1. General description of the variables used in the diameter growth model according  
3 to species.

4

5 Table 2. Parameter estimates and related statistics for the radial growth models for oak  
6 species.

7

8 <Figure list>

9 Figure. 1. Diagram of a sample plot using the expanded iterative search method ( $S_i$ : subject  
10 tree  $i$ ,  $PC_j$ :  $j^{th}$  primary competitor of  $S_i$ ,  $SC_{jk}$ :  $k^{th}$  secondary competitor of  $PC_j$ ).

11 Figure. 2. Annual radial growth for different ages and diameters, as simulated using the  
12 *SQiD* model, with a constant CI of 1.5.

13 Figure. 3. Annual radial growth for different combinations of Hegyi's competition index  
14 (*HgCI*) and age, as simulated using the *SQiD* model, with a constant DBH of 15 cm.

15 Figure. 4. Annual radial growth for different combinations of Hegyi's competition index  
16 (*HgCI*) and the current DBH, as simulated using the *SQiD* model, with a constant tree  
17 age of 30 years

## 1 &lt;Tables&gt;

2 Table 1. General description of the variables used in the diameter growth model according to species.

Variables	Total (217)				<i>Q. variabilis</i> (94)				<i>Q. accutissima</i> (66)				<i>Q. mongolica</i> (57)			
	mean	min	Max	S.D.	mean	min	Max	S.D.	mean	min	Max	S.D.	mean	min	Max	S.D.
Age (year)	36.1	11	103	10.3	35.4	17	73	8.1	34.9	11	55	7.5	39.7	19	103	15.2
DBH (cm)	18.5	8.0	35.3	5.9	20.1	8.8	35.3	5.9	17.9	8.5	33.0	5.3	17.8	8.0	29.5	5.5
Height (m)	14.8	6.0	23.0	3.4	15.9	1.7	21.0	3.2	14.4	6.0	23.0	3.6	14.1	6.0	21.0	3.1
HgCI	2.7	0.2	8.8	1.3	2.7	0.8	7.9	1.4	2.5	0.2	5.6	0.9	2.8	1.0	8.8	1.4
PAI (cm)	0.231	0.046	0.521	0.111	0.244	0.052	0.499	0.108	0.227	0.046	0.521	0.124	0.216	0.050	0.459	0.099
Ao (year)	41	20.3	72.3	9.5	35.8	28.3	69.0	7.2	34.9	28.3	43.0	4.1	42.5	20.3	72.3	13.8
Ho (m)	14.3	13.2	20.3	2.5	17.4	13.7	20.3	1.7	23.2	13.7	20.0	1.9	15.9	13.2	20.0	2.2
Aspect (°)	181.4	10	355	82.6	193.7	20	350	49.8	176.5	120	210	27.9	166.7	10	355	144.9
AI	0.91	0.00	1.94	0.4	0.46	0.11	1.89	0.37	0.36	0.00	1.08	0.30	1.27	0.11	1.94	0.55
Elevation (m)	284.9	195	460	53.7	267.4	195	460	35.9	262.9	195	330	29.8	333.2	220	460	60.8
Slope (°)	25.2	7	39	7.9	25.3	7	39	7.8	23.2	7	39	8.1	27.4	15	36	7.4

*HgCI: Hegyi's competition index, PAI: Periodic Annual Increment of tree radius, Ao: dominant stand age,*

*Ho: dominant stand height, AI: Aspect index = 2 |1-Aspect/180|.*

3

4

5

6

1  
2  
3  
4  
5

Table 2. Parameter estimates and related statistics for the radial growth models for oak species.

Parameters	Tree species and growth models								
	<i>Q. variabilis</i>			<i>Q. acutissima</i>			<i>Q. mongolica</i>		
	<i>SQiD</i>	<i>SQD-1</i>	<i>SQD-2</i>	<i>SQiD</i>	<i>SQD-1</i>	<i>SQD-2</i>	<i>SQiD</i>	<i>SQD-1</i>	<i>SQD-2</i>
$a_0$	5.4314*	0.0866*	5.8147*	1.0107*	0.1234*	0.0599*	3.2970*	5.6265*	1.6409*
$a_1$	-0.7901***	-1.1056***	-0.9735***	-0.3042**	-0.4127***	-0.2243**	-0.6376***	-0.5397***	-0.5122**
$a_2$	0.2297**	-0.0109 <sup>ns</sup>	0.3167***	0.2717**	0.2963**	0.1299*	0.2740*	0.4230**	0.2336*
$b$	-0.6449***	-0.8702***	-0.5950***	-0.7825***	-0.7584***	-0.6326***	-0.7265***	-0.6781***	-0.7846***
$c_1$		1.1442***	0.4858***		0.7020***	0.5960**		-0.4225*	-0.2498*
$c_2$		0.7878***	-0.0093 <sup>ns</sup>		-0.0426 <sup>ns</sup>	0.1026 <sup>ns</sup>		-0.0616 <sup>ns</sup>	0.0155 <sup>ns</sup>
$c_3$			0.0135 <sup>ns</sup>			0.1496**			0.0845 <sup>ns</sup>
$\sqrt{MSE}$	0.0814	0.0761	0.0782	0.0876	0.0870	0.0770	0.0694	0.0677	0.0660
$R^2$	0.45	0.54	0.51	0.53	0.55	0.66	0.47	0.54	0.59
$AIC$	-467.75	-478.62	-472.52	-299.21	-298.73	-318.25	-291.36	-314.39	-316.05

\*\*\*: significant at a level of 0.001, \*\*: significant at a level of 0.01, \*: significant at a level of 0.05,  
<sup>ns</sup>: not-significant,  $\sqrt{MSE}$ : the root of mean square of error,  $R^2$ : coefficient of determination,  $AIC$ : Akaike's Information Criteria.

1 <Figures>




Figure. 1. Diagram of a sample plot using the expanded iterative search method ( $S_i$ : subject tree  $i$ ,  $PC_j$ :  $j^{\text{th}}$  primary competitor of  $S_i$ ,  $SC_{jk}$ :  $k^{\text{th}}$  secondary competitor of  $PC_j$ ).




Figure. 2. Annual radial growth for different ages and diameters, as simulated using the SQiD model, with a constant CI of 1.5.

1




Figure. 3. Annual radial growth for different combinations of Hegyi`s competition index (HgCI) and age, as simulated using the SQiD model, with a constant DBH of 15 cm

2




Figure. 4. Annual radial growth for different combinations of Hegyi`s competition index (HgCI) and the current DBH, as simulated using the SQiD model, with a constant tree age of 30 years.

3