

Uncertainties in emissions estimates of greenhouse gases and air pollutants in India and their impacts on regional air quality

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2017 Environ. Res. Lett. 12 065002

(<http://iopscience.iop.org/1748-9326/12/6/065002>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 147.125.57.21

This content was downloaded on 13/06/2017 at 08:38

Please note that [terms and conditions apply](#).

You may also be interested in:

[Explosive growth in African combustion emissions from 2005 to 2030](#)

C Liousse, E Assamoi, P Criqui et al.

[The last decade of global anthropogenic sulfur dioxide: 2000–2011 emissions](#)

Z Klimont, S J Smith and J Cofala

[Exceedances of air quality standard level of PM_{2.5} in Japan caused by Siberian wildfires](#)

Kohei Ikeda and Hiroshi Tanimoto

[Decadal-scale trends in regional aerosol particle properties and their linkage to emission changes](#)

Bin Zhao, Jonathan H Jiang, Yu Gu et al.

[Rising critical emission of air pollutants from renewable biomass based cogeneration from the sugar industry in India](#)

S K Sahu, T Ohara, G Beig et al.

[Carbon reductions and health co-benefits from US residential energy efficiency measures](#)

Jonathan I Levy, May K Woo, Stefani L Penn et al.

[The impact of European legislative and technology measures to reduce air pollutants on air quality, human health and climate](#)

S T Turnock, E W Butt, T B Richardson et al.

[Air pollution co-benefits of low carbon policies in road transport: a sub-national assessment for India](#)

Shivika Mittal, Tatsuya Hanaoka, Priyadarshi R Shukla et al.

Environmental Research Letters



LETTER

Uncertainties in emissions estimates of greenhouse gases and air pollutants in India and their impacts on regional air quality

OPEN ACCESS

RECEIVED

29 May 2016

REVISED

2 April 2017

ACCEPTED FOR PUBLICATION

12 April 2017

PUBLISHED

26 May 2017

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Eri Saikawa¹, Marcus Trail^{1,2}, Min Zhong¹, Qianru Wu¹, Cindy L Young^{1,3}, Greet Janssens-Maenhout^{4,5}, Zbigniew Klimont⁶, Fabian Wagner^{6,7,8}, Jun-ichi Kurokawa⁹, Ajay Singh Nagpure¹⁰ and Bhola Ram Gurjar¹¹

¹ Department of Environmental Sciences, Emory University, 400 Dowman Drive, Atlanta, GA, 30319, United States of America

² Georgia Environmental Protection Division, 2 Martin Luther King Jr. Drive, Atlanta, GA, 30334, United States of America

³ School of Earth and Atmospheric Sciences, Georgia Institute of Technology, 311 Ferst Drive, Atlanta, GA, 30332, United States of America

⁴ European Commission, Joint Research Centre, Institute for Environment and Sustainability, Via Fermi, 2749, 21027 Ispra VA, Italy

⁵ Ghent University, Campus Ardoyen, Ghent-Zwijnaarde, Belgium

⁶ International Institute for Applied Systems Analysis, Schloßplatz 1, 2361 Laxenburg, Austria

⁷ Woodrow Wilson School of Public and International Affairs, Princeton University, Princeton, NJ, 08544, United States of America

⁸ Andlinger Center for Energy and the Environment, Princeton University, Princeton, NJ, 08542, United States of America

⁹ Asia Center for Air Pollution Research, 1182 Sowa, Nishi-ku, Niigata, Niigata, 950-2144, Japan

¹⁰ Humphrey School of Public Affairs, University of Minnesota, 301 19th Avenue S, Minneapolis, MN, 55455, United States of America

¹¹ Department of Civil Engineering, Indian Institute of Technology Roorkee, Roorkee 247667, Uttarakhand, India

E-mail: eri.saikawa@emory.edu

Keywords: emissions, India, air pollution, carbon dioxide, inventories

Supplementary material for this article is available [online](#)

Abstract

Greenhouse gas and air pollutant precursor emissions have been increasing rapidly in India. Large uncertainties exist in emissions inventories and quantification of their uncertainties is essential for better understanding of the linkages among emissions and air quality, climate, and health. We use Monte Carlo methods to assess the uncertainties of the existing carbon dioxide (CO₂), carbon monoxide (CO), sulfur dioxide (SO₂), nitrogen oxides (NO_x), and particulate matter (PM) emission estimates from four source sectors for India. We also assess differences in the existing emissions estimates within the nine subnational regions. We find large uncertainties, higher than the current estimates for all species other than CO, when all the existing emissions estimates are combined. We further assess the impact of these differences in emissions on air quality using a chemical transport model. More efforts are needed to constrain emissions, especially in the Indo-Gangetic Plain, where not only the emissions differences are high but also the simulated concentrations using different inventories. Our study highlights the importance of constraining SO₂, NO_x, and NH₃ emissions for secondary PM concentrations.

1. Introduction

Understanding the spatial and temporal distribution of greenhouse gases (GHGs) and air pollutant precursor emissions is vital to the implementation of appropriate climate and air quality mitigation measures. Obtaining accurate anthropogenic emissions estimates is especially critical in India, which is currently the world's third largest emitter of carbon dioxide (CO₂) from fossil fuel combustion and from industrial processes (production of cement, metals

and chemicals) behind China and USA (Olivier *et al* 2015), after India surpassed Russia in 2010. Half of the top twenty most polluted cities in the world are in India (WHO 2016) and India is ranked the third worst of 180 countries for PM_{2.5} (particulate matter with an aerodynamic diameter less than 2.5 μm) exposure (Hsu *et al* 2016). Several emissions inventories have been developed recently either specifically for India or for larger regions that include India (Streets *et al* 2003, Garg *et al* 2006, Ohara *et al* 2007, Klimont *et al* 2009, 2013, 2016, Zhang *et al* 2009, EC-JRC/PBL 2011,

Lu *et al* 2011, Smith *et al* 2011, Sahu *et al* 2012, Kurokawa *et al* 2013, Pandey *et al* 2014, Sadavarte and Venkataraman 2014, IEA 2014, Janssens-Maenhout *et al* 2015). These inventories are being used in the air quality and climate model simulations to better understand air pollution and climate change in India, in Asia, and globally.

Several studies have compared emissions inventories in India and other Asian countries (Garg *et al* 2006, Granier *et al* 2011, Kurokawa *et al* 2013, Klimont *et al* 2016). However, most studies focus their analysis on similarities and differences in national total emissions and do not analyze regional scale or source sector level emissions. These studies also do not include the recent national emissions inventories. Pandey *et al* (2014) and Sadavarte and Venkataraman (2014) evaluated their Indian emissions inventory against several but not against many others, including the Regional Emissions inventory in ASia version 2.1 (REAS) and Emissions Database for Global Atmospheric Research version 4.2 (EDGAR), the two most commonly used datasets for Asian and global emissions, respectively.

In this study, we first compare emission inventories of anthropogenic, combustion-related surface emissions of CO₂ and air pollutant precursors (carbon monoxide CO, sulfur dioxide SO₂, nitrogen oxides (NO_x = NO + NO₂), and particulate matter with an aerodynamic diameter less than 10 μm, PM₁₀) for each of the four source sectors and compare existing estimates within subnational regions in India. We then use Monte Carlo sampling to assess the uncertainties of the existing emissions estimates per sector and per species. We further conduct a chemical transport model simulation with two different gridded emissions to validate them and assess their impacts on air quality.

2. Methodology

2.1. Datasets

This work relies on existing emissions inventories that provide estimates at provincial and/or national levels in India. We use the following eight inventories to compare and to assess uncertainties at the source sector level, using Monte Carlo sampling methods: 1) EDGAR v4.2 (EDGAR) (EC-JRC/PBL 2011); 2) REAS v2.1 (REAS) (Kurokawa *et al* 2013); 3) INTEX-B (Zhang *et al* 2009); 4) National emissions inventory for residential and road transportation emissions (Nagpure–Gurjar) (Nagpure and Gurjar 2012); 5) National emissions inventory for all sectors (Sadavarte–Venkataraman) (Pandey *et al* 2014, Sadavarte and Venkataraman 2014); 6) IEA (2014), 7) GAINS (Amann *et al* 2011, Klimont *et al* 2016); and 8) Lu *et al* (2011). The emissions inventories analyzed in this paper were developed using a similar bottom-up methodology, where emissions were calculated as the

product of activity data, such as fuel consumption, and fuel- and technology-dependent emission factors. Table 1 describes the details of each inventory including years, source sectors, and species covered, its horizontal resolution, proxies used, as well as its coverage. In addition to these eight that provide sector-level emissions, we also use total CO₂, CO, SO₂ and NO_x emissions estimates by Garg *et al* (2006), total SO₂ emissions estimates by Smith *et al* (2011) and Klimont *et al* (2013), as well as the total NO_x emissions estimates by Ghude *et al* (2012).

2.2. Emissions comparison

We compared emissions of CO₂, CO, SO₂, NO_x, and PM₁₀ from four source sectors, including power plants (power), industrial combustion and processes (industry), domestic combustion (domestic), and road and non-road transportation (transport) for the years between 2000 and 2010 at national and nine subnational regions. Table S1 available at stacks.iop.org/ERL/12/065002/mmedia describes how we categorize sources for each inventory into the four described above. For the national level analysis, we used seven inventories (EDGAR, REAS, Nagpure–Gurjar, Sadavarte–Venkataraman, IEA, GAINS, and Garg) for CO₂, seven inventories (EDGAR, REAS, INTEX-B, Nagpure–Gurjar, Sadavarte–Venkataraman, GAINS, and Garg) for CO and NO_x, seven inventories (EDGAR, REAS, INTEX-B, Nagpure–Gurjar, Sadavarte–Venkataraman, GAINS, and Lu) for SO₂, and four inventories (EDGAR, REAS, INTEX-B, and GAINS) for PM₁₀. For the subnational level analysis, we used five inventories (EDGAR, REAS, INTEX-B, Nagpure–Gurjar, and GAINS).

We also conducted one million Monte Carlo samplings, choosing an emissions inventory per sector per year randomly. We sampled a normal distribution for most but a few emission estimates in Sadavarte–Venkataraman inventory (CO all sectors; SO₂ transport and domestic; and NO_x all sectors), which assumed a log-normal distribution. We used the standard deviation (SD) values reported by each inventory if such information was available. For the inventories where this was not provided, the relative uncertainty estimates provided by REAS for each sector were used. The mean and the SD of the newly-composed emissions estimates were then calculated from one million samples. The uncertainty is reported as the 95% confidence interval, following Kurokawa *et al* (2013).

2.3. Air quality simulation

In order to assess how differences in emissions inputs affect air quality simulations, we conducted two simulations using the Weather Research and Forecasting model coupled with Chemistry (WRF-Chem) version 3.5 (Grell *et al* 2005), which has been validated previously in Asia (Zhong *et al* 2016). We chose two gridded emissions inventories, EDGAR

Table 1. Description of emissions inventories used for this study.

	Years	Source Sectors	Species	Horizontal Resolution	Proxies for allocating emissions	Coverage	Reference
EDGAR	1970–2008	energy, industrial processes, product use, agriculture, large scale biomass burning, and other anthropogenic sources	CO ₂ , N ₂ O, SO ₂ , CO, PM ₁₀ , NO _x	0.1° × 0.1°	Rural, urban, and total populations, roads, railways, power plant maps, and selected industrial production	Global	EC-JRC/PBL 2011
REAS v2	2000–2008	power plants, combustible and non-combustible sources in industry, on-road and off-road sources in transportation, residential and agricultural sources	CO ₂ , N ₂ O, SO ₂ , CO, PM ₁₀ , PM _{2.5} , NO _x , BC, OC, NH ₃ , NMVOC	0.25° × 0.25°	Rural, urban, and total populations, as well as road network	33 Asian countries	Kurokawa <i>et al</i> 2013
INTEX-B	2006	power plants, industry, residential, and transportation	SO ₂ , CO, PM ₁₀ , PM _{2.5} , NO _x , BC, OC, NMVOC	0.5° by 0.5°	spatial proxies at 1km × 1km resolution	22 Asian countries	Zhang <i>et al</i> 2009
Nagpure–Gurjar	2001–2011	residential and road-transport	CO ₂ , N ₂ O, SO ₂ , CO, PM ₁₀ , NO _x	provincial level	NA	India	
Sadavarte–Venkataraman	1996–2015 (projected using 2010 data)	industry, transportation, residential, and ‘informal industries’ including brick production and processing operations for food and agricultural products	CO ₂ , N ₂ O, SO ₂ , CO, PM _{2.5} , NO _x	25 × 25 km (0.25° × 0.25°)	Census data, urban population, road network, and district-level production data	India	Pandey <i>et al</i> 2014, Sadavarte and Venkataraman 2014
IEA	1960–2012	fossil fuel combustion	CO ₂	national level	NA	Global	IEA, 2014
GAINS	1990–2030 (5 year intervals, projection starting in 2015)	energy, domestic, industrial combustion and processes, road and non-road transportation and agriculture	CO, NO _x , PM ₁₀ , PM _{2.5} , CO ₂ , SO ₂	0.5° × 0.5°	RCP and GEA (Global Energy Assessment) sectoral proxies, population, and selected industrial plants location (e.g. smelters)	Global	Amann <i>et al</i> 2011, Klimont <i>et al</i> 2016 www.iiasa.ac.at/web/home/research/researchPrograms/air/Global_emissions.html

and REAS, and performed simulations in January and July of 2008. These two months were chosen to simulate the seasonality in air pollutant concentrations. January is the dry winter month with mostly high air pollutant concentrations, whereas July is in the middle of the monsoon season with relatively low concentrations (Gaur *et al* 2014). For O₃, it is similar and July tends to have higher mixing ratio than in January. The model covered the entire India, with a horizontal resolution of 20 × 20 km and 31 vertical levels. The initial and lateral chemical boundary conditions were taken from a present-day simulation of the NOAA Geophysical Fluid Dynamics Laboratory (GFDL) global chemistry-climate model AM3

(Naik *et al* 2013). The meteorological data were obtained from the National Center for Environmental Prediction (NCEP) Global Forecast System final gridded analysis datasets. We used Carbon-Bond Mechanism version Z (CBMZ) (Zaveri and Peters 1999) for gas-phase chemistry and the Model for Simulating Aerosol Interactions and Chemistry (MOSAIC) (Zaveri *et al* 2008) for aerosol chemistry. The model simulation was spun-up for ten days before the beginning of each monthly simulation.

2.4. Surface observations

We compared our air quality simulations with existing instantaneous surface observations of SO₂ and NO_x




Figure 1. Time series of mean emissions and one standard deviation (green shade) from Monte Carlo results and mean and one standard deviation of different emission inventory estimates for Indian national CO₂, CO, SO₂, NO_x and PM₁₀. Different inventory is represented by a different style line or point. Sadavarte–Venkataraman emissions are represented by S–V and Nagpure–Gurjar emissions are represented by N–G in the legends.

concentrations (Maharashtra Pollution Control Board 2016), and monthly average CO and O₃ mixing ratio (Kumar *et al* 2012) for January and July 2008. In addition, we used Aerosol Optical Depth (AOD) retrieved from the MODerate Resolution Imaging Spectrometer (MODIS) instrument aboard the Terra satellite and compared it with the simulated AOD from WRF-Chem. MODIS provides AOD retrievals at a resolution of 10 × 10 km. In this study, we use Level 2 and Collection 6 aerosol optical thickness at 550 nm. We calculated the monthly mean AOD values from daily observations of AOD over land and ocean.

3. Results

We present the mean and the SD of our Monte Carlo samplings for the national total emissions estimates for the five species in figure 1, as well as the estimates and SD provided by existing emissions inventories. The detailed means and SD of each sector per species from the Monte Carlo simulations are listed in table S2.

3.1. CO₂

Our Monte Carlo analysis illustrates that there is up to ±76% uncertainty in India's national total CO₂ emissions estimates when we combine all the existing estimates equally. This is larger than the REAS' existing uncertainty of ±49% (figure 1). The largest uncertainty comes from the domestic sector, with the maximum being ±149%. The IEA domestic CO₂ emissions are the lowest of all the inventories at 63 Tg CO₂ yr⁻¹ in 2000, contributing to only 13% of REAS domestic CO₂ emissions. This large difference can be explained in part because IEA does not include biofuel emissions in their estimates, similar to GAINS and Garg.

The smallest uncertainty stems from the power sector, which is the primary source of CO₂ emissions in India. EDGAR, REAS, GAINS, Sadavarte–Venkataraman, and IEA power sector CO₂ emissions continuously increase over time, with Sadavarte–Venkataraman emissions being the highest of all at all times.

Using EDGAR, REAS, GAINS, and Nagpure–Gurjar, we further compared the emissions in the nine




Figure 2. Map of India regions and 2005 emissions of CO₂, CO, SO₂, NO_x, and PM₁₀ from EDGAR, REAS, GAINS, and Nagpure–Gurjar datasets from industrial (ind), transport (tra), power (pow), domestic (dom) sectors. Provinces included in each region are listed in supplementary table S4.

subnational regions in 2005 (figure 2). The largest difference of 75% SD in the domestic sector exists in region 8 on the Indo-Gangetic Plain (IGP). This is also the region with the largest difference for the regional total CO₂ (47% SD) and for transport (44%) among EDGAR, REAS, and GAINS. Another large difference of 73% SD in industry come from region 1, which is also on the IGP, and the largest difference for the power sector is found in region 3 (44%), where the greatest number of large (> 2000 MW) power plants exist.

3.2. CO

The majority of CO emitted in India stems from the domestic sector, making up 69%–70%, 49%–56%, 76%–79%, and 64%–77% of total EDGAR, REAS, GAINS, and Sadavarte–Venkataraman, respectively.

The uncertainty of domestic sector CO emissions from the Monte Carlo simulations is on average ±195% and is the largest of all sectors. This value is equivalent to the ±192% uncertainty provided by the REAS inventory for the same sector. On a regional scale, we find the largest differences in domestic emissions within the region 2 on IGP in 2005 (figure 2) being 4.4 Tg yr⁻¹ between GAINS and Nagpure–Gurjar.

The uncertainty of industrial CO emissions from the Monte Carlo samplings is the second largest among the four sectors with ±154% and is larger than the ±118% uncertainty by REAS. REAS industrial CO emissions are consistently high in all nine regions with the largest difference of 2.3 Tg yr⁻¹ (55% and 310% larger than EDGAR and GAINS, respectively) occurring in region 2 on the IGP.

While transport is not as large of a CO source as the domestic and industry sectors, the difference between the REAS and EDGAR estimates has grown from 521 Gg yr⁻¹ (within 10%) in 2000 to 6278 Gg yr⁻¹ by 2008 with REAS emissions over three times larger than those of EDGAR. The difference among the four inventories in the transport sector is most apparent in regions 2, 4, and 7 with larger than 50% SD. It is most likely that Nagpure–Gurjar until 2008 and REAS assumed much larger emissions either from super-emitters or significantly lower penetration of vehicles equipped with catalytic converters; the comparison of NO_x emissions in transport also support this.

3.3. SO₂

Fossil fuel combustion in the power sector contributes the largest portion of SO₂ emissions in India but has the least uncertainty of ±46%. Power plant SO₂ emissions make up 61%–63% of total SO₂ in EDGAR, 47%–50% of REAS, 54%–57% of GAINS, and 70%–75% of Sadavarte–Venkataraman. Although the relative contributions of the power sector to total India SO₂ emissions for EDGAR and REAS differ, power sector emissions agree well for the two inventories between 2000 and 2008. Power plant emissions are within 10% of each other and both increase by 46%–47% from 2000 to 2008.

For SO₂ emissions, our Monte Carlo analysis shows the largest ±99% uncertainty, on average, in the domestic sector. Industry and transport share ±69 and ±66% uncertainty, respectively, on average. Because of the large differences among the inventory estimates, the national SO₂ emissions have an average uncertainty of ±48% over time, which is higher than any of the individual inventory estimates (figure 1). These values are also higher than the calculated uncertainty of ±16% in Streets *et al* (2003) for Asian SO₂ emissions.

The difference is apparent also on a subnational scale in every region (figure 2). In region 6, where the REAS total SO₂ emissions are the highest, their industrial SO₂ are 14.5 times that of EDGAR. Although the total magnitude is not as large as in region 6, region 1 on the IGP has the highest difference (132% SD) in industrial SO₂ with REAS estimating 34 (six) times greater than EDGAR (GAINS). Region 1 also has the highest difference for the total SO₂ with 108% SD. Besides region 7, where the power sector difference is the largest with 63%, REAS total SO₂ emissions are always the highest, followed by GAINS and EDGAR.

3.4. NO_x

Transport NO_x emissions differ substantially among inventories and while the transport sector is the largest NO_x source in REAS and GAINS, it is the second largest in EDGAR and Sadavarte–Venkataraman. The average uncertainty from the Monte Carlo methods for the transport NO_x emissions is the second highest of all sectors (118%). This large difference confirms


that part of the difference is due to the assumptions about the fleet characteristics (i.e., share of super-emitting vehicles and penetration of vehicles with catalytic converters) assumed in each inventory. EDGAR, without emissions from super-emitting vehicles, has the second lowest transport emissions throughout the years. Excluding the emissions from super-emitting vehicles in GAINS, the difference between GAINS and EDGAR transport NO_x emissions is reduced to 28% from 49% with super-emitting vehicle emissions. For CO, super-emitters are mainly gasoline vehicles but for NO_x, they can be either gasoline or diesel. We therefore find even larger uncertainties in transport NO_x emissions than in CO. Within the same region, the ratio of the transport NO_x emissions among the four inventories remaining the same as that of transport CO emissions provides further confirmation of this assumption. In all regions, REAS transport emissions are at least three times as large as those of EDGAR, confirming larger emissions either from super-emitters and/or lower penetration of control technology, similarly to CO.

The largest NO_x source sector in India in EDGAR and Sadavarte–Venkataraman and the second largest in REAS and GAINS is power generation. The trend is the same in all inventories and the only noticeable difference is that GAINS and Sadavarte–Venkataraman (REAS) have approximately 1 (0.3) Tg/year difference from EDGAR. This sector has the second best agreement among the inventories and the average uncertainty is ±83%. The best agreement is seen in the industrial emissions, where the uncertainty from the Monte Carlo simulation is, on average, ±65%. Both of these are, however, still larger than the uncertainties in the existing inventories.

NO_x domestic emissions show the largest uncertainty of ±196% from the Monte Carlo samplings, much higher than the ±37% in Streets *et al* (2003) for Asian NO_x emissions (2003). In all regions, it is clear that EDGAR and GAINS have similarly low domestic emissions, with much higher REAS and Nagpure–Gurjar emissions. The average SD among the four inventories is the second highest with 31% and region 1 again has the largest difference for the domestic sector (80%) as well as for total emissions (89%).

3.5. PM₁₀

Primary PM emissions typically include black carbon (BC), organic carbon (OC), metals, and dust; composition varies strongly between sources. Ambient PM composition includes, beyond primary components, also a variety of secondary compounds, such as sulfates, nitrates, ammonium, metals and other organic and inorganic compounds. REAS and EDGAR emissions of PM₁₀ in India do not show good agreement in any of the sectors explored in this paper. Indeed, uncertainty in all sectors from the Monte Carlo simulation exceeds ±120%. PM₁₀ emissions from the power sector in EDGAR are over three times larger than those of REAS




and GAINS and six times larger than those of Sadaverte–Venkataraman estimates. It is not only the magnitude of emissions from power sector that is different among the inventories, but the inventories also have a varying emissions trend. While EDGAR, REAS, and Sadaverte–Venkataraman show a 44, 58, and 45% increase in power sector emissions over time, respectively, GAINS estimates an 18% decrease. Since power capacity has grown in the considered period, different trends are likely due to the assumptions on penetration and efficiency of control equipment (electrostatic precipitators, fabric filters) on newly build power plants.

Domestic sector is the largest source sector for PM₁₀ in REAS, GAINS, and Sadaverte–Venkataraman and is the second largest in EDGAR. This is also the sector with highest uncertainty of, on average, $\pm 315\%$ in the Monte Carlo samplings. EDGAR domestic sector emissions are more than twice (50%) as large as those of REAS (Sadaverte–Venkataraman). However, unlike the varying growth rate shown for the power sector emissions, the trend over time is consistent among all inventories. Industry sector is the second largest in GAINS and Sadaverte–Venkataraman and the third largest in EDGAR and REAS. The uncertainty of, on average, $\pm 123\%$ from

Table 2. Regional difference of monthly mean (a) emissions between EDGAR and REAS and (b) concentrations between simulations using EDGAR and REAS. Results are shown with deviations of REAS from EDGAR in January and July of 2008.

Regions	(a)									
	2008 January (REAS-EDGAR)/EDGAR (%)					2008 July (REAS-EDGAR)/EDGAR (%)				
	PM ₁₀	CO	SO ₂	NH ₃	NO _x	PM ₁₀	CO	SO ₂	NH ₃	NO _x
1	-14	28	238	96	94	-16	26	242	95	123
2	-7	16	19	185	52	-10	15	12	185	70
3	-70	47	-53	77	-36	-70	47	-56	77	-35
4	-50	4	29	131	95	-52	4	19	131	99
5	-52	-4	33	101	63	-54	-5	27	101	64
6	-62	3	13	85	-7	-64	2	1	85	-2
7	-38	19	15	70	76	-40	16	1	70	105
8	-24	4	-5	163	17	-29	0	-11	162	53
9	-29	44	86	175	100	-46	9	75	168	117

Regions	(b)															
	2008 January (REAS-EDGAR)/EDGAR (%)									2008 July (REAS-EDGAR)/EDGAR (%)						
	PM ₁₀	NO ₃	NH ₄	SO ₄	O ₃	CO	SO ₂	NO ₂	PM ₁₀	NO ₃	NH ₄	SO ₄	O ₃	CO	SO ₂	NO ₂
1	45	140	85	36	9	7	122	84	24	73	52	12	20	3	193	96
2	55	107	74	23	11	13	21	42	21	96	60	6	14	6	23	59
3	5	109	46	0.4	10	17	-30	-24	-18	71	46	0	14	10	-38	-11
4	19	148	60	5	11	7	6	44	-5	89	86	23	20	2	17	69
5	19	203	60	14	11	4	10	33	-16	42	53	6	9	-2	4	36
6	38	124	80	32	12	11	18	-3	-11	33	30	-7	7	-2	-11	-2
7	72	152	123	67	10	20	50	72	11	87	61	5	20	5	1	70
8	43	95	75	32	11	13	-1	13	23	104	66	12	18	2	-16	36
9	52	91	82	55	2	20	85	100	43	152	108	55	17	4	67	120



this sector in the Monte Carlo simulation is the smallest for PM₁₀.

On the subnational scale, REAS total PM₁₀ emissions are the lowest among the three inventories in all regions (figure 2). EDGAR PM₁₀ emissions from the industrial and transport sector are, however, constantly lower than REAS or GAINS in every region, with national average EDGAR just 6% of REAS transport PM₁₀ in 2008. Transport sector has

more than 100% SD in all years, except 2000 and 2005. The main reason for this large difference is due to the lack of super-emitters in EDGAR as mentioned earlier and there is good agreement between REAS and GAINS transport PM₁₀ emissions. However, even after omitting super-emitter contribution, GAINS transport emissions are still 450% higher than EDGAR emissions, which indicates that there are also other differences in emission factors and




Figure 4. (Continue)

technology trends among inventories and possibly the way non-exhaust emissions (road, tire, and break wear) are represented.

3.6. Emission impacts on air quality

Figure 3 compares the simulated monthly mean surface CO, SO₂, NO₂, and O₃ mixing ratios, as well as PM₁₀ concentrations in January and July 2008, using

the REAS and EDGAR emissions inventories. Table 2 highlights the differences in emissions between the two inventories and in simulated concentrations by region. We compare these two simulations with existing observations of CO and O₃ mixing ratios, SO₂ and NO_x concentrations, and AOD in figure 4. The simulation with REAS emissions leads to higher CO but the regional difference is less than 20% in most

regions. The model always underestimates but REAS does a better job in reproducing them. For O_3 , the simulation using REAS produces a slightly higher mixing ratio than EDGAR in all regions, but the difference is often less than 12% (7 ppbv). The model usually overestimates and EDGAR thus reproduces better, except for one site in Ananatpur.

Although the two simulations show similar spatial distributions of NO_2 and SO_2 surface concentrations, their magnitudes differ substantially. The difference is particularly apparent in regions 1 and 9, where the simulation using REAS emissions estimates 36%–55% higher SO_2 and 84%–100% higher NO_2 due to 86%–238% and 95%–100% higher emissions, respectively. The comparison of NO_x and SO_2 concentrations with surface measurements indicate that the model overestimates in cities but usually underestimates NO_x in rural areas.

The REAS model simulation reproduces SO_2 better in general, and NO_x better in most non-urban regions, while the EDGAR model simulation reproduces O_3 better. There is also significant overestimation of NO_x in urban areas and significant underestimation in rural areas using both inventories. These suggest first that transport emissions are possibly overestimated in REAS, as transport sector emissions are mainly from the urban centers. We find that India's O_3 is largely sensitive to NO_x (table 2), as found in Sharma *et al* (2016), and considering that REAS transport emissions were estimated high for all species affirms this hypothesis. Second, the spatial distribution of the different proxies (i.e. population) used by the two inventories may be the cause of the differences in gridded emissions, as reflected in the model simulations.

The simulation using REAS produces 19%–72% (6 – $24 \mu\text{g m}^{-3}$) higher surface concentrations of PM_{10} in most regions than those using EDGAR, even though the primary PM_{10} emissions is lower in REAS. The highest difference occurs in region 7 (72%) and we find that this is due to over 150, 120, and 65% differences in nitrate, ammonium, and sulfate aerosols in the simulation using REAS emissions. SO_2 , NO_x , and NH_3 emissions in most regions are much greater in REAS as compared to those in EDGAR (table 2). Larger emissions of these species lead to a greater production of secondary inorganic aerosols in the model, and consequently, to higher PM_{10} concentrations despite the lower primary PM_{10} emissions. Spatial and temporal emissions of SO_2 , NO_x , and NH_3 play an important role in secondary PM formation, as found in EMEP (2016). We further find that REAS reproduces the MODIS AOD better than when using EDGAR emissions in region 7, confirming the importance of these SO_2 , NO_x , and NH_3 emissions on PM concentrations.

Differences in emissions estimates affect concentrations and mixing ratios of pollutants in varying ways, depending on a region. Emission differences are

large in regions 1, 2, 3 and 9 but we find the largest difference in concentrations and mixing ratios of various pollutants in region 7 in January. In July, the largest differences in air quality and emissions for most species are both in regions 1, 3 and 9. We find that this is due to pollution transport from neighboring regions and to seasonal changes in prevailing wind patterns and precipitation. Constraining the emissions in IGP (regions 1, 2, 7, 8, and 9) through field emission measurement campaigns and enhanced surface measurement network, therefore, will be most useful for the development of air pollution mitigation strategies in India.

4. Conclusions

In this study, various inventories of anthropogenic CO_2 and air pollutant precursor emissions are compared in India on national, regional, and sectoral scales. For the global CO , SO_2 , and NO_x emissions, the difference between inventories is 28%, 42%, and 17% in 2000, respectively (Granier *et al* 2011). In India, we find the differences of total CO , SO_2 , and NO_x emissions among existing inventories to be 15%, 62%, and 58% in 2000. However, our analysis of regional and sector-specific emissions revealed that much higher differences exist at the sector level than the national total emissions. Our Monte Carlo results also highlight higher uncertainties in existing emissions estimates both at the national total and source sector levels than currently considered for most species. Although we can infer whether some differences result from activity data or emission factors, in order to fully understand the reasons behind discrepancies among the multiple inventories, data needs to be transparent and available. Our model results on the impact of emissions uncertainties highlights the importance of constraining emissions through field emission measurement campaigns and enhanced surface measurement network in the IGP region for understanding the local and regional air quality. Our study further highlights the importance of constraining SO_2 , NO_x , and NH_3 emissions for secondary PM formation.

Acknowledgments

We thank Vaishali Naik and Larry W. Horowitz for providing global AM3 simulation. We also thank the editor and three anonymous reviewers for constructive comments to improve the paper. This study was partly supported by the National Science Foundation (grant number AGS-1350021). Bhola R. Gurjar thanks the Max Planck Society (Germany) for the support received through Max Planck Partner Group of Max Planck Institute for Chemistry, Mainz (Germany) and Indian Institute of Technology Roorkee (India). REAS is supported by the Global Environmental Research Fund

of the Ministry of the Environment Japan (S-7 and S-12). The NCEP GFS data used for this study are from the Research Data Archive (RDA), which is maintained by the Computational and Information Systems Laboratory (CISL) at the National Center for Atmospheric Research (NCAR). The data are available at <http://rda.ucar.edu/datasets/ds083.2/> (National Centers for Environmental Prediction, National Weather Service, NOAA, US Department of Commerce, 2000). We would like to acknowledge high-performance computing support from Yellowstone (ark:/85065/d7wd3xhc) provided by NCAR's Computational and Information Systems Laboratory, sponsored by the National Science Foundation.

References

- Amann M *et al* 2011 Cost-effective control of air quality and greenhouse gases in Europe: modeling and policy applications *Environ. Modell. Softw.* **26** 1489–501
- EMEP 2016 Transboundary particulate matter, photo-oxidants, acidifying and eutrophying components (http://emep.int/publ/reports/2016/EMEP_Status_Report_1_2016.pdf)
- European Commission, and Joint Research Centre (JRC)/Netherlands Environmental Assessment (PBL) 2011 Emission Database for Global Atmospheric Research (EDGAR), release version 4.2 (<http://edgar.jrc.ec.europa.eu>)
- Garg A, Shukla P R and Kapshe M 2006 The sectoral trends of multigas emissions inventory of India *Atmos. Environ.* **40** 4608–20
- Gaur A, Tripathi S N, Kanawade V P, Tare V and Shukla S P 2014 Four-year measurements of trace gases (SO₂, NO_x, CO, and O₃) at an urban location, Kanpur, in Northern India *J. Atmos. Chem.* **71** 283–301
- Ghude S D, Pfister G G, Jena C, van der A R J, Emmons L K and Kumar R 2012 Satellite constraints of nitrogen oxide (NO_x) emissions from India based on OMI observations and WRF-Chem simulations *Geophys. Res. Lett.* **40** 423–8
- Granier C *et al* 2011 Evolution of anthropogenic and biomass burning emissions of air pollutants at global and regional scales during the 1980–2010 period *Clim. Change* **109** 163–90
- Grell G A, Peckham S E, Schmitz R, McKeen S A, Frost G, Skamarock W C and Eder B 2005 Fully coupled 'online' chemistry within the WRF model *Atmos. Environ.* **39** 6957–75
- Hsu A *et al* 2016 2016 *Environmental Performance Index* New Haven CT Yale University (www.epi.yale.edu)
- IEA 2014 *Energy Statistics of OECD Countries and Non-OECD Countries* (Paris: International Energy Agency)
- Janssens-Maenhout G *et al* 2015 HTAP_v2.2: a mosaic of regional and global emission gridmaps for 2008 and 2010 to study hemispheric transport of air pollution *Atmos. Chem. Phys.* **15** 11411–32
- Klimont Z *et al* 2009 Projections of SO₂, NO_x, and carbonaceous aerosols emissions in Asia *Tellus* **61 B** 602–17
- Klimont Z, Smith S J and Cofala J 2013 The last decade of global anthropogenic sulfur dioxide: 2000–2011 emissions *Env. Res. Lett.* **8** 014003
- Klimont Z, Kupiainen K, Heyes C, Purohit P, Cofala J, Rafaj P, Borken-Kleefeld J and Schöpp W 2016 Global anthropogenic emissions of particulate matter including black carbon *Atmos. Chem. Phys. Discuss* submitted (<https://doi.org/10.5194/acp-2016-880>)
- Kumar R, Naja M, Pfister G G, Barth M C, Wiedinmyer C and Brasseur G P 2012 Simulations over South Asia using the weather research and forecasting model with Chemistry (WRF-Chem): chemistry evaluation and initial results *Geosci. Model Dev.* **5** 619–48
- Kurokawa J, Ohara T, Morikawa T, Hanayama S, Janssens-Maenhout G, Fukui T, Kawashima K and Akimoto H 2013 Emissions of air pollutants and greenhouse gases over Asian regions during 2000–2008: regional emission inventory in Asia (REAS) version 2 *Atmos. Chem. Phys.* **13** 11019–58
- Lu Z, Zhang Q and Streets D G 2011 Sulfur dioxide and primary carbonaceous aerosol emissions in China and India, 1996–2010 *Atmos. Chem. Phys.* **11** 9839–64
- Maharashtra Pollution Control Board 2016 Data available online at (www.mpcb.gov.in/index.php) (Last Accessed: 17 December 2016)
- Naik V, Horowitz L W, Fiore A M, Ginoux P, Mao J, Aghedo A M and Levy H 2013 Impact of preindustrial to present-day changes in short-lived pollutant emissions on atmospheric composition and climate forcing *J. Geophys. Res. Atmos.* **118** 8086–110
- Nagpure A S and Gurjar B R 2012 Development and evaluation of vehicular air pollution inventory model *Atmos. Environ.* **59** 160–69
- Ohara T, Akimoto H, Kurokawa J, Horii N, Yamaji K, Yan X and Hayasaka T 2007 An Asian emission inventory of anthropogenic emission sources for the period 1980–2020 *Atmos. Chem. Phys.* **7** 4419–44
- Olivier J G J, Janssens-Maenhout G, Muntean M and Peters J A H W 2015 Trends in global CO₂ emissions: 2015 Report *PBL Netherlands Environmental Assessment Agency* The Hague
- Pandey A, Sadavarte P, Rao A B and Venkataraman C 2014 Trends in multi-pollutant emissions from a technology-linked inventory for India: II. Residential, agricultural and informal industry sectors *Atmos. Environ.* **99** 341–52
- Sadavarte P and Venkataraman C 2014 Trends in multi-pollutant emissions from a technology-linked inventory for India: I. Industry and transport sectors *Atmos. Environ.* **99** 353–64
- Sahu S K, Beig G and Parkhi N S 2012 Emerging pattern of anthropogenic NO_x emission over Indian subcontinent during 1990s and 2000s *Atmos. Pollut. Res.* **3** 262–9
- Sharma S, Chatani S, Mahtta R, Goel A and Kumar A 2016 Sensitivity analysis of ground level ozone in India using WRF-CMAQ models *Atmos. Environ.* **131** 29–40
- Smith S J, van Aardenne J, Klimont Z, Andres R J, Volke A and Arias S D 2011 Anthropogenic sulfur dioxide emissions: 1850–2005 *Atmos. Chem. Phys.* **11** 1101–16
- Streets D G *et al* 2003 An inventory of gaseous and primary aerosol emissions in Asia in the year 2000 *J. Geophys. Res.* **108** 8809
- WHO 2016 Ambient (outdoor) air pollution in cities database 2016 (www.who.int/phe/health_topics/outdoorair/databases/cities/en/)
- Zaveri R A, Easter R C, Fast J D and Peters L K 2008 Model for simulating aerosol interactions and chemistry (MOSAIC) *J. Geophys. Res. Atmos.* **113** D13204
- Zaveri R A and Peters L K 1999 A new lumped structure photochemical mechanism for large-scale applications *J. Geophys. Res. Atmos.* **104** 30387–415
- Zhang Q *et al* 2009 Asian emissions in 2006 for the NASA INTEX-B mission *Atmos. Chem. Phys.* **9** 5131–53
- Zhong M, Saikawa E, Liu Y, Naik V, Horowitz L W, Takigawa M, Zhao Y, Lin N-H and Stone E A 2016 Air quality modeling with WRF-Chem v3.5 in East Asia: sensitivity to emissions and evaluation of simulated air quality *Geosci. Model Dev.* **9** 1201–18