

Received date: 13-Feb-2016

Accepted date: 21-Apr-2016

Article Type: Opinion

Title: Reducing emissions from agriculture to meet the 2°C target

Running head: A 2°C target for agriculture

List of authors: E. Wollenberg^{1,20}, M. Richards^{1,20}, P. Smith^{17,18†}, P. Havlík^{9†}, M. Obersteiner^{9†}, F.N. Tubiello^{7†}, M. Herold^{22†}, P. Gerber^{7,22†}, S. Carter^{22†}, A. Reisinger¹⁴, D. van Vuuren^{15†}, A. Dickie^{2†}, H. Neufeldt^{8†}, B.O. Sander^{13†}, R. Wassmann^{13†}, R. Sommer^{3†}, J.E. Amonette^{16†}, A. Falcucci^{7†}, M. Herrero^{6†}, C. Opio^{7†}, R. Roman-Cuesta^{4,22†}, E. Stehfest¹⁵, H. Westhoek¹⁵, I. Ortiz-Monasterio^{5†}, T. Sapkota^{5†}, M.C. Rufino^{4†}, P.K. Thornton^{1,10}, L. Verchot^{4†}, P.C. West^{12†}, J.-F. Soussana¹¹, T. Baedeker²¹, M. Sadler²¹, S. Vermeulen^{1,19}, B.M. Campbell^{1,3}

Institute or laboratory of origin

¹ CCAFS: CGIAR Research Program on Climate Change, Agriculture and Food Security

² CEA: California Environmental Associates

³ CIAT: International Center for Tropical Agriculture

⁴ CIFOR: Center for International Forestry Research

⁵ CIMMYT: International Maize and Wheat Improvement Center

⁶ CSIRO: Commonwealth Scientific and Industrial Research Organisation

⁷ FAO: Food and Agriculture Organization of the United Nations

⁸ ICRAF: World Agroforestry Centre

⁹ IIASA: International Institute for Applied Systems Analysis

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi:

10.1111/gcb.13340

This article is protected by copyright. All rights reserved.

¹⁰ ILRI: International Livestock Research Institute

¹¹ INRA: French National Institute for Agricultural Research

¹² IONE: Institute on the Environment, University of Minnesota

¹³ IRRI: International Rice Research Institute

¹⁴ NZAGRC: New Zealand Agricultural Greenhouse Gas Research Centre

¹⁵ PBL: Netherlands Environmental Assessment Agency

¹⁶ PNNL: Pacific Northwest National Laboratory

¹⁷ Scottish Food Security Alliance-Crops

¹⁸ U Aberdeen: University of Aberdeen

¹⁹ U Copenhagen: University of Copenhagen

²⁰ UVM: University of Vermont

²¹ WB: World Bank

²² WUR: Wageningen University and Research Centre

[†]Also provided data

Corresponding author: Eva Wollenberg, lini.wollenberg@uvm.edu, Gund Institute for Ecological Economics, University of Vermont, 617 Main Street, Burlington Vermont 05405 USA +1 802 656 0890

Keywords: climate change, mitigation, agriculture, target, United Nations Framework Convention on Climate Change (UNFCCC), integrated assessment modeling, policy

Type of paper: Opinion

Abstract

More than 100 countries pledged to reduce agricultural greenhouse gas (GHG) emissions (Richards *et al.*, 2015a) in the 2015 Paris Agreement of the United Nations Framework Convention on Climate Change. Yet technical information about how much mitigation is needed in the sector versus how much is feasible remains poor. We identify a preliminary global target for reducing emissions from agriculture of ~ 1 GtCO_{2e}/yr by 2030 to limit warming in 2100 to 2°C above pre-industrial levels. Yet plausible agricultural development pathways with mitigation co-benefits deliver only 21 to 40 % of needed mitigation. The target indicates that more transformative technical and policy options will be needed, such as methane inhibitors and finance for new practices. A more comprehensive target for the 2°C limit should be developed to include soil carbon and agriculture-related mitigation options. Excluding agricultural emissions from mitigation targets and plans will increase the cost of mitigation in other sectors or reduce the feasibility of meeting the 2°C limit.

Introduction

The 2015 Paris Agreement of the United Nations Framework Convention on Climate Change (UNFCCC) aims to hold the rise in global average temperatures by 2100 to “well below 2°C above pre-industrial levels and to pursue efforts to limit the temperature increase to 1.5°C above pre-industrial levels.” A surprisingly large number of countries—at least 119—voluntarily pledged to reduce their agricultural greenhouse gas (GHG) emissions for the agreement in their statements of Intended Nationally Determined Contributions to the UNFCCC. (Richards *et al.*, 2015a). Yet how much mitigation is needed in agriculture to meet a global target versus how much is feasible remains poorly understood (Del Grosso & Cavigelli, 2012; Cafaro, 2013).

Current pledges reflect countries' interests and capacities and are limited to available technical options. Meanwhile, scenarios indicate that agricultural and agriculture-related emissions, including non-CO₂ emissions, will constitute the largest sector of surplus emissions in the future, as other sectors are projected to reduce their emissions to the maximal extent by 2030, so agriculture is critical to meeting global climate targets (Bajzelj *et al.*, 2014; Gernaat *et al.*, 2015). Excluding agricultural emissions from mitigation targets will increase the cost of mitigation in other sectors (Reisinger *et al.*, 2013) or reduce the feasibility of meeting the 2°C limit.

A global target for reduced emissions from agriculture based on meeting the 2°C limit as a threshold for warming would show the shared effort required, and in turn guide countries' ambitions, drive the development of new low emissions options and assess the global relevance of mitigation contributions. We identify here a preliminary target to guide this process. We also examine its implications by comparing the target with plausible future mitigation pathways, showing that vastly more effort is needed.

Agriculture contributes ~5.0 to 5.8 GtCO₂e/yr or ~11% of total anthropogenic GHG emissions, not including land use change (Smith *et al.*, 2014). Developing countries collectively produce the majority of agriculture-related emissions globally and are where emissions are expected to rise the fastest (Smith *et al.*, 2014). Agricultural emissions are also significant at national levels, contributing an average of 35% of emissions in developing countries and 12% in developed countries (Richards *et al.*, 2015b).

We define agricultural net emissions as the methane (CH₄) and nitrous oxide (N₂O) emissions and carbon sequestration resulting from the production of crops, livestock and agroforestry on farms. Agriculture-related emissions and opportunities for mitigation also occur in the supply chain (transport, processing fertilizer production, post harvest loss), and due to land

use change and consumption patterns (diet and food waste). One of the challenges of developing a sectoral mitigation target linked to the 2°C goal is defining the boundaries of the sector. The tools and data available currently shape how global emissions reductions are allocated to the sector. Most models use 2°C climate scenarios that focus only on non-CO₂ emissions in agriculture, as soil carbon is highly variable and involves assumptions related to organic matter inputs, carbon-nitrogen ratios, depth and bulk density, and timing of saturation, while global data on carbon in biomass, such as agroforestry, is comparatively weak. Carbon sequestration is also reversible. As a result, the target presented here is for only non-CO₂ emissions. We acknowledge the importance of other sources and sinks, however, and provide aspirational targets for the other components as preliminary guidance.

Scenarios that limit warming by 2°C

To determine the emissions budget necessary to limit warming in 2100 to no more than 2°C above pre-industrial levels, we used a scenario prepared for the Intergovernmental Panel on Climate Change (IPCC) known as Representative Concentration Pathway (RCP) 2.6 (van Vuuren *et al.*, 2011). The RCP 2.6 scenario represents 2.6 W/m² radiative forcing in 2100, or ~450 ppm of CO₂e in 2100, which results in a 66% or “likely” chance of staying below the 2°C warming limit (van Vuuren *et al.*, 2011). The RCP2.6 is one of four reference scenarios used to model concentration pathways for the IPCC.

We then compared the emissions in this desirable scenario against the business-as-usual emissions in agriculture from three integrated assessment models (IAM): Integrated Assessment of Global Environmental Change (IMAGE) (van Vuuren *et al.*, 2011), Global Change Assessment Model (GCAM) (Wise *et al.*, 2014) and Model for Energy Supply Strategy

Alternatives and their General Environmental Impact (MESSAGE) (Reisinger *et al.*, 2013).

Using this approach differs from previous estimates of agriculture's contribution that identify the wedges of mitigation possible (Pacala & Sokolow, 2004), allocate mitigation proportional to current emissions (Del Grosso & Cavigelli, 2012) or examine contributions to the total emissions budget in 2030 necessary for 2°C (Hedenus *et al.*, 2013). By using the sectoral emissions in the RCP2.6 scenario as the target, we generated a goal consistent with a 2°C pathway and based on a coherent least-cost approach across sectors.

The three IAM models used to compare the desirable 2°C degree and business-as usual worlds produce slightly different scenarios, but use similar assumptions to achieve the RCP 2.6 pathway, including: significantly increased carbon prices relative to current prices, e.g., IMAGE used 80 USD per tCO₂e in 2030 and 160 USD per tCO₂e in 2050; increased food production to meet the needs of a larger population and shifts in consumer demand; and maintaining current rates of food insecurity in the population, not eliminating it entirely. As noted previously, the models only account for non-CO₂ gases in agriculture, not soil carbon sequestration. They do, however, include bioenergy with carbon capture and storage (BECCS) to achieve the negative emissions needed to offset increases driven by an increasing population and consumption, as well as carbon sequestration associated with land use change. More details on data and methods are provided in the Supporting Information (S1-S3).

A 2030 goal

The resulting scenarios indicate that **a preliminary goal for agricultural non-CO₂ mitigation by 2030 to stay within the 2°C limit is 0.92 to 1.37 GtCO₂e/yr or about 1 GtCO₂e**. This is an annualized, not cumulative, goal. The target assumes an allowable emissions budget of 6.15 to 7.78 GtCO₂e for agriculture in 2030 (Table 1). The goal represents an 11-18%

reduction relative to the scenarios' respective 2030 business-as-usual baselines. Our estimate falls in the range of 0.3 to 2.0 GtCO₂e/yr for land-based CH₄ and N₂O emissions reductions reported by Smith *et al.* (2014) in the idealized implementation of the 2°C scenario for 2010-2050. The goal would contribute ~4-5% of the 26 GtCO₂e/yr in mitigation needed across all sectors in 2030 to achieve the 2°C limit; business-as usual emissions for all sectors in the same year are ~68 GtCO₂e (New Climate Economy, 2014).

<insert Table 1 here>

As a target for 2030, this is a near-term goal only. The scenarios show that the contribution of agriculture would need to increase in 2050 to 2.51 GtCO₂e (IMAGE) and 2.63 GtCO₂e (GCAM), reaching a maximum of 2.91 GtCO₂e in 2070-2080 using IMAGE and 4.20 GtCO₂e in 2100 using GCAM. Despite the models' different trajectories, all scenarios indicate the ongoing importance of agricultural emissions for decades to come.

Is the goal achievable?

Assuming that 1 GtCO₂e/yr in 2030 is a reasonable order of magnitude for reducing non-CO₂ emissions in the agriculture sector, is it feasible? We examined this question using the best comprehensive scientific evidence available and tested two plausible development pathways: one that reflects widespread dissemination of technical agronomic practices at prices of up to 20 USD per tCO₂e; and one based on intensified production of crops and livestock with increases in efficiency, also at prices of up to 20 USD per tCO₂e. Both pathways rely on existing practices that improve, or at least do not compromise, food productivity.

The pathway for widespread dissemination was tested by summing the mitigation achieved across agricultural technologies demonstrated to reduce non-CO₂ emissions and shows that agricultural non-CO₂ GHG emissions could be reduced by up to 0.40 GtCO₂e/yr in 2030 globally (Smith *et al.*, 2008, 2013). This technology-by-technology estimate includes livestock management, cropland management and paddy rice management practices used by the IPCC, but excludes practices related to soil carbon due to the need for consistency with the 2°C scenarios. This pathway would require implementing improved technologies with nearly universal adoption globally.

The second pathway of intensifying livestock and crop production and increasing economic efficiency was tested using the Global Biosphere Management Model (GLOBIOM). This pathway reduced agricultural non-CO₂ emissions by up to 0.21 GtCO₂e/yr in 2030 (Havlík *et al.*, 2014). The estimate reflects five broad crop and livestock sector-related structural transformations, such as transitioning from extensive rangeland systems to more efficient and productive livestock production, with accompanying improvements in livestock feed quality, breeding, reproductive efficiency, health and grassland management, and re-allocation of production to GHG-efficient regions. Soil carbon was also not included in this analysis.

Comparing the two pathways against the idealized RCP 2.6 scenarios (Figure 1) indicates that current agronomic and policy interventions compatible with food production would achieve only 21 to 40% of the needed mitigation. Neither technological dissemination as considered in Smith *et al.* (2008, 2013) nor large-scale transformation of crop and livestock production systems as analyzed by Havlík *et al.* (2014) contributes the required reduction at low costs. Even if implemented jointly, the results would fall short of the necessary mitigation, and the interventions are unlikely to be additive. Examining the mitigation possible in specific

agricultural subsectors also shows that only a fraction of the mitigation needed would be achievable with current technologies (Table S3).

<Insert Figure 1 here>

How to reduce emissions further?

The large gap between desired mitigation outcomes and plausible outcomes indicates that more transformative technical and policy options will be needed to reduce non-CO₂ emissions or that mitigation from other sources will be needed to offset them. New low emissions technologies are in the pipeline for agriculture, but vastly more effort and urgency is necessary to make options operational. Many are high-tech solutions not likely to be widely available soon, especially in the developing world. Promising options include recently developed methane inhibitors that reduce dairy cow emissions by 30% while increasing body weight without affecting milk yields or composition (Hristov *et al.*, 2015). Work is in progress to identify cattle breeds that produce less methane (Herd *et al.*, 2013) and wheat and maize varieties that inhibit the production of nitrous oxide (Subbarao *et al.*, 2015). Evidence suggests it may be possible to manage soil-plant microbial processes to increase the stability of soil organic matter and thereby retain carbon in the soil longer (Paustian *et al.*, 2016, Cotrufo *et al.*, 2013). These are each potentially transformative options, but they are not yet enough to create the menus of options needed for diverse agroecological systems and farmers to meet a mitigation target for 2°C. Coordinated research effort and investment among countries towards high-impact, quickly implementable technical options, especially for new breeds and varieties that can be easily accessed and do not require completely new management practices or inputs, is key.

More ambitious policy mechanisms also will be needed to create incentives for improved information systems and for farmers to use new practices at large scales. More productive

Accepted Article

agricultural practices, finance of low emissions agricultural development, innovative means for valuing carbon reductions, and use of government or supply chain incentives to meet sustainability standards for reduced emissions will all likely be needed. The finance and technology mechanisms in the 2015 Paris Agreement are a good start, but complementary effort will be needed at national and subnational levels, especially to engage farmers and producer organizations. Strong technical assistance for farmers, including farmer innovation hubs, two-way technical support via cell-phones, web-based information portals and farmer-to-farmer exchange, will be essential to foster changes in behavior and locally relevant options. As rapidly implementing new farming practices at large scales is risky, especially given climate uncertainties, monitoring and iterative improvement of mechanisms will be vital to provide feedback for further improvements.

The need for increased global food production by 2050 presents an opportunity to introduce mitigation measures as co-benefits of agricultural development and support farmers to leapfrog to more sustainable low emissions practices. Investments in mitigation could thereby hasten agricultural development. Special effort will be needed to ensure that new technologies are relevant, affordable and accessible to farmers in the developing world.

Other targets for agriculture

Targets linked to the 2°C limit are also needed for carbon sequestration and agriculture-related mitigation options, which can have equal or larger impacts on mitigation than practices to reduce non-CO₂ and may help offset CO₂ emissions. Improving models to produce these additional targets is a priority.

In the absence of models that enable calculations of these targets, we estimated aspirational targets for agriculture-related emissions sources based on what is achievable

globally at low costs. Where available, we used economic potentials. Soil carbon sequestration is the largest potential sink compatible with food production, mitigating ~ 1.2 GtCO_{2e}/yr in 2030 at 20 USD per tCO_{2e} (Smith *et al.*, 2014; Williamson, 2016), although its effects are easily reversed with tillage or soil disturbance. Reducing land use change due to clearing for agriculture would mitigate by 1.71 to 4.31 GtCO_{2e}/yr in 2030 at USD 20/tCO_{2e} (Carter *et al.*, 2015).

Decreasing food loss and waste by 15% (estimates vary from 30 to 50% for lost or wasted food) would reduce emissions by 0.79 to 2.00 GtCO_{2e}/yr (Stehfest *et al.*, 2013). Shifting dietary patterns, based on the diet recommended by the World Health Organization (Stehfest *et al.*, 2013) or in response to increases in carbon prices (Havlík *et al.*, 2014), would mitigate 0.31 to 1.37 GtCO_{2e}/yr in 2030. See S1-S3 for details on methods. Based on these proxy estimates, a more comprehensive goal for agriculture-related emissions would be on the order of ~ 5 -9 GtCO_{2e}/yr, or about 27% of the mitigation needed across all sectors. This estimate is consistent with Del Grosso and Cavigelli's (2012) estimate for a similar set of options.

Targets also can be organized by supply chains to mobilize action for specific subsectors or products. In the livestock supply chain, a major source of emissions globally, emissions could be reduced by about 1.77 GtCO_{2e}/yr (Gerber *et al.*, 2013). Since food production will need to increase in the coming decades, a target based on the GHG efficiency of agricultural products, or emissions intensity (GHG per unit product), is a useful secondary indicator to guide ambition and mark progress.

Conclusion

We propose that the global institutions concerned with agriculture and food security set a sectoral target to guide more ambitious mitigation and track progress toward goals. To be policy-relevant, a target for mitigation in agriculture must help achieve the 2°C warming limit while

also assuring food security. Using the RCP2.6 scenario, we identified ~1 GtCO₂e/yr by 2030 as a preliminary 2°C-based target for reducing agricultural non-CO₂ emissions. Plausible development pathways fall far short of this goal, and coordination of high-impact technical and policy interventions will be needed, including options that meet the needs of farmers in the developing world.

The proposed target is based on the best available scientific evidence, but can be improved. A more comprehensive 2°C-based target is needed that includes the full menu of options for mitigation related to agriculture. For more transformative impacts, the potential of emerging technical and policy options also should be tested using the RCP2.6 or similar scenarios. Better understanding of the sensitivity of a target to different carbon prices, alternative mitigation pathways and varied levels of food security—including full food security globally—would support more robust quantification and understanding of impacts. Better estimates of uncertainties are also needed. Aligning scenarios with a consistent emissions baseline, such as FAOSTAT's projections for agricultural emissions, or country's reported emissions would enable verification and more harmonized analysis. Scenarios for limiting warming to 1.5°C also will be needed, as even 2°C is expected to result in extensive damage and the mandate in the Paris Agreement to pursue this 1.5°C. Downscaling the target to the country levels is needed to inform countries' revised submissions of Nationally Determined Contributions to the UNFCCC (Höhne *et al.*, 2014).

As more countries seek to address climate change in the agriculture sector, linking national targets to global 2°C threshold can guide research agendas, agricultural development and national farm policy. Analysis of the investment needed in agriculture to reach the 2°C goal will inform what is economically desirable and where trade-offs might occur with other sectors.

Without the guidance of a 2°C-based goal in agriculture, much effort will be driven by what is technically or politically feasible, rather than by what is necessary. Better understanding of the gaps will show where further investment and accelerated action are really needed.

Acknowledgments

Data are presented in the Supporting Information and available from authors upon request. Thank you to Julianna White, Tapan Adhya, William Hohenstein, Tim Searchinger, Kitty Cardwell, Elise Golan and several anonymous reviewers for their comments and contributions. This work was undertaken as part of the CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS), which is a strategic partnership of CGIAR and Future Earth. This research was carried out with funding by the European Union (EU) and technical support from the International Fund for Agricultural Development (IFAD). The views expressed in the document cannot be taken to reflect the official opinions of CGIAR, Future Earth, or donors.

References

- Alexandratos N, Bruinsma J (2012) World agriculture towards 2030/2050: The 2012 Revision. Food and Agriculture Organization of the United Nations (FAO), Rome.
- Bajželj B, Richards KS, Allwood JM, Smith P, Dennis JS, Curmi E, Gilligan CA (2014) Importance of food-demand management for climate mitigation. *Nature Climate Change*, **4**, 924-929.
- Beach RH, Deangelo BJ, Rose S, Li C, Salas W, Delgrosso SJ (2008) Mitigation Potential and Costs for Global Agricultural Greenhouse Gas Emissions. *Agricultural Economics*, **38**, 109–115.

Accepted Article

Cafaro P (2013) Avoiding Catastrophic Climate Change: Why Technological Innovation is Necessary but not Sufficient. *Ethics and Emerging Technologies*. New York, NY: Palgrave Macmillan, 424-438.

Carter S, Herold M, Rufino MC, Neumann K, Kooistra L, Verchot L (2015) Mitigation of agriculture emissions in the tropics: comparing forest land-sparing options at the national level. *Biogeosciences Discussions* **12**, 5435–5475.

Cotrufo, MF, Wallenstein, M, Boot, CM, Denef, K & Paul, EA (2013) The Microbial Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition with soil organic matter stabilization: do labile plant inputs form stable soil organic matter? *Glob. Change Biol.* **19**, 988–995. □

DeAngelo BJ, de la Chesnaye F, Beach RH, Sommer A, Murray BC (2006) Methane and nitrous oxide mitigation in agriculture. *The Energy Journal*, **27**, 89–108.

Del Grosso, SJ, Cavigelli MJ (2012) Climate stabilization wedges revisited: can agricultural production and greenhouse gas reduction goals be accomplished? *Frontiers in Ecology and the Environment*, **10**, 571-578.

Gerber PJ, Steinfeld H, Henderson B *et al.* (2013) *Tackling climate change through livestock – A global assessment of emissions and mitigation opportunities*. Food and Agriculture Organization of the United Nations (FAO), Rome.

Gernaat DEHJ, Calvin K, Lucas PL *et al.* (2015) Understanding the contribution of non-carbon dioxide gases in deep mitigation scenarios. *Global Environmental Change*, **33**, 142–153.

Havlík P, Valin H, Herrero M *et al.* (2014) Climate change mitigation through livestock system transitions. *Proceedings of the National Academy of Sciences of the United States of America*, **111**, 3709-3714.

Hedenus F, Wirsenius S, Johansson DJA (2014) The importance of reduced meat and dairy consumption for meeting stringent climate change targets. *Climatic Change*, **124**, 79-91.

Herd RM, Bird SH, Donoghue KA, Arthur PF, Hegarty RF (2013) Phenotypic associations between methane production traits, volatile fatty acids and animal breeding traits. *Proceedings of the Association for the Advancement of Animal Breeding and Genetics*, **20**, 286-89.

Höhne N, den Elzen M, Escalante D (2014) Regional GHG reduction targets based on effort sharing: a comparison of studies. *Climate Policy*, **14**, 122-147.

Hristov AN, Oh J, Giallongo F *et al.* (2015) An inhibitor persistently decreased enteric methane emission from dairy cows with no negative effect on milk production. *Proceedings of the National Academy of Sciences of the United States of America*, **112**, 10663–10668.

Kriegler E, Weyant P, Blanford G *et al.* (2013) The role of technology for achieving climate policy objectives: overview of the EMF27 study on global technology and climate policy strategies. *Climatic Change* **123**, 353-367.

Lucas PL, van Vuuren DP, Olivier JGJ, den Elzen MGJ (2007) Long-term reduction potential of non-CO₂ greenhouse gases. *Environmental Science and Policy* **10**, 85–103.

New Climate Economy (2014) Better growth, better climate: The New Climate Economy Report. Washington, DC.

Paustian, K, J Lehmann, S Ogle, D Reay, GP Robertson & P. Smith (2016) Climate-smart soils. *Nature* **532**, 49–57.

Pacala S, Socolow R (2004) Stabilization wedges: solving the climate problem for the next 50 years with current technologies. *Science*, **305**, 968-972.

Powlson DS, Whitmore AP, Goulding KWT (2011) Soil carbon sequestration to mitigate climate change: a critical re-examination to identify the true and the false. *European Journal of Soil Science*, **62**, 42–55.

Reisinger A, Havlík P, Riahi K, van Vliet O, Obersteiner M, Herrero M (2013) Implications of alternative metrics for global mitigation costs and greenhouse gas emissions from agriculture. *Climatic Change*, **117**, 677–690.

Richards M, Bruun T, Campbell BM, Huyer S, Kuntze V, Gregersen LE, Madsen STN, Oldvig MB, Vasileiou I (2015a) How countries plan to address agricultural adaptation and mitigation: An analysis of Intended Nationally Determined Contributions. CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS), Copenhagen.

Richards MB, Wollenberg E, Buglion-Gluck S (2015b) Agriculture's contributions to national emissions. CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS), Copenhagen.

Smith P, Martino D, Cai Z *et al.* (2008) Greenhouse gas mitigation in agriculture. *Philosophical Transactions of the Royal Society B*, **363**, 789-813.

Smith P, Haberl H, Popp A *et al.* (2013) How much land-based greenhouse gas mitigation can be achieved without compromising food security and environmental goals? *Global Change Biology*, **19**: 2285–2302.

Smith P, Bustamante M, Ahammad H *et al.* (2014) Agriculture, Forestry and Other Land Use (AFOLU). In: *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds. Edenhofer O, Pichs-Madruga R, Sokona Y *et al.*) Cambridge University Press, Cambridge.

Accepted Article

Stehfest E, Berg M V D, Woltjer G, Msangi S, Westhoek H. 2013. Options to reduce the environmental effects of livestock production - Comparison of two economic models. *Agricultural Systems* **114**:38-53.

Subbarao GV, Yoshihashi T, Worthington M *et al.* (2015) Suppression of soil nitrification by plants. *Plant Science* 233, 155-164.

van Vuuren DP, Stehfest E, den Elzen MGJ *et al.* (2011) RCP2.6: Exploring the possibility to keep global mean temperature increase below 2 degrees C. *Climatic Change*, **109**, 95–116.

Williamson, P (2016). Scrutinizing CO₂ removal methods. *Nature* 530: 153-155.

Wise M, Calvin K, Kyle P, Luckow P, Edmonds J (2014) Economic and physical modeling of land use in GCAM 3.0 and an application to agricultural productivity, land, and terrestrial carbon. *Climatic Change Economics*, **5**, 145003.

Supporting Information

Appendix S1. Overview of methods, including Tables S1-S4

Appendix S2. Data sources and methods, including Figure S1 and Tables S5-S10

Appendix S3. 2030 reference levels

Appendix S4. References

Table 1. GHG emissions and mitigation needed in the agriculture sector in 2030 to avoid exceeding 2°C

Model	Model category	Basis for non-CO ₂ mitigation	Baseline 2030 emissions GtCO ₂ e	450 ppm scenario emissions GtCO ₂ e	Mitigation modeled
IMAGE RCP 2.6 (Van Vuuren <i>et al.</i> , 2011)	Recursive dynamic partial equilibrium model	US-EPA MAC curves based on Lucas <i>et al.</i> (2007)	7.52	6.15	1.37
GCAM (Wise <i>et al.</i> , 2014)	Recursive dynamic partial equilibrium model	US-EPA MAC curves based on DeAngelo <i>et al.</i> (2006)	8.97	7.78	1.19
MESSAGE (Reisinger <i>et al.</i> , 2013)	Intertemporal optimization general equilibrium model	US-EPA MAC curves based on Beach <i>et al.</i> (2008)	8.58	7.66	0.92

