

Journal Pre-proof

Comparative analysis of greenhouse gas emission inventory for Pakistan: Part I energy and industrial processes and product use

Kaleem Anwar Mir, Chunkyoo Park, Pallav Purohit, Seungdo Kim



PII: S1674-9278(20)30020-4

DOI: <https://doi.org/10.1016/j.accre.2020.05.002>

Reference: ACCRE 172

To appear in: *Advances in Climate Change Research*

Received Date: 2 January 2020

Revised Date: 9 March 2020

Accepted Date: 13 May 2020

Please cite this article as: Mir, K.A., Park, C., Purohit, P., Kim, S., Comparative analysis of greenhouse gas emission inventory for Pakistan: Part I energy and industrial processes and product use, *Advances in Climate Change Research*, <https://doi.org/10.1016/j.accre.2020.05.002>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Copyright © 2020, National Climate Center (China Meteorological Administration). Production and hosting by Elsevier B.V. on behalf of KeAi. All rights reserved.

Title Page Information

Comparative analysis of Greenhouse gas emission inventory for Pakistan – Part I: Energy and industrial processes and product use

Kaleem Anwar Mir^{a,e,*}, Chunkyoo Park^b, Seungdo Kim^c, Pallav Purohit^d

^aResearch Centre for Climate Change & Energy, Department of Environmental Sciences & Biotechnology, Hallym University, Chuncheon, 24252, Republic of Korea
Email: kaleemanwar.mir@gmail.com
Designation: PhD Student

^bMinistry of Environment, Government Complex-Sejong, Sejong-si, 30103, Republic of Korea
Email: ckpark1991@naver.com
Designation: Vice Minister

^cResearch Centre for Climate Change & Energy, Department of Environmental Sciences & Biotechnology, Hallym University, Chuncheon, 24252, Republic of Korea
Email: sdkim@hallym.ac.kr
Designation: POSCO Professor/Director

^dInternational Institute for Applied Systems Analysis, Schlossplatz 1, A-2361, Laxenburg, Austria
Email: purohit@iiasa.ac.at
Designation: Research Scholar

^eGlobal Change Impact Studies Centre, Ministry of Climate Change, Government of Pakistan, Islamabad, 44000, Pakistan
Email: kaleemanwar.mir@gmail.com
Designation: Scientific Officer

* Corresponding author.

E-mail address: kaleemanwar.mir@gmail.com (Kaleem Anwar Mir)

1 Comparative analysis of greenhouse gas emission inventory for 2 Pakistan: Part I energy and industrial processes and product use

3 Kaleem Anwar MIR ^{a,d*}, Chunkyoo PARK^b, Pallav PUROHIT ^c, Seungdo KIM ^a

4
5 ^a *Research Centre for Climate Change and Energy, Department of Environmental Sciences and Biotechnology,*
6 *Hallym University, Chuncheon, 24252, Republic of Korea*

7 ^b *Ministry of Environment, Government Complex-Sejong, Sejong-si, 30103, Republic of Korea*

8 ^c *International Institute for Applied Systems Analysis, Laxenburg, A-2361, Austria*

9 ^d *Global Change Impact Studies Centre, Ministry of Climate Change, Government of Pakistan,*
10 *Islamabad,44000, Pakistan*

11 12 Abstract

13 In order to further improve the accuracy and reliability and reduce uncertainties in the national GHG
14 inventories for Pakistan, this study call for using 2006 IPCC Guidelines, to help to identify the
15 national targets for GHG mitigation with respect to the nationally determined contributions (NDCs).
16 GHG (CO₂, CH₄, and N₂O) inventories for Pakistan have been developed by conducting a detailed
17 sectoral assessment of IPCC source sectors, energy, industrial processes and product use (IPPU),
18 agriculture, forestry and other land use (AFOLU), and the waste sector. Further, sector wise
19 comparative analysis of GHG inventories (1994–2017) based on the 2006 and 1996 IPCC Guidelines
20 have also been presented. Results indicated an average relative difference of 4% in total GHG
21 emissions (CO₂ equivalent) from energy sector between 2006 and 1996 IPCC Guidelines. With 3.6%
22 average annual growth rate based on 2006 IPCC Guidelines, CO₂ from energy sector remained the
23 most abundant GHG emitted, followed by CH₄ and N₂O. While the average absolute difference in
24 emissions of CH₄ and N₂O from the energy sector is notable, the total estimated GHG emissions by
25 2006 IPCC Guidelines duplicate those by 1996 IPCC Guidelines. In the mineral industry with 2006
26 IPCC Guidelines, an average annual growth rate of 6.7% is observed, contributing 64% of total IPPU
27 sector CO₂ emissions. Nevertheless, the relative difference between the two Guidelines in overall
28 IPPU sector emissions remained negligible. There might be a need for switching to 2006 IPCC
29 Guidelines to consider more parameters such as additional source sectors and new default emission
30 factors that fit into national circumstances.

31 Keywords

32 Greenhouse gas, Emission inventory; Energy sector; Industrial processes and product use; Pakistan

33 1. Introduction

* Corresponding author.

E-mail address: kaleemanwar.mir@gmail.com (K.A. Mir)

1 Implementation of the Paris Agreement (UNFCCC, 2015) requires all parties to report their national
2 anthropogenic greenhouse gas (GHG) emissions to the United Nations Framework Convention on
3 Climate Change (UNFCCC) on a regular basis and to frequently analyze the sum of global emissions
4 in the process of global stocktaking (UN, 1992; UNFCCC, 2015). An essential part of the Paris
5 Agreement is the transparency framework, to ensure accurate, transparent, comparable, consistent and
6 complete reporting of GHG inventories, building on the methodologies developed by the
7 Intergovernmental Panel on Climate Change (IPCC) (Bergamaschi et al., 2018). Reporting for non-
8 Annex I Parties to the UNFCCC is implemented through national communications (NCs) and biennial
9 update reports (BURs), subject to the availability of financial support (Zhu and Wang, 2013). The
10 Government of Pakistan submitted its Initial National Communication (Pak-INC) (UNFCCC, 2003a)
11 to the UNFCCC on November 15, 2003 and Second National Communication (Pak-SNC) (UNFCCC,
12 2019) on August 9, 2019 with national GHG inventories for the years 1994 (UNFCCC, 2003a) and
13 2015 (GCISC, 2017), respectively. These two inventories were prepared following the Revised 1996
14 IPCC Guidelines for National GHG Inventories (hereinafter referred to as the 1996 GLs) (IPCC,
15 1997). Moreover, GHG inventories for the years 2008 (ASAD, 2016), and 2012 (GCISC, 2016; Mir et
16 al., 2017) have also been completed in indigenous capacities using the same 1996 GLs. Nevertheless,
17 since the publication of 2006 IPCC Guidelines for National GHG Inventories (hereinafter referred to
18 as 2006 GLs) (IPCC, 2006) and UNFCCC Decision 24/CP.19 (UNFCCC, 2014), though it is not
19 mandatory (Yona, 2020) but Pakistan should make efforts to prepare its national GHG inventories
20 using the latest 2006 GLs. This would also be in accordance with the criteria set out in the revised
21 UNFCCC guidelines for preparing the national communications of non-Annex I countries (UNFCCC,
22 2003b), which encourages developing countries to use the latest available methodologies. Further, the
23 2006 GLs have several advantages over the 1996 GLs in terms of additional sources, new default
24 emission factors, guidance on choosing appropriate estimation methods for individual inventory
25 categories, and cross-sectoral good practice guidance e.g. key category analysis to identify most
26 important inventory categories (Breidenich, 2011). It might at first result in facing the challenge of
27 modifying Pakistan's nationally determined contributions (NDCs) (UNFCCC, 2016) in line with the
28 new inventories estimates.

29 With the exception of ASAD (2009), the 2006 GLs have never been implemented for calculating
30 national GHG emissions in Pakistan. In addition, neither higher Tiers nor country-specific parameters
31 were used or identified in Pak-INC and Pak-SNC, the default approach (Tier 1) was applied in both
32 inventories by using 1996 GLs (UNFCCC, 2003a, 2019; GCISC, 2017). There is an urgent need of at
33 least shifting to 2006 GLs which might consider more parameters (e.g. additional source sectors, new
34 default emission factors) that fit into national circumstances and reduce the uncertainty of GHG
35 emissions estimates. This study deals with a more technical, improved, and comprehensive time series
36 (1994–2017) evaluation of Pakistan's GHG emissions inventories using the latest 2006 GLs and their

1 comparison with those prepared following the old 1996 GLs. This would provide an insight into: the
2 difference in GHG emission quantities between both GLs; the mitigation priorities that need to be
3 considered in future; and the importance of applicability of 2006 GLs which has been hardly
4 recognized by developing countries.

5 **2. Methods and data**

6 **2.1 Data sources**

7 The activity data used was acquired from official national government documents such as: Pakistan
8 Energy Year Book by the Hydrocarbon Development Institute of Pakistan (HDIP, 1994, 2008, 2012,
9 2015, 2107) , Pakistan Economic Survey by the Ministry of Finance (MoF 2008, 2012, 2015, 2017),
10 Pakistan Agricultural Statistics by the Ministry of National Food Security and Research (MoNFSR
11 2019), Industrial Statistics from the Ministry of Industries and Production Year Book (MoIP, 2017);
12 and Pakistan Forest Resources Assessment (FAO, 2015) by the Ministry of Climate Change (MoCC).
13 In addition, country specific information from a few international sources such as the Food and
14 Agriculture Organization (FAO), the World Bank (WB), the United States Geological Survey
15 (USGS), and the United Nations (UN) was also accessed.

16 At the time of current study, the GHG inventories data estimated by UNFCCC Non-Annex I National
17 Greenhouse Gas Inventory Software for four years (1994, 2008, 2012, and 2015) based on the Tier 1
18 approach in the 1996 GLs, was available (UNFCCC 2003a; ASAD, 2009, 2016; GCISC, 2016, 2017;
19 Mir et al. 2017). In addition, Pakistan's latest GHG inventory (2017) was estimated as part of the
20 present work using the same 1996 GLs methodology and data sources as used in previously available
21 inventories to maintain the consistency. Following the 1996 GLs, the source sectors included in all
22 these five-year (1994, 2008, 2012, 2015, and 2017) GHG inventories were energy, industrial
23 processes, agriculture, land use change and forestry (LUCF), and waste. The linear statistical
24 interpolation method was then applied to the detailed data points of the five-year inventories at the
25 national, sectoral, and sub-sectoral levels to develop the estimates for the missing intermediate years.
26 This completed the development of time series (1994–2017) of the estimates based on the 1996 GLs.
27 The year 1994 was considered as the base year and 2017 as the latest year. The same data points of
28 the five-year inventories were then updated and estimated following the latest 2006 GLs together with
29 the corresponding sectoral worksheets by considering same source sectors, method, and data.

30 **2.2 Emission estimation**

31 In general, both sets of GLs follow the same methodological approach (IPCC, 1997, 2006). This
32 approach involves integrating information on the level of human activity, known as Activity Data,
33 with the quantified emission coefficients per unit activity, known as the Emission Factor. Therefore,
34 the fundamental equation that was used to calculate the GHG emissions from different source sectors
35 is: Emissions = Activity Data × Emission Factor. Although the country-specific sectoral activity data
36 for multiple years based on national official statistics have been used, all the emission factors were
37 the default values provided by both the 1996 and 2006 GLs. The simplest Tier 1 sectoral approach,
38 typically requiring the most basic and least disaggregated activity details along with the default
39 emission factors of IPCC, was used to calculate the emissions of three GHGs, CO₂, CH₄, and N₂O.

40 **2.3 Missing data management**

41 According to the IPCC GLs, the national GHG inventories must be recorded in the calendar year in
42 which the atmospheric emissions occur (IPCC, 2006). Nonetheless, if the unavailability of sufficient
43 data prevents compliance with this rule, emissions can be calculated using information from other
44 years by applying reasonable splicing techniques such as overlapping, surrogate approach,

1 interpolation, and trend extrapolation to complete the time series (IPCC, 2006). In this analysis, since
2 the inventories information from 1994 to 2017 were available for five distinct years (1994, 2008,
3 2012, 2015, and 2017), the data for the missing intermediate years were interpolated linearly to
4 complete the time series. This method seems practical as the overall trend tended to be stable, and real
5 emissions for missing intermediate years are unlikely to vary significantly from the
6 predicted interpolation values.


7 **3. Time series GHG inventories using 2006 GLs**

8 **3.1 Energy sector**

9 Two versions of the CO₂ emissions of the energy sector exist: the Sectoral Approach and the
10 Reference Approach (RA). Under the convention, the SA is used for inventory development whereas
11 the Reference Approach is used for verification purposes only (quality control activity). The Sectoral
12 Approach is based on data collected from the Pakistan Energy Year Book (HDIP, 1994, 2008, 2012,
13 2015, 2017) and additional source-specific information. The Sectoral Approach combines fossil fuel
14 consumption statistics with bottom-up information and calculations based on fuel consumption
15 models. On the other hand, the Reference Approach employs a top-down strategy based on the
16 apparent consumption of energy by Pakistan as reflected in the energy production statistics of the
17 Pakistan Energy Year Book. Within the dataset, the differences in energy consumption and CO₂
18 emissions between Reference Approach and Sectoral Approach are determined. For both methods, the
19 CO₂ emissions and difference in energy consumption are below 5% for the entire period and do not
20 need any clarification according to the 2006 GLs. In addition, the default 100% oxidation rate for
21 various types of fuel combustion is used in the new 2006 GLs based inventories compared to the old
22 1996 GLs based, where the default oxidation rate used for coal, natural gas, and oil was 98%, 99.5%,
23 and 99%, respectively.

24 The energy sector is the most important source of GHG emissions in Pakistan. The CO₂ produced by
25 the energy sector is the most abundant GHG released into the atmosphere, followed by CH₄ and N₂O
26 (Fig. 1a), with an average annual growth rate of 3.6%. With the exception of a marked increase
27 between 2015 and 2017 owing to a substantial increase in the amount of energy consumed by new
28 power plants for electricity generation, GHG emissions from the energy sector continued to expand
29 over the period examined. The almost flat pattern in the period 2008–2012 is the result of the energy
30 crises in Pakistan. The main types covered by the energy sector are fuel combustion and fugitive
31 emissions from fuels. Four source categories dominate the GHG emissions in Pakistan's energy sector.
32 The energy industries (mainly electricity generation) and manufacturing sectors are the primary
33 sources of GHG emissions (Fig. 1b). Transport (mainly road transport) and other sectors (commercial,
34 residential, and agriculture) also play an important role in national GHG emissions.

35



1

2 **Fig. 1. GHG emissions from the energy sector of Pakistan using 2006 GLs during 1994–2017 by**
 3 **gas (a), and by sub-sector (b)**

4 3.1.1 Fuel combustion

5 The fuel combustion sub-sector constitutes, among other sub-sectors (fugitive emissions), the largest
 6 share representing more than 97% of total emissions from the energy sector. Combustion activities
 7 include both stationary and mobile combustion operations that represent almost all combustion
 8 activities in Pakistan. The fuel combustion sub-sector primarily comprises four categories, specifically
 9 the energy industries (power), manufacturing industries, transportation, and other sectors
 10 (commercial, residential, and agriculture). Table 1 provides a comprehensive overview of the amount
 11 of GHG emissions produced by the fuel combustion categories. The category of energy industries
 12 adds most of the GHG emissions of the energy sector, representing 31% of the total average GHG
 13 emissions of this sector. This is followed by the manufacturing industries, which contributed about
 14 30%. The GHG emissions from the transport category and other sectors are lower than other
 15 categories and represented an average of about 21% and 15% of the total GHG emissions of the
 16 energy sector, respectively. The fugitive emissions have a significantly reduced share (3%) of the total
 17 national GHG emissions of Pakistan.

18 It is observed that the average annual growth rate of the GHG emissions (approximately 2.5%) is
 19 comparatively low for the manufacturing industries, despite this category being the second largest
 20 contributor to GHG emissions in the energy sector. The gas-consuming manufacturing industries
 21 include steel mills, cement, fertilizer (as consumers of fuel) and general industries in Pakistan. The
 22 major coal consumers in the manufacturing sector are the steel, cement, and brick kiln industries. In
 23 addition, the natural gas used in gas processing plants belonging to other energy industries (sub-sector
 24 of manufacturing of solid fuels and other energy industries) has the highest average annual growth
 25 rate of nearly 8%. Similar to the energy sector as a whole, CO₂ holds the largest amount and share of
 26 the GHG emissions in transport sector. Owing to the significant contribution of road transport (90%

1 with an average annual growth rate of approximately 4.4%) the transport sector emissions increased
2 by approximately 4.2% on average. The combined average annual growth rate for rail, navigation,
3 and other transportation and domestic aviation was 2.1% and 1.8% respectively.

4 It is observed that all categories in the energy sector had a positive annual growth rate in GHG
5 emissions during 1994–2017. This specifies the increase in fuel consumption in this sector due to
6 increase in energy demand owing to urbanization, economic growth and population in Pakistan. The
7 GHG emissions resulting from fuel combustion are mainly associated with the amount of fuel burned
8 in respective sectors. The CO₂ emissions from these categories are determined on the basis of the fuel
9 used by each sector and the carbon content of the fuel, irrespective of the combustion technology or
10 emission control technology in use. The carbon content of fuel is relative to the default IPCC values,
11 but for the purpose of calculating GHG emissions, the gross calorific value (GCV) is translated to the
12 net calorific value (NCV) according to the 2006 GLs.

13 3.1.2 Fugitive emission

14 This sub-sector consists of three sources of fugitive emissions (primarily CH₄), solid fuels, oil and gas
15 systems, and other energy production. The first two categories of fugitive CH₄ emissions are
16 estimated in this study. The total fugitive CH₄ emission constitutes 3% of the total GHG
17 emission from the energy sector. However, an average annual growth rate of 2.5% is noted for
18 fugitive CH₄ emissions, as shown in Table 1. The relatively small percentage of GHG released from
19 this group is mainly because of Pakistan's lower production of oil, gas, and coal. There is, however, a
20 significant difference between the estimates using the 2006 GLs and 1996 GLs, respectively,
21 especially for fugitive CH₄ emissions from the production and distribution of natural gas. This
22 distinction is attributable to the implementation of very high emission factors by using the 1996 GLs.
23 Nevertheless, the emission factor of the 2006 GLs has been revised and has a reduced range relative to
24 that of the 1996 GLs.

1 **Table 1. GHG emissions from the energy sector of Pakistan for 1994–2017 (2006 GLs) (unit: Mt CO₂-eq)**

Year	Fuel combustion								Fugitive emissions			Total ^d (A+B)
	Energy industries		Manufacturing industries	Transport			Other sectors ^b	Total (A)	Coal mining	Oil & natural gas	Total (B)	
	Electricity generation	Other energy industries ^a		Road	Domestic aviation	Rail & other						
1994	22.07	2.74	27.67	15.91	1.34	0.66	12.92	83.31	1.03	1.20	2.23	85.54
1995	23.48	2.79	28.96	16.71	1.37	0.72	13.47	87.50	1.04	1.29	2.33	89.83
1996	24.89	2.83	30.25	17.50	1.40	0.78	14.03	91.69	1.05	1.39	2.44	94.13
1997	26.29	2.88	31.54	18.30	1.44	0.85	14.58	95.87	1.07	1.49	2.55	98.42
1998	27.70	2.92	32.83	19.09	1.47	0.91	15.14	100.06	1.08	1.58	2.66	102.72
1999	29.11	2.97	34.12	19.89	1.51	0.97	15.69	104.24	1.09	1.68	2.77	107.01
2000	30.52	3.01	35.40	20.68	1.54	1.03	16.24	108.43	1.10	1.77	2.87	111.31
2001	31.93	3.06	36.69	21.47	1.57	1.09	16.80	112.62	1.11	1.87	2.98	115.60
2002	33.34	3.10	37.98	22.27	1.61	1.15	17.35	116.80	1.13	1.97	3.09	119.89
2003	34.74	3.15	39.27	23.06	1.64	1.22	17.91	120.99	1.14	2.06	3.20	124.19
2004	36.15	3.19	40.56	23.86	1.67	1.28	18.46	125.18	1.15	2.16	3.31	128.48
2005	37.56	3.24	41.85	24.65	1.71	1.34	19.02	129.36	1.16	2.25	3.41	132.78
2006	38.97	3.28	43.14	25.45	1.74	1.40	19.57	133.55	1.17	2.35	3.52	137.07
2007	40.38	3.33	44.42	26.24	1.77	1.46	20.12	137.73	1.19	2.45	3.63	141.36
2008	41.79	3.38	45.71	27.04	1.81	1.53	20.68	141.92	1.20	2.54	3.74	145.66
2009	41.18	4.74	44.05	27.81	1.85	1.39	21.25	142.28	1.16	2.56	3.72	146.00
2010	40.58	6.10	42.39	28.58	1.90	1.25	21.83	142.64	1.12	2.58	3.71	146.35
2011	39.98	7.47	40.73	29.36	1.95	1.12	22.40	143.00	1.09	2.60	3.69	146.69
2012	39.38	8.83	39.07	30.13	1.99	0.98	22.97	143.36	1.05	2.63	3.68	147.03
2013	40.98	7.84	39.21	31.41	1.93	0.95	23.48	145.79	1.06	2.60	3.66	149.45
2014	42.59	6.85	39.34	32.68	1.87	0.92	23.99	148.22	1.06	2.58	3.65	151.87
2015	44.20	5.85	39.47	33.95	1.80	0.89	24.50	150.65	1.07	2.56	3.63	154.29
2016	46.11	10.66	44.40	38.58	1.91	0.98	25.59	168.21	1.13	2.63	3.76	171.97
2017	48.02	15.46	49.33	43.20	2.01	1.07	26.68	185.78	1.20	2.69	3.89	189.66
CAGR ^c	3.4%	7.8%	2.5%	4.4%	1.8%	2.1%	3.2%	3.5%	0.7%	3.6%	2.5%	3.5%

^a This includes fossil fuel combustion in petroleum refining and gas processing plants.

^b This includes fossil fuel combustion in the residential, commercial/institutional, and agricultural/forestry/fishing sectors.


^c CAGR is the compound annual growth rate, calculated by the formula $(\text{latest value}/\text{base value})^{(1/\text{no. of years})} - 1$.

^d May not sum similar to the total due to rounding.

1 3.2 Industrial processes and product use

2 The industrial process and product use (IPPU) sector is the third largest contributor to total GHG
 3 emissions from Pakistan after the AFOLU sector. IPPU-emitted GHGs vary from other sectors as they
 4 consist of CO₂, hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), and sulfur hexafluoride (SF₆)
 5 (all from product use). Fluorinated GHG emissions (including HFCs, PFCs, and SF₆) have never been
 6 reported because of the lack of activity data in Pakistan. CO₂ emissions show a steady increase
 7 (average annual growth rate of 4.2%) throughout the time series as a result of continued growth in
 8 cement production in the mineral industry which is the primary contributor to industrial process
 9 emissions. The emissions from the IPPU sector are mainly from the mineral, chemical, and metal
 10 industries. Fig. 2 shows GHG emissions from the IPPU sector of Pakistan using 2006 GLs.

11



12

13 **Fig. 2. GHG emissions from the industrial process and product use (IPPU) sector of Pakistan,**
 14 **during 1994–2017 using 2006 GLs**

15 Table A1 provides an aggregate overview of GHG emissions from the IPPU sector and the percentage
 16 share. The largest average annual growth rate (6.7%) is observed in the mineral industry sub-sector,
 17 which mainly includes cement production. GHG emissions associated with imports and exports of
 18 clinker bricks is assumed to be zero in Pakistan as assumed in past national GHG inventories. The
 19 other category in the mineral industry is lime production, which forms 5% of the share of the mineral
 20 industry in this subsector. The remaining 95% is that of cement production. The proportion of
 21 limestone and dolomite extraction/production data (MoF, 2017) in Pakistan (mainly in the Pakistan's
 22 steel industry) remained the same (4% of limestone extraction/production; 73% of dolomite
 23 extraction/production) as in the Pak-INC inventory by assuming the fact that the capacity of Pakistan
 24 steel has not increased since 1994 and has remained constant. In the IPPU sector the overall share of
 25 the mineral industry is 64% followed by the chemical industry (27%).

26 Further 2006 GLs also require data regarding the mass of the lime (high-calcium lime and dolomite
 27 lime) produced rather than the mass of limestone or dolomite produced/extracted. Therefore, further
 28 calculation was carried out to convert limestone/dolomite (CaCO₃/CaMg(CO₃)₂) production/usage
 29 data into the mass of lime (high calcium lime and dolomite lime) produced by using the conversion
 30 factor given in the 2006 GLs i.e., 1 t of lime (CaO) requires the calcination of 1.785 t of CaCO₃. The
 31 Pakistan Economic Survey provides data on the production of soda ash rather than the use of soda
 32 ash, therefore the emissions from soda ash production are reported in the chemical industry according
 33 to the 2006 GLs, whereas emissions resulting from the use of soda ash are included in the respective
 34 end-use sectors in which soda ash is used. Thus, based on the availability of data, the chemical
 35 industry worksheet of 2006 GLs was used for CO₂ emissions from soda ash production. According to
 36 the 2006 GLs, soda ash emissions are included in the emissions of the chemical industry, whereas in
 37 previous inventories they are included in the mineral industry worksheet and applied to the overall

1 emissions from the mineral industry. In terms of metal production in 2017, the value is zero because
2 the Pakistan's Steel industry was shut down in 2017 and has not yet been operational. This is why the
3 metal production emission growth rate dropped to -100% due to zero value in the year 2017.

4 **4. Comparative analysis of GHG inventories using 2006 and 1996 GLs**

5 **4.1 Source categorization and global warming potential**

6 It is evident that under 2006 GLs, energy and waste remain separate sectors. However, industrial
7 processes, and solvent and other product use are integrated as one sector –IPPU. To deal with
8 emissions from the non-energy use of fuels, 2006 GLs clearly establishes the boundary between the
9 energy sector and IPPU compared to 1996 GLs, and such emissions are now reported mainly in the
10 IPPU sector. The 2006 GLs introduces a broader concept of 'excluded carbon' for the non-energy use
11 of fuels which includes not only 'stored carbon' (old term in 1996 GLs) but also carbon used and
12 emitted as CO₂ in other sectors quite often within the IPPU (not just in the energy sector). The 2006
13 GLs also merge agriculture, and land use change and forestry as the agriculture, forestry and other
14 land use (AFOLU) sector to facilitate effective use of information. The source categorization also
15 varies on a more disaggregated level between the two versions of IPCC GLs.

16 Another major distinction between the 2006 GLs and the 1996 GLs is the defined global warming
17 potential (GWP) values. Decision 17/CP.8 under the convention states that the GWP values provided
18 by the IPCC in its Second Assessment Report based on the effects of GHGs over a 100-year time
19 horizon should be used by non-Annex I Parties (UNFCCC, 2003b). However, the Annex-I countries
20 from 2015 onwards will use the GWP values provided by the IPCC Fourth Assessment Report (IPCC,
21 2007) as agreed by Decision 24/CP.19 (UNFCCC, 2014). The GWP values used in reporting
22 aggregated emissions based on 1996 GLs were taken from IPCC in its Second Assessment Report.
23 Therefore, the same GWP values are also used in reporting aggregated emissions based on 2006 GLs
24 to keep the consistency.


25

26 **4.2 Comparison by sector**

27 **4.2.1 Energy sector overall difference**

28 Figure 3 shows the overall gap in the time series of 2006 and 1996 GLs estimates. Although the gap is
29 noticeable for CH₄ and N₂O, the total CO₂-eq emissions according to the 2006 GLs replicate those
30 based on the 1996 GLs because the CO₂ emissions are dominant and demonstrate similar emission
31 quantities over the period of interest. The significant difference in CH₄ emission quantities is due to
32 the large inconsistency present in the CH₄ emission factors between two GLs for estimating fugitive
33 emissions from oil (production, transport, and refining) and natural gas (processing, and distribution)
34 operations.

1



2


3 **Fig. 3. Energy sector overall gap during 1994–2017 of (a) CO₂, (b) CH₄, (c) N₂O, and (d) CO₂-eq**
 4 **emissions in Pakistan, 2006 vs. 1996 GLs**

5 **4.2.2 Energy sector sub-sectoral differences**

6 **4.2.2.1 Energy industries**

7 Figure 4 shows the difference between the emission (CO₂, CH₄, N₂O, and the total) estimates from the
 8 2006 and 1996 GLs in the energy industries. These industries primarily cover fuel consumption for
 9 electricity generation, gas processing plants, and petroleum refining. The total amounts of the total
 10 estimated by both GLs overlay each other and show quantities that are almost similar. The trends in
 11 the estimates of CO₂ emissions are also similar to those of CO₂-eq. However, the CH₄ emissions
 12 remained uniform until 2015, after which they doubled (0.001 Mt to 0.002 Mt) in 2017 (with the 2006
 13 GLs) because Pakistan did not include the fuel consumption for petroleum refinery in the inventories
 14 based on the 1996 GLs. Petroleum refining was also considered for the other years (1994, 2008, 2012,
 15 2015) based on the estimates using the 2006 GLs, but the CH₄ emissions overlapped with the
 16 estimates using the 1996 GLs because of the low consumption level. The trend in N₂O emissions in
 17 2008–2017 (based on already available 1996 GLs estimates) overlays the origin line as the value
 18 reported in the 2008, 2012, 2015 and 2017 summary tables is zero. This is simply because the
 19 UNFCCC software rounds off decimals when generating the summary tables. It means that the value
 20 is not necessarily zero, but very close to zero. The same trend can also be seen in the manufacturing
 21 and transport sectors; however, the reason for this is the same as before. In 1994, the fuel
 22 consumption for petroleum refining was very small, the reason for the 1994 N₂O emission value being
 23 the same for the estimates using both GLs. In summary, it is appropriate to mention here that the

1 emission factors of the energy industries were the same for both GLs, and that the marginal difference
 2 arose because of the inclusion of additional source category in the energy industries.



3
 4 **Fig. 4. Difference in GHG emissions in the energy industries in Pakistan during 1994–2017,**
 5 **2006 vs. 1996 GLs**

6 **4.2.2.2 Manufacturing industries**

7 Figure 5 shows the emissions gap in the energy sector for the manufacturing industries. The
 8 manufacturing industries include the fuel used in Pakistan steel mills, the cement industry, the
 9 fertilizer industry, and other general industries. Production of steel and cement in Pakistan is believed
 10 to use coking coal and other bituminous coal, imported mostly from other countries. The default
 11 emission factor in both GLs is the same for coking coal and other bituminous coal. However, it is
 12 assumed that Pakistan's brick kiln industry uses sub-bituminous coal, which has a higher emission
 13 factor compared with coking coal and other bituminous coal. For the estimates of both GLs, the
 14 difference between CO₂ and CO₂-eq is not significant. However, the difference is visible for both CH₄
 15 and N₂O. The CH₄ emissions by 1996 GLs were higher than the 2006 GLs because of higher CH₄
 16 emission factors for oil, gas, and coal defined in 1996 GLs. The N₂O emissions obtained with the
 17 2006 GLs are more than those estimated with the 1996 GLs because of the higher emission factor for
 18 coal (other bituminous and sub-bituminous) in the 2006 GLs. There is no difference in the fuel
 19 consumption data that were used to calculate emissions following both GLs.





Fig. 5. Difference in GHG emissions in the manufacturing industries in Pakistan during 1994–2017, 2006 vs. 1996 GLs

4.2.2.3 Transport sector

Figure 6 shows the difference in GHG emissions between the 2006 and 1996 GLs in the transport sector. The trends for CO₂-eq and CO₂ overlay, with the exception of the year 2012, in which the estimates with the 1996 GLs are higher than those with the 2006 GLs. The reason for the different estimate for the year 2012 is that the value for diesel consumption used in the 1996 GL based inventory for 2012 for the mobile agricultural/forestry/fishing sector was taken from a reference energy scenario generated by the Pakistan Integrated Energy Model (Pak-IEM, 2010). This value was projected on the basis of the model rather than being the actual value based on national statistics, which, however, is lower than the projected value. This discrepancy would explain the difference in total emissions for 2012. The difference between CH₄ and N₂O emissions is attributable to the fact that the 2006 GLs provide updated fuel-specific gasoline, diesel, and natural gas emission factors that are higher than those estimated using the 1996 GLs. That is why the 2006 GLs based emission quantities of CH₄ and N₂O from transport sector are higher compared to 1996 GLs.

1
2




3
4
5
6

Fig. 6. Difference in GHG emissions in the transport sector in Pakistan during 1994–2017, 2006 vs. 1996 GLs

4.2.2.4 Other sectors

7 Figure 7 shows the emission gap for other sectors including residential, commercial/institutional, and
 8 agriculture/forestry/fishing between the 2006 and 1996 GLs. The trend for all GHGs and CO₂
 9 emissions are quite similar except for the year 2012 in which the estimates for the 1996 GLs are
 10 higher than the estimates for 2006 GLs. The reason for this is the use of the scenario value modeled
 11 by Pakistan Integrated Energy Model (Pak-IEM, 2010) for diesel consumption in agricultural
 12 transport in 2012. This modeled diesel value is higher than the actual value based on national
 13 statistics, with the result being the gap in total emissions for 2012. Nevertheless, the CH₄ and N₂O
 14 emissions estimated according to the 2006 GLs exhibit flat patterns, despite some variations in the
 15 estimates based on the 1996 GLs (Fig. 7b, 7c). The reason for the difference is that the value of the
 16 1994 emission inventory for these two GHGs was not available; therefore, the values are based on
 17 expert judgment. Furthermore, the default CH₄ and N₂O emission factors in the category of
 18 agriculture/forestry/fishing (for mobile combustion only) were updated in 2006 GLs (CH₄ 4.15 kg
 19 (TJ)⁻¹; N₂O 28.6 kg (TJ)⁻¹ compared to the 1996 GLs (CH₄ 5 kg (TJ)⁻¹; N₂O 0.6 kg (TJ)⁻¹).

1
2




3
4
5
6

Fig. 7. Difference in GHG emissions in other sectors (residential, commercial, and agriculture) of energy sector in Pakistan during 1994–2017, 2006 vs. 1996 GLs

4.2.2.5 Fugitive emissions


7 Figure 8 shows the fugitive CH₄ emissions and the considerable difference between the estimates
8 based on the 2006 GLs and 1996 GLs. This difference can be attributed to the application of very high
9 emission factors taken from the 1996 GLs for natural gas production and distribution. In the 2006
10 GLs, the emission factors were revised and have a reduced range compared to the 1996
11 GLs. This clarifies why the estimates with the 2006 GLs have lower emission values than those with the 1996
12 GLs.



1
2 **Fig. 8. Difference in fugitive GHG emissions in Pakistan during 1994–2017, 2006 vs. 1996 GLs**
3 **4.2.3 IPPU sector overall and sub-sectoral differences**

Journal Pre-proof

1 Figure 9 shows the difference between the 2006 GLs and 1996 GLs for GHG emissions of the IPPU
 2 sector. The difference is negligible for the mineral industry, which is mostly concerned with cement
 3 production. The data on cement production was taken from Pakistan Economic Surveys and all
 4 cement production is presumed to be mainly that associate with Portland cement in Pakistan.
 5 According to the 2006 GLs, a default value of 95% is used to indicate the clinker fraction in cement.
 6 However, a difference exists for the chemical industry because of the use of different methodologies
 7 of both GLs. The estimates of the 1996 GLs use the urea production to calculate the amount of CO₂,
 8 whereas the 2006 GLs use the amount of ammonia that was produced. Similarly, soda ash emissions
 9 are reported in the chemical industry, according to the 2006 GLs, whereas in previous inventories the
 10 emissions were considered under the mineral industry. In terms of the metal industry (iron and steel
 11 production), the CO₂ emissions were estimated as zero in 2017. This is because the Pakistan steel
 12 industry was shut down in 2017, resulting in no coke or pig iron production according to national
 13 statistics. The trend for the metal industry also shows that iron and steel production in Pakistan has
 14 continued to decline since 1994.



15

16 **Fig. 9. IPPU sector overall gap of CO₂ emissions in Pakistan during 1994–2017, 2006 vs. 1996**
 17 **GLs**

18 **5. Conclusions**

19 The study found that the 2006 GLs based total CO₂ equivalent emissions from the energy and IPPU
 20 sectors show a relative difference of 4% and –1% compared to those based on 1996 GLs, respectively.
 21 In the energy sector, with an average annual growth rate of 3.6%, CO₂ is the most abundant GHG
 22 released into the atmosphere followed by CH₄ and N₂O. Although the gap in energy sector emissions
 23 is notable for CH₄ and N₂O, the overall CO₂ equivalent emissions based on the 2006 GLs

1 replicate those reported by using the 1996 GLs. This is because the CO₂ emissions that are dominant,
2 show similar amounts over the time series. Further, the fugitive emissions under energy sector
3 represent significant gap between the two Guidelines. For the IPPU sector, the gap is relatively large
4 in the chemical industry, but the difference in total GHG emissions is marginal between both GLs.
5 However, the 2006 GLs endeavor to upgrade the accuracy and reliability of GHG inventories as a
6 result of incorporating new sources and revised emission factors. For that reason, the GHG
7 inventories by 2006 GLs is quite different from that by 1996 GLs. It is therefore recommended that
8 Pakistan implement the 2006 GLs to improve the GHG inventories, as this has rarely been adopted by
9 developing countries. This would also help to better identify the national targets for GHG reduction
10 with respect to NDCs.

11 The results of the study are limited to Tier 1 method (using default emission factors) of IPCC GLs due
12 to the unavailability of country-specific emission factor data. Although decision tree in 2006 GLs for
13 estimating GHG emissions clearly recommends using higher tiers for key categories (key category
14 analysis is described in section 3 of part II of this paper), most developing countries suffer from a lack
15 of resources and experts, forcing to adopt Tier 1 method. In the long run, Pakistan should devote to
16 develop Tiers 2 and 3 method reflecting country- and plant-specific emission characteristics. For the
17 time being, Tier 1 method has been applied broadly despite of high uncertainty. In general, the
18 uncertainties in the activity data are smaller as it is compiled and frequently reported by national
19 statistical agencies, which may have already identified the uncertainties associated with data as part of
20 their data collection procedures.

21 It should be noted that Pakistan has a well-managed system to compile and publish the national
22 energy statistics annually. However, it must be made more dynamic by collecting the data in a format
23 consistent with the GHG inventories requirement. This will further improve the data consistency and
24 reduce data conversion errors, such as unit conversion. Information on the country-specific fuel
25 properties such as carbon content, carbon oxidation factor, and fuel energy content – NCV is
26 recommended to be reported in national energy statistics publication. Further, efforts should be made
27 for road transport (a key category) to collect distance travelled data of vehicles by type and fuel to
28 reflect true vehicle emissions under a higher Tier approach. The cement industry and the ammonia
29 production has appeared to be the key categories in the IPPU sector. It is therefore recommended that
30 the data concerning clinker production or the use of carbonate be made available and reported as part
31 of national industry statistics for the cement industry. For the ammonia industry, instead of estimating
32 emissions based on the ammonia production data, efforts should be made to get details on the
33 total fuel demand for ammonia production. It would undoubtedly benefit Pakistan to achieve more
34 reliable and accurate estimates of GHG emissions from energy and IPPU sector under a higher Tier.

35 **Conflict of interest**

36 The authors declare no conflict of competing interest.

37 **Acknowledgments**

38 We sincerely acknowledge the Global Change Impact Studies Centre (Ministry of Climate Change,
39 Government of Pakistan) and Applied Systems Analysis Division (Pakistan Atomic Energy
40 Commission, Government of Pakistan) for making available the data of previous GHG emissions
41 inventories for Pakistan. We also thank the editors and three anonymous reviewers for their careful
42 reading of the manuscript and insightful comments and suggestions.

43 **Appendix A.**

44 Table A1. GHG emissions and percentage share of the IPPU sector of Pakistan (1994–2017) using the
45 2006 GLs

Journal Pre-proof


1 References

- 2 ASAD (Applied Systems Analysis Division, Pakistan Atomic Energy Commission, Government of
3 Pakistan), 2009. Greenhouse Gas Emission Inventory of Pakistan for the Year 2007–2008.
- 4 ASAD, 2016. Revised Greenhouse Gas Emission Inventory of Pakistan for the Year 2007–2008.
- 5 Bergamaschi, P., A. Danila, R. F. Weiss, et al., 2018. Atmospheric monitoring and inverse modelling
6 for verification of greenhouse gas inventories. EUR 29276 EN, Publications Office of the
7 European Union, Luxembourg, ISBN 978-92-79-88938-7, doi:10.2760/759928, JRC111789.
- 8 Breidenich, C., 2011. Improving reporting of national communications and GHG inventories by non-
9 annex I parties under the climate convention. Natural Resources Defense Council, White Paper.
- 10 FAO (Food and Agriculture Organization of the United Nations), 2015. Global Forest Resource
11 Assessment 2015 – Country Report (Pakistan).
- 12 GCISC (Global Change Impact Studies Centre, Ministry of Climate Change, Government of
13 Pakistan), 2016. Greenhouse Gas Emission Inventory of Pakistan for the Year 2011–2012.
14 GCISC-RR-19.
- 15 GCISC, 2017. Greenhouse Gas Emission Inventory of Pakistan for the Year 2014–2015. GCISC-RR-
16 20.
- 17 HDIP (Hydrocarbon Development Institute of Pakistan, Ministry of Energy, Government of
18 Pakistan), 1994. Pakistan Energy Yearbook 1994.
- 19 HDIP, 2012. Pakistan Energy Yearbook 2012.
- 20 HDIP, 2015. Pakistan Energy Yearbook 2015.
- 21 HDIP, 2017. Pakistan Energy Yearbook 2017.
- 22 HDIP, 2008. Pakistan Energy Yearbook 2008.
- 23 IPCC, 1996. Climate Change 1995: The Science of Climate Change: Contribution of Working Group
24 I to the Second Assessment Report of the Intergovernmental Panel on Climate Change.
25 Cambridge University Press, Cambridge.
- 26 IPCC, 1997. Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Cambridge
27 University Press, Cambridge.
- 28 IPCC, 2001. Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the
29 Third Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge
30 University Press, Cambridge and New York.
- 31 IPCC, 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Cambridge University
32 Press, Cambridge.
- 33 IPCC, 2007. Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to
34 the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge
35 University Press, Cambridge.
- 36 Mir, K. A., Purohit, P., Mehmood, S., 2017. Sectoral assessment of greenhouse gas emissions in
37 Pakistan. *Environ Sci Pollut Res* 24, 27345-27355. <https://doi.org/10.1007/s11356-017-0354-y>
- 38 MoF (Ministry of Finance, Government of Pakistan), 2008. Pakistan Economic Survey 2007-08.
39 Retrieved from http://www.finance.gov.pk/survey_0708.html


- 1 MoF, 2012. Pakistan Economic Survey 2011-12. Retrieved from
2 http://www.finance.gov.pk/survey_1112.html
- 3 MoF, 2015. Pakistan Economic Survey 2014-15. Retrieved from
4 http://www.finance.gov.pk/survey_1415.html
- 5 MoF, 2017. Pakistan Economic Survey 2016-17. Retrieved from
6 http://www.finance.gov.pk/survey_1617.html
- 7 MoIP (Ministry of Industries and Production, Government of Pakistan), 2017. Year Book 2016-2017.
8 Retrieved from <http://www.moip.gov.pk/frmDetails.aspx>
- 9 MoNFSR (Ministry of National Food Security & Research, Government of Pakistan), 2019.
10 Agricultural Statistics of Pakistan 2016-17. Retrieved from
11 <http://www.mnfsr.gov.pk/frmDetails.aspx>
- 12 Pak-IEM (International Resources Group for Asian Development Bank and Ministry of Planning and
13 Development, Government of Pakistan), 2010. Pakistan Integrated Energy Model (Pak-IEM)
14 Final Report – Volume II.
- 15 UN (United Nations), 1992. United Nations Framework Convention on Climate Change.
- 16 UNFCCC (United Nations Framework Convention on Climate Change), 2003a. Pakistan’s Initial
17 National Communication on Climate Change.
- 18 UNFCCC, 2003b. Report of the Conference of the Parties on its Eighth Session, held at New Delhi
19 from 23 October to 1 November 2002. FCCC/CP/2002/7/Add.2, Decision 17/CP.8.
- 20 UNFCCC, 2014. Report of the Conference of the Parties on its Nineteenth Session, held in Warsaw
21 from 11 to 23 November 2013. FCCC/CP/2013/10/Add.3, Decision 24/CP.19.
- 22 UNFCCC, 2015. Report of the Conference of the Parties on its twenty-first session, held in Paris from
23 30 November to 13 December 2015. FCCC/CP/2015/10/Add.1, Decision 1/CP.21.
- 24 UNFCCC, 2016. Pakistan’s Intended Nationally Determined Contribution.
- 25 UNFCCC, 2019. Pakistan’s Second National Communication on Climate Change.
- 26 Yona, L., Cashore, B., Jackson, R. B., et al., 2020. Refining national greenhouse gas inventories.
27 *Ambio*, 1-6. <https://doi.org/10.1007/s13280-019-01312-9>
- 28 Zhu, S.-L., Wang, W.-T., 2013. Development of national greenhouse gas emissions inventories in the
29 context of international climate negotiations. *Adv. Clim. Change Res.* 4(1), 60-68.
30 <https://doi.org/10.3724/SP.J.1248.2013.060>


31

(a)




(b)







(a)




(b)





(c)





(d)




















Conflict of Interest

The authors declare no conflict of interest.

Journal Pre-proof