


Journal Pre-proof

Radiative effects of reduced aerosol emissions during the COVID-19 pandemic and the future recovery

Stephanie Fiedler, Klaus Wyser, Joeri Rogelj, Twan van Noije



PII: S0169-8095(21)00422-1

DOI: <https://doi.org/10.1016/j.atmosres.2021.105866>

Reference: ATMOS 105866

To appear in: *Atmospheric Research*

Please cite this article as: S. Fiedler, K. Wyser, J. Rogelj, et al., Radiative effects of reduced aerosol emissions during the COVID-19 pandemic and the future recovery, *Atmospheric Research* (2021), <https://doi.org/10.1016/j.atmosres.2021.105866>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier B.V.

Radiative effects of reduced aerosol emissions during the COVID-19 pandemic and the future recovery

Stephanie Fiedler^{a,b,*}, Klaus Wyser^c, Joeri Rogelj^{d,e}, Twan van Noije^f

^a*University of Cologne, Institute of Geophysics and Meteorology, Cologne, Germany*

^b*Hans-Ertel-Centre for Weather Research, Climate Monitoring and Diagnostics, Bonn/Cologne, Germany*

^c*Rosby Centre, Swedish Meteorological and Hydrological Institute, Sweden*

^d*Grantham Institute, Imperial College London, United Kingdom*

^e*International Institute for Applied Systems Analysis, Laxenburg, Austria*

^f*Royal Netherlands Meteorological Institute, De Bilt, Netherlands*

Abstract

The pandemic in 2020 caused an abrupt change in the emission of anthropogenic aerosols and their precursors. We estimate the associated change in the aerosol radiative forcing at the top of the atmosphere and the surface. To that end, we perform new simulations with the CMIP6 global climate model EC-Earth3. The simulations use the here newly created data for the anthropogenic aerosol optical properties and an associated effect on clouds from the simple plumes parameterization, based on revised SO₂ and NH₃ emission scenarios. Our results highlight the small impact of the pandemic on the global aerosol radiative forcing in 2020 compared to the CMIP6 scenario SSP2-4.5 of the order of +0.04 Wm⁻², which is small compared to the natural year-to-year variability in the radiation budget. Natural variability also limits the ability to detect a meaningful regional difference in the anthropogenic aerosol radiative effects. We identify the best chances to find a significant change in radiation at the surface during cloud-free conditions for regions that were strongly polluted in the past years. The post-pandemic recovery scenarios indicate a spread in the aerosol forcing of -0.68 to -0.38 Wm⁻² for 2050 relative to the pre-industrial, which translates to a difference of +0.05 to -0.25 Wm⁻² compared to the 2050 baseline from SSP2-

*Corresponding author

Email address: stephanie.fiedler@uni-koeln.de (Stephanie Fiedler)

4.5. This spread falls within the present-day uncertainty in aerosol radiative forcing and the CMIP6 spread in aerosol forcing at the end of the 21st century. We release the new MACv2-SP data for studies on the climate response to the pandemic and the recovery scenarios. Our 2050 forcing estimates suggest that sustained aerosol emission reductions during the post-pandemic recovery cause a stronger climate response than in 2020, i.e., there is a delayed influence of the pandemic on climate.

Keywords: anthropogenic aerosols, radiative forcing, CC v1.1.19 pandemic, scenarios, CMIP6, CovidMIP

1. Introduction

The aerosol burden in 2020 is affected by reduced emissions of anthropogenic aerosols and their precursors associated with the global COVID-19 pandemic. Many countries have witnessed a reduction in socio-economic activities and lockdowns. The associated decline in traffic and industrial productivity have led to marked regional reductions in atmospheric pollution improving the air quality (e.g., van Heerwaarden et al., 2021; Ranjan et al., 2020). Figure 1 illustrates the observed anomaly in the mid-visible aerosol optical depth, τ' , for northern hemisphere spring in 2020, calculated as the difference against the 20-year spring climatology from NASA's MODIS satellite product MOD08_M3 v6.1 (Acker and Leptoukh, 2007; Fatnick et al., 2015), assessed in more detail by Sanap (2021). Pronounced negative τ' in spring 2020 are identified for Eastern and Southern Asia as well as the Northwest Pacific. Even Europe and North America, where the mean anthropogenic aerosol burden is relatively small, have seen less aerosols during spring 2020. Albeit an influence of negative regional trends in anthropogenic aerosol emissions due to legislation (e.g., Sun et al., 2018; Chu et al., 2020; Cherian and Quaas, 2020), and variability in aerosol emissions from natural sources (e.g., Riemer et al.; Song et al., 2021), the pandemic is a clear contributor to τ' , consistent with other findings. For instance, the pandemic restrictions in most of the European countries occurs in parallel with excep-




Figure 1: Observed anomaly in aerosol optical depth for northern hemisphere spring 2020. Shown is the anomaly in the aerosol optical depth (τ') at 550 nm in March–May 2020 against the climatology of the same months for 2000–2020 from MODIS.

tionally blue skies and new extremes in surface irradiance (van Heerwaarden et al., 2021). Improved air quality associated with COVID-19 lockdowns were also documented for India (Srivastava et al., 2021; Zhang et al., 2021), Iran (Broomandi et al., 2020), and the United States (Berman and Ebisu, 2020).

25 The emission reductions due to the COVID-19 pandemic are thought to potentially influence climate (e.g., Forster et al., 2020a; Gettelman et al.; Yang et al., 2020), but the emission reductions are not considered in the contemporary climate simulations of the Coupled Model Intercomparison Project phase 6 (CMIP6, Eyring et al., 2016), used for assessing climate changes by the Intergovernmental Panel on Climate Change (IPCC). The construction of the CMIP6
 30 scenarios of anthropogenic emissions has therefore been revisited (Forster et al., 2020b). Based on the new emission data, the scientific community plans to investigate the co-occurrence of climate anomalies and the COVID-19 impacts on global air quality. To this end, a new climate model inter-comparison project
 35 for 2015–2050 with the revised emission scenarios has been proposed (Covid-MIP, Lamboll et al., 2021) under the umbrella of the Detection and Attribution Model Intercomparison Project (DAMIP, Gillett et al., 2016) endorsed by CMIP6. Nine CMIP6 models in the sixth assessment report of the IPCC (IPCC, 2021) use the simple plumes aerosol parameterization MACv2-SP (Fiedler et al.,

40 2017; Stevens et al., 2017). Some of them like EC-Earth3 and MPI-ESM1.2 participate in CovidMIP and therefore need MACv2-SP input data consistent with the new emission data of CovidMIP. We here derive these new input data for MACv2-SP and provide it for use in climate studies.

The aim of the present study is to estimate the impact of the COVID-19 pandemic on the radiative forcing of anthropogenic aerosols for the pandemic year 45 2020 and the new recovery scenarios in 2050. A first estimate of the radiative forcing associated with aerosol-radiation interaction from a complex aerosol-climate model has been made for spring 2020 (Weber et al., 2020). Here, we provide the first estimates of the effective radiative forcing for the entire years 50 2020 and 2050 from MACv2-SP that account for both aerosol-radiation and aerosol-cloud interactions as well as the large impact of the variability in rapid adjustments. To this end, we derive and describe the new MACv2-SP data for the anthropogenic aerosol optical properties and an associated effect on clouds for 2015–2050 from the revised aerosol emissions (Forster et al., 2020b). These 55 emission data cover different recovery pathways after the pandemic ranging from fossil-fuel based to green developments into the future. We use the here newly constructed MACv2-SP data in the CMIP6 model EC-Earth3 (Döscher et al., 2021), which uses MACv2-SP as standard to represent anthropogenic aerosols. EC-Earth3 simulates aerosol-radiation and aerosol-cloud interactions including 60 both the Twomey effect and cloud adjustments with MACv2-SP. We perform new atmosphere-only experiments with EC-Earth3 and estimate the effective radiative forcing (ERF) of the anthropogenic aerosols in 2020 and 2050 at the top of the atmosphere and the surface. Details of our methods are given in Section 2, followed by our results in Section 3, and a discussion with conclusions in 65 Section 4.

2. Methods

2.1. Emissions of SO_2 and NH_3

Forster et al. (2020b) developed five new scenarios to explore the impact of

the COVID-19 pandemic on current and future anthropogenic emissions. These

70 scenarios are:

- A baseline scenario (*base*) without any impact of the COVID-19 pandemic and the measures to contain it;
- A two-year-blip scenario (*blp*) that assumes emissions return to the baseline scenario after a two-year reduction in emissions due to temporary societal lockdowns and disruptions;
- 75 • A fossil-fuel recovery scenario (*ff*) that assures the recovery from the COVID-19 economic downturn preferentially supports polluting fossil-fuel-based economic sectors;
- A moderate and strong green recovery scenario (called *mg* and *sg*, respectively)
- 80 • (respectively) that assume different levels of preferential stimulus of green sectors during the recovery from COVID-19 (e.g., see Andrijevic et al., 2020) that would lead to limiting global mean temperature increase relative to preindustrial levels to well below $2^{\circ}C$ and to $1.5^{\circ}C$, respectively (Forster et al., 2020b).

85 The near-term evolution of SO_2 and NH_3 during the COVID-19 lockdown period uses the activity scaling method of Forster et al. (2020b). For the extensions beyond the COVID-19 lockdown period and until 2050, the large-scale global relationships between greenhouse gases, aerosols and aerosol precursors as found in detailed emissions scenarios derived with integrated assessment models is used

90 (Lamboll et al., 2020b). That is the emission data produced with the python package Silicone v1.0.0 for inferring emission data of multiple chemical species. It allows to fill missing data of one species based on reports on emissions of other species, e.g., N_2O projections for a given CO_2 emission using the relationship of N_2O and CO_2 from another scenario. Emissions evolutions of SO_2 and NH_3

95 compatible with each of the above scenarios have been estimated, based on the relationships found in the scenario ensemble compiled and assessed as part of the Intergovernmental Panel on Climate Change's (IPCC) Special Report on

Global Warming of 1.5°C (Rogelj et al., 2018; Huppmann et al., 2018). The dataset defines eight emission sectors, namely agriculture, energy, waste, transportation, international shipping, and solvent production and application as well as industrial, residential, and commercial sectors plus a remaining fraction classified as other.

For illustrating the difference in emissions due to the pandemic in 2020, we show the difference in the emission sum of SO_2 and NH_3 emissions from all sectors for 2020 for the *blp* scenario against *base* (Figure 2). The reduction in the emissions of SO_2 plus NH_3 due to the pandemic is globally visible, with particularly pronounced impacts on the emissions in Asia and the Pacific coast of South America measured by the absolute differences (Figure 2a). Emission changes for 2020 associated with the pandemic translate to regional differences by up to -37% at the Western coast of South America and mostly -14% over the oceans relative to *base*, while reductions in Asian countries are at the order of a few percent (Figure 2b). In the global area-weighted mean, the reduction of the SO_2 plus NH_3 emission sum for 2020 is -11% in *blp* against *base*. The reduction in SO_2 emissions (-14%) is slightly stronger than for NH_3 emissions (-10%). The percentage reductions in SO_2 and NH_3 emissions per sector are similar. Strongest reductions in the 2020 emission sums are associated with transportation (-20%), international shipping (-14%) and industrial sources (-14%) in *blp* relative to *base*. There are no differences for agriculture, waste, and solvents production and applications. Emissions of SO_2 and NH_3 are used to construct the MACv2-SP data in the next Section.

2.2. Anthropogenic aerosol parameterization

We use the emissions of SO_2 and NH_3 from Forster et al. (2020b) to create the new input data for the novel simple-plumes parameterization MACv2-SP (Fiedler et al., 2017; Stevens et al., 2017) in use for representing anthropogenic aerosol effects in climate models of CMIP6 (e.g., Fiedler et al., 2019a; Mauritsen et al., 2019). MACv2-SP prescribes month-to-month and year-to-year changes of the three dimensional fields of anthropogenic aerosol optical prop-




Figure 2: Aerosol emission difference for 2020 due to pandemic. Shown are the reduction in the emission sum of SO_2 and NH_3 for the *blp* scenario against *base* (a) as absolute difference in gm^{-2} per year and (b) as percentage change per year relative to *base*. All values are based on sums for the year 2020 over all emission sectors.




Figure 3: MACv2-SP τ_a in 2050. Shown is the annual mean in anthropogenic aerosol optical depth (τ_a) at 550 nm for the (a) fossil-fuel based, (b) moderate green, and (c) strong green scenario. Grey symbols mark the centers of the aerosol plumes associated with industrial pollution (circles) and emissions from both industry and biomass burning (rectangles).

erties and associated effects on clouds. Temporal changes from 1850 to 2100 have been derived by scaling the anthropogenic aerosol optical depth of 2005 with the CMIP6 emission amounts of SO_2 and NH_3 (Stevens et al., 2017; Fiedler et al., 2019b). Emissions of SO_2 and NH_3 are chosen for consistency with earlier data constructions for MACv2-SP, motivated by the assumption that temporal changes in other anthropogenic aerosol species correlate with them (Stevens et al., 2017), e.g., NO_x . Large-scale co-variability in NO_x , SO_2 , and NH_3 is for instance also found for the emission data used here (Forster et al., 2020a). Evaluating the MACv2-SP approach against the aerosol optical depth from a complex aerosol-climate model indicates similar regional scaling behaviours (Fiedler et al., 2019b). Moreover, our MACv2-SP data in CovidMIP models generate results that fall within the spread from aerosol-climate model simulations with more complex aerosol treatments (Jones et al., 2021). A complete technical description of MACv2-SP is given by Stevens et al. (2017). Again previous data for MACv2-SP did not account for the effect of the pandemic on anthropogenic emissions. We therefore create here new MACv2-SP input data, based on the emission data sets for 2015–2050 accounting for the COVID-19 pandemic and the four recovery scenarios from Forster et al. (2020b).

For creating the new MACv2-SP data, we scale the anthropogenic aerosol optical depth and the effect on clouds from 2005 to other years by multiplying scaling factors in the centre of the nine plumes, marked in Figure 3a. Mathe-

mathematical functions in MACv2-SP use the values in the plume centers to create the
 150 three dimensional distribution of the aerosol extinction. As an example, Figure
 3 shows the maps of the mid-visible anthropogenic aerosol optical depth in 2050
 from MACv2-SP for three scenarios that we derive here. Technically, we create
 annual scaling factors for each plume center and each year to be represented in
 MACv2-SP.

155 Our method for constructing the scaling factors for MACv2-SP is similar to
 the method for the CMIP6 scenarios (Fiedler et al., 2019b). The anthropogenic
 aerosol optical depth τ_i in each plume center $i = 1, \dots, 9$ is scaled with the
 emission scaling factor E_i for the years t with:

$$\tau_i(t) = E_i(t)\tau_i(2005) \quad (1)$$

We use $\tau_i(2005)$ at 550 nm from Stevens et al. (2017). The scaling factors $E_i(t)$
 160 are constructed from the anthropogenic emission ϵ_{ik} of the species k . These are
 the gridded emission data for SO_2 and NH_3 from the emission data version 4
 (Forster et al., 2020b). These emission data are provided on a longitude-latitude
 grid with a resolution of $0.5^\circ \times 0.7^\circ$ (Figure 2). The calculation of $E_i(t)$ follows
 Fiedler et al. (2019a):

$$E_i(t) = \frac{\sum_{k=1,2} w_k [\epsilon_{ik}(t) - \epsilon_{ik}(1850)]}{\sum_{k=1,2} w_k [\epsilon_{ik}(2005) - \epsilon_{ik}(1850)]} \quad (2)$$

165 The emissions of 1850 and 2005 are taken from the CMIP6 historical emission
 data. We consider emissions from all anthropogenic sectors provided by Forster
 et al. (2020b), and include open burning emissions from the CMIP6 scenario
 SSP2-4.5. SSP2-4.5 is the baseline for the experiments to be carried out in
 CovidMIP (Lamboll et al., 2021). The anthropogenic emissions ϵ_{ik} are inte-
 170 grated values over the 10×10 grid boxes surrounding the plume center. The
 weights w_k for the two species are $w_1 = 0.645$ for SO_2 and $w_2 = 0.355$ for NH_3 ,
 representing the forcing ratio of sulphate against ammonium for present day
 (Stevens et al., 2017).

Figure 4 shows examples of the scaling factors for each of the nine aerosol
 175 plume centers i . Contrasting the baseline against the two-year-blip data illus-




Figure 4: Example of the scaling factors for the aerosol optical depth in the centre of the aerosol plumes for two scenarios. Shown are the annual scaling factors for the color-coded plumes calculated from the (solid) baseline and (dashed) two-year-blip emissions for the years provided in the data. The pandemic period 2019–2023 is marked with grey shading.

trates the reduction of E_i for 2020–2021. The reduction for E_i is particularly strong for the plumes over India and South America. These are consistent with emission reductions in anthropogenic aerosols and their precursors in many regions across Asia for 2020 (Figure 2). In South America, the reductions are particularly strong near the Pacific coast. Comparably smaller changes are seen in Europe, Africa and North America, consistent with relatively less emission reductions.

MACv2-SP typically uses one $E_i(t)$ per decade with linear interpolation in between. This is the same here, except that we also construct the scaling factors for the individual years around 2020 for consistency with the new emission data considering the pandemic. We note that the observed small-scale structures in the aerosol burden, e.g., like in local observations, cannot be created with MACv2-SP owing to the design and purpose of this parameterization (Stevens et al., 2017). MACv2-SP is a fast and flexible parameterization of anthropogenic aerosols to represent their effects on climate. As such results from using these individual years in later studies should be interpreted as estimates for the large-

scale influence and not as local constraints of the forcing for individual years.

Aerosol absorption is prescribed with the single scattering albedo of $\omega_0 = 0.93$ for industrial plumes and $\omega_0 = 0.87$ for plumes additionally affected by biomass burning, marked in Figure 3a, accounting for regional differences in the aerosol mixture. The asymmetry parameter, $\gamma = 0.63$, is constant. MACv2-SP uses the Angstrom exponent $\alpha = 2$ to interpolate the aerosol optical properties for different wavelengths.

Additionally to the aerosol optical properties, MACv2-SP prescribes aerosol effects on the cloud droplet number concentrations N . The latter is induced with the prefactor η_N to be multiplied with N in the host model:

$$\eta_N = 1 + \frac{dN}{N} = \frac{\ln[1000(\tau_a(\phi, \lambda, t) + \gamma(\phi, \lambda, t)) + 1]}{\ln[1000\tau_b(\phi, \lambda, t) + 1]} \quad (3)$$

The background aerosol optical depth (τ_b) is a simplified representation that follows the plume structure for τ_a to parameterize the aerosol effect on clouds. Host models can multiply η_N with N in the radiation transfer calculation to induce a Twomey effect only, e.g., in MPI-ESM1.2, or in the cloud microphysics to allow further rapid adjustments of clouds, e.g., in EC-Earth3 used here. Additional documentation of MACv2-SP and details on the application in climate studies is given elsewhere (e.g., Fiedler et al., 2017; Stevens et al., 2017; Fiedler et al., 2019a).

The new MACv2-SP input data are provided as supplementary material for use in climate studies, e.g., in CovidMIP (Lamboll et al., 2021). It covers the baseline and scenarios that assume recoveries after the COVID-19 pandemic that intensify the use of fossil fuels, follow a moderate or strong green pathway, and return to a business as usual pathway after the assumed two-year interruption by the pandemic in 2020 and 2021.

2.3. Model experiment design

We estimate the ERF of the anthropogenic aerosol reduction in 2020 and the ERF spread associated with the recovery scenarios in 2050 from the new data. To this end, we perform atmosphere-only simulations with EC-Earth3 and

220 compute the ERF of the anthropogenic aerosols at the top of the atmosphere and the surface, i.e., the instantaneous radiative effects plus the rapid adjustments in the atmosphere.

EC-Earth3 is an Earth system model participating in CMIP6 (Döscher et al., 2021). EC-Earth consists of an atmosphere and land surface model adapted
225 from cycle 36r4 of the European Centre for Medium-Range Weather Forecasts (ECMWF) Integrated Forecasting System (IFS), documented online (ECMWF, 2010a,b, last accessed 4 Jan 2021), coupled to the physical ocean and sea-ice model from NEMO3.6 (Madec and the NEMO team, 2016), and a river runoff model based on CaMa-Flood v3.6.2 (Yamazaki et al., 2013). The model grid
230 chosen here has a horizontal resolution of about 80 km and 91 vertical levels up to 0.01 hPa.

The implementation of MACv2-SP in EC-Earth3 is such that the model accounts for aerosol-radiation interactions and aerosol-cloud interactions, including Twomey and cloud lifetime effects (e.g., Fiedler et al., 2019a). Natural
235 tropospheric aerosols are prescribed with a monthly climatology of pre-industrial aerosol optical properties and concentrations representative for the year 1850. This climatology was produced in an offline simulation of TM5, the atmospheric chemistry and aerosol component of EC-Earth3-AerChem (van Noije et al., 2021). TM5 includes contributions from sulphate, black carbon, primary and
240 secondary organics, sea salt, mineral dust, ammonium, nitrate and methane sulfonic acid. Shortwave radiative effects of tropospheric aerosols are determined by the combined pre-industrial and anthropogenic contributions to the aerosol extinction, absorption (single-scattering albedo), and asymmetry factor; their longwave radiative effects are calculated based on the pre-industrial mass con-
245 centrations of different aerosol species. Cloud droplet number concentrations are calculated from the pre-industrial aerosol number and mass concentrations following the activation scheme from (Abdul-Razzak and Ghan, 2000), and modified by the anthropogenic enhancement factor provided by MACv2-SP. Both the cloud droplet effective radius and the rate of autoconversion of cloud liquid
250 to rain depend on the cloud droplet number concentration (van Noije et al.,

2021). Consequently, the anthropogenic aerosol forcing in EC-Earth3 includes a contribution from the cloud lifetime or second indirect effect.

Our simulations use annually repeating aerosol optical properties and associated effects on clouds for the years 2020 and 2050. We run experiment with
 255 the same setup except for these changes in the anthropogenic aerosols:

- *base*: new MACv2-SP properties from the baseline for the year 2020 (*2020-base*) and 2050 (*2050-base*).
- *2020-blp*: new MACv2-SP properties from the two year-blip for 2020,
- *2050*: new MACv2-SP properties for 2050 from the recovery scenarios
 260 that are either primarily fossil-fuel based (*2050-ff*), moderate (*2050-mg*) or strong green (*2050-sg*)
- *pi*: without anthropogenic aerosol effects for calculating the ERF in 2020 and 2050 relative to the pre-industrial.

Our setup of these experiments follows Wyser et al. (2020). These are pre-
 265 industrial atmosphere-only experiments, as typical for effective radiative forcing calculations from contemporary climate model experiments (Pincus et al., 2016; Fiedler et al., 2019; Smith et al., 2020). We prescribe annually repeating pre-industrial boundary conditions like for a *piClim-control* experiment in the Radiative Forcing Model Inter-comparison project (RFMIP, Pincus et al.,
 270 2016), i.e., monthly climatology for sea-surface temperatures and sea ice, here calculated from 120 years of the model’s pre-industrial control experiment for CMIP6. All our experiments are run for 55 years. The first five simulation years are discarded in our analyses. We compute 50-year averages for ERF, substantially reducing the impact of natural year-to-year variability on the estimate
 275 (Fiedler et al., 2019a).

Table 1: Global means from the new MACv2-SP data and EC-Earth3 experiments. Shown are the ERF at the top of the atmosphere (ERF_{TOA}) and at the surface (ERF_{SFC}) calculated against pi as the mean \pm 95% confidence interval. (*) marks differences to the baseline that are statistically significant at the 95% level.

Experiment	$\overline{\tau_a}$	$\overline{\eta_N}$	ERF_{TOA} [Wm^{-2}]	ERF_{SFC} [Wm^{-2}]
2020-base	0.021	1.060	-0.661 ± 0.087	-1.449 ± 0.047
2020-blp	0.019	1.056	-0.622 ± 0.072	-1.338 ± 0.038 (*)
2050-base	0.019	1.054	-0.631 ± 0.068	-1.324 ± 0.035
2050-ff	0.020	1.057	-0.675 ± 0.081	-1.407 ± 0.041 (*)
2050-mg	0.012	1.038	-0.382 ± 0.080 (*)	-0.871 ± 0.036 (*)
2050-sg	0.014	1.040	-0.461 ± 0.069 (*)	-0.987 ± 0.039 (*)

3. Results

3.1. Annual means of τ_a and η_N

We show the global annual means of the anthropogenic aerosol optical depth ($\overline{\tau_a}$) and the prefactor for inducing aerosol effects on clouds ($\overline{\eta_N}$) from MACv2-SP in Figure 5. Note that we here show results from the new MACv2-SP data derived and provided in the framework of this study, but we only use the MACv2-SP data for 2020 and 2050 in time-slice simulations with EC-Earth3 later. We find that both $\overline{\tau_a}$ and $\overline{\eta_N}$ clearly reduce during the pandemic, e.g., by 0.002 for 2020 compared to the baseline and by 0.005 compared to 2005. These translate to a reduction by about 10% and 25% for the global $\overline{\tau_a}$ in 2020 and 2050. The associated effect on $\overline{\eta_N}$ is consistent with the change in $\overline{\tau_a}$, with a reduction of $\overline{\eta_N}$ by 0.004 compared to baseline and 0.017 compared to 2005, i.e., a global reduction in η_N by about 0.5–1%. All data sets have the same emissions for 2015–2023, except the baseline. Hence results from MACv2-SP other than the baseline are identical with the two-year-blip results for this period (Figure 5 and Table 1).

The post-pandemic recovery of τ_a and η_N strongly depends on the scenario. The $\overline{\tau_a}$ in 2030 is close to the value in 2005 in the baseline and two-year-blip




Figure 5: MACv2-SP τ_a in 2015–2050. Shown are the global annual means in (a) the anthropogenic aerosol optical depth (τ_a) at 550 nm and (b) the scaling factor the cloud droplet number concentration (η_N) for the scenarios. The pandemic period 2019–2023 is marked with grey shading. The baseline and two-year blip scenario overlap for 2030–2050. All scenarios except baseline overlap for 2015–2030. The 2005 value from the historical scaling is marked as horizontal line.

scenario, larger in the fossil fuel scenario, and substantially smaller for both
 295 green scenarios (Figure 5). By 2050 all scenarios point to a decrease of $\bar{\tau}_a$ relative
 to 2005, with the strongest reduction in the moderate green not the strong green
 scenario. This might be counter intuitive since a stronger green scenario might
 suggest cleaner air. The smaller $\bar{\tau}_a$ in the moderate green scenario is due to the
 lower emissions of NH_3 compared to the strong green by about -3.5×10^{-6} kg
 300 $\text{m}^{-2} \text{s}^{-1}$ in 2050, integrated over the globe and sectors. Although the emissions
 of SO_2 are smaller in the strong green scenario, the total effect of NH_3 and SO_2
 on the scaling factor E_i (Section 2) leads to a slightly larger $\bar{\tau}_a$ in the strong
 green scenarios.

The assumption of larger NH_3 emissions in the strong green recovery scenario
 305 compared to the moderate green recovery scenario is the result of changes in
 the structure of the economy in low emissions scenarios, particularly related to
 agricultural practice and energy provision. Stringent emissions scenarios to an
 increasing degree rely on more efficient food and biomass production to support
 a growing world population while generating low-carbon energy and enable pos-
 310 sibilities of carbon-dioxide sequestration (Popp et al., 2017). The fertilizer use

required to achieve this results in an increase in NH_3 emissions. Furthermore, stringent climate change mitigation scenarios often rely on very high shares of renewable energy, which have intermittent power generation properties and thus require energy storage technologies to bridge gaps in supply. One technology
 315 that can store energy across seasons and even multiple years is the production of ammonia (Society, 2020). The projected increased use of ammonia, as a fuel and for energy storage, results in larger projected emissions of NH_3 due to leakage and due to imperfect transport or storage. Slightly higher NH_3 emissions in the strong green (*sg*) than the moderate (*mg*) recovery scenarios are hence
 320 consistent with the general understanding of the technologies and practices that would be required for a transformation to a strongly decarbonized society.

3.2. Hemispheric asymmetry in τ_a

The spatial distribution of τ_a , measured by the hemispheric asymmetry, is qualitatively similar across the scenarios, but the magnitudes differ. Figure 6
 325 illustrates the hemispheric asymmetry A :

$$A = \frac{\overline{\tau_a(\phi)} - \overline{\tau_a(-\phi)}}{2} \quad (4)$$

using the zonal averages $\overline{\tau_a(\phi)}$ at the same geographical latitudes on the northern (ϕ) and southern hemisphere ($-\phi$). All scenarios have larger A in the tropics and sub-tropics than further poleward (Fig. 6), consistent with the CMIP6 scenarios (Fiebler et al., 2019a). For 2030, the fossil fuel and baseline scenarios
 330 have substantially higher A than for 2020. In the middle of the 21st century, A in the fossil fuel and baseline scenarios are more similar to each other and close to A from 2020, consistent with little differences in the spatial distribution of the emissions.

The temporal behaviour for A in the green scenarios is different to the fossil-fuel dominated scenarios, i.e., they are close to A from 2020 in 2030, but both
 335 much lower in 2050. Both green scenarios in 2030 have particularly similar A to 2020 in the tropics and slightly larger differences poleward. In 2050, the green scenarios are still similar to each other, but have overall smaller A compared to




Figure 6: Hemispheric asymmetry for MACv2-SP τ_a . Shown is the hemispheric asymmetry (\bar{A}) as function of the sinus of the geographical latitude for (a) 2030 and (b) 2050 for all scenarios. Baseline and two-year blip are identical for these years. We mark the values of the two-year blip scenario in 2020 as a reference.

2020, e.g., a reduction by 50% in the maximum around $\phi = 24^\circ$. This reflects
 340 the decrease in τ_a due to improved air quality in a green recovery (compare Fig.
 5).

3.3. Seasonal cycle in τ_a

The month-to-month changes in τ_a from MACv2-SP is dominated by the
 biomass burning season. These lead to tropical maxima in τ_a between July
 345 and October and November and February (Fig. 7a–b). Overall, the seasonal
 and zonal patterns in the two-year-blip scenario and baseline are very similar
 (not shown). Note here again that all data sets are identical with the two-year-
 blip in 2020, except the baseline.

In 2050, the overall seasonal pattern in τ_a remains qualitatively similar, but
 350 the magnitudes strongly depend on the scenario. The fossil fuel and baseline
 scenario are very similar (not shown), but the green scenarios show large reduc-
 tions in τ_a relative to baseline (Figure 8a–c). These are primarily projected in
 the northern hemisphere equatorward of 50°N . Here, the strongest reductions
 occur between June and October in the moderate green scenario. Again the
 355 stronger reduction in τ_a in the moderate than the strong green scenario is asso-




Figure 7: MACv2-SP τ_a patterns in 2020. Shown are the annual cycles of the anthropogenic aerosol optical depth (τ_a) at 550 nm as (a) zonal means and (b) zonal means weighted by $\bar{\tau}_a$ for the two-year-blip scenario.




Figure 8: MACv2-SP τ_a patterns in 2050. Shown is the annual cycle of the anthropogenic aerosol optical depth (τ_a) at 550 nm as (a) zonal mean for the baseline and (b–c) difference of the green scenarios relative to baseline for 2050.

ciated with the larger emissions of NH_3 in the strong green recovery leading to larger τ_a than for the moderate green scenario.

3.4. Global radiative forcing

We calculate the global ERF of anthropogenic aerosols from our experiments with the new MACv2-SP data at the top of the atmosphere and at the surface, ERF_{TOA} and ERF_{SFC} . For 2020, the impact of the reduction in anthropogenic aerosols compared to the baseline is a less negative ERF_{TOA} by about $+0.04 \text{ Wm}^{-2}$ (Table 1). This reduction is small compared to the year-to-year variability in the model, reflected by the confidence intervals about the mean of $\pm 0.07 \text{ Wm}^{-2}$ to $\pm 0.09 \text{ Wm}^{-2}$ across our ensemble of model simulations. Again these estimates are based on fifty years of simulations with annually repeating aerosol patterns. It will therefore be difficult to disentangle any differences in the TOA radiation budget due to reductions in aerosols during the pandemic from differences arising due to natural variability in both observations and small ensembles of simulations. We identify a larger and statistically significant difference in ERF_{SFC} associated with the aerosol reduction during the pandemic compared to the baseline of the order of 0.1 Wm^{-2} (Table 1). This implies that radiation observations at the surface and sufficiently many model estimates for ERF_{SFC} can be more informative for quantifying the influence of the pandemic on the global radiation and energy budget than estimates for the TOA.




Figure 9: Probability density function for the global mean ERF in 2050. Shown are the occurrence frequency of annual mean ERF at (a) the top of the atmosphere (TOA) and (b) the surface (SFC) for the color-coded scenarios in 2050. ERF is calculated from 50 years of data from our EC-Earth3 experiments with anthropogenic aerosols (*2050-base*, *2050-ff*, *2050-mg*, *2050-sg*) against the pre-industrial control experiment (*pi*).

The spread in ERF_{TOA} of anthropogenic aerosols due to the different scenarios for 2050 is -0.68 Wm^{-2} to -0.38 Wm^{-2} (Table 1). Compared to the baseline, these are differences of $+0.05$ to -0.25 Wm^{-2} . The least negative ERF_{TOA} occurs for the moderate green scenario, consistent with the lowest τ_a across the
 380 MACv2-SP data associated with the lower NH_3 emissions than in the strong green scenario. We obtain ERF_{TOA} of the anthropogenic aerosols for the green scenarios that are statistically significant different compared to the baseline. Baseline and the fossil-fuel based scenarios, however, yield very similar ERF_{TOA} for 2050, consistent with small differences in $\bar{\tau}_a$ and $\bar{\eta}_N$ for the two scenarios
 385 (Figure 9a). The ERF_{SFC} for 2050 is more negative than ERF_{TOA} (Figure 9b) and has a smaller 95% confidence interval of about $\pm 0.04 \text{ Wm}^{-2}$ compared to ERF_{TOA} . We therefore find for all scenarios a statistically significant difference in ERF_{SFC} relative to 2050-base (Table 1).




Figure 10: Differences in effective radiative effects due to the pandemic in 2020. Shown are differences between *2020-blp* and *2020-base* for the effective radiative effects in (left) all-sky (F_{all}), (middle) clear-sky (F_{clr}) and (right) cloudy-sky (F_{cld}) at (top) the top of the atmosphere and (bottom) the surface. Hatched areas mark regions where the differences are not statistically significant at the 95% confidence level.

3.5. Pattern of radiative effects

390 We assess the spatial distribution of the radiative effects associated with the anthropogenic aerosols. To this end, we calculate the effective radiative effects in all-sky (F_{all}), clear-sky (F_{clr}), and cloudy-sky (F_{cld}) using the relationship:

$$F_{all} = (1 - f)F_{clr} + fF_{cld}, \quad (5)$$

with the total cloud cover (f). Figure 10 and 11 show the results for the top of the atmosphere (TOA) and at the surface.

395 For 2020, we find some evidence for regionally significant differences in the radiative effects at TOA associated with anthropogenic aerosol reductions due to the pandemic. We measure significance at the 95% confidence level. Significant regional signals at TOA are rare, and primarily limited to F_{clr} (Figure 10). This is primarily explained by the strong variability of clouds. The difference in F_{clr} at TOA indicates less negative radiative effects over some regions, e.g., offshore of typically polluted regions in Asia. This is consistent with the
400 expectation of a more negative aerosol radiative effect for a given aerosol plume

over a dark ocean surface than over a relatively brighter land surface. Less aerosols during 2020 therefore coincide with a less negative radiative effect of anthropogenic aerosols over the ocean than over land, seen as positive anomaly in Figure 10. At the surface, the regional differences in the described radiative effects are more pronounced and spatially further extended in F_{clr} , covering large parts in Southeast Asia and East Asia both over land and ocean, and are also seen in F_{all} for some regions (Figure 10). Surface measurements in these regions could potentially help to constrain the aerosol effects on climate. Much of the radiative effects occur over oceans, where the measurement network is typically sparse. Efforts to collect necessary observations during this unique situation could involve sun photometer measurements aboard research vessels as part of the Maritime Aerosol Network (e.g., Spirnov et al., 2009) and in-situ measurements aboard aircrafts based on existing expertises (e.g., Zuidema et al., 2016).

The scenario differences in the radiative effects at TOA for 2050 are largest in South, Southeast and East Asia (Figure 11). Here, F_{clr} shows significant differences with less negative radiative effects in the moderate green scenario compared to the baseline by up to 3 Wm^{-2} . This result is in agreement with the expectation of less aerosol scattering and absorption associated with the lower τ_a . The pattern of F_{clr} is again more inhomogeneous than F_{clr} . Scenario differences for F_{clr} are again not significant for most world regions. The overall pattern for F_{all} differences at the surface is qualitatively similar to TOA, but the magnitude and spatial extent of significant differences are larger at the surface. Based on these results, significant scenario differences for the radiative effects associated with anthropogenic aerosols are primarily confined to the tropics and sub-tropics close to hotspots for industrial and biomass burning activity at present.




Figure 11: As Fig. 10 but for the differences between the moderate green (*2050-mg*) against the baseline (*2050-base*) scenario in 2050.

430 4. Discussion and conclusion

We show the anthropogenic aerosol optical properties and the associated effect on clouds from MACv2-SP. The new MACv2-SP data is based on the COVID-19 emission data from Forster et al. (2020a) for use in CovidMIP (Jones et al., 2021; Lamboll et al., 2021). Our results point to a reduction in the global anthropogenic aerosol optical depth by 10% due to the pandemic compared to the baseline. Scenarios for the post-pandemic recovery indicate a continuous increase in aerosols until 2030 in scenarios without more green technologies and a general decrease in all scenarios for 2030–2050. The spread in the anthropogenic aerosol optical depth in 2050 is 0.012 (medium green scenario) to 0.02 (fossil fuel scenario). These values fall within the lower end of the spread in 2050 obtained from the original CMIP6 aerosol scenarios (Fiedler et al., 2019b).

First estimates of the effective radiative forcing (ERF) associated with the new MACv2-SP data are calculated from several hundred years of atmosphere-only simulations with EC-Earth3. The results highlight a weaker (less negative) aerosol ERF of the order of 10% during the pandemic relative to the baseline. Such small ERF differences require long averaging, hence our 50 years of simulations for each aerosol pattern assessed here. The small change in aerosol ERF

for the time of the pandemic is not expected to induce a global climate response that is clearly detectable in light of model-internal variability. Even regional
450 radiative effects are rather difficult to detect at the top of the atmosphere. We find, however, more significant effective radiative effects at the surface in regions typically more strongly polluted by aerosols. Any attempt to use the pandemic period to constrain aerosol effects should therefore focus on areas in South and East Asia, primarily assessing radiative effects at the surface paired
455 with a validation of the representation of observed regional emissions. This may involve station observations for radiation and aerosols, e.g., from aerosol networks (Xin et al., 2015; Giles et al., 2019). Our experiments suggest that aerosol radiative effects are expected offshore of land with major pollution in the past decades. We therefore propose to also use other measurements, e.g., from
460 sun photometers aboard research vessels or in-situ instruments from aircrafts. Measurements outside of clouds might be particularly beneficial, although much research focuses on aerosol-cloud interactions. Our model results indeed suggest that there are better chances to obtain a signal in clear-sky conditions at the surface rather than in cloudy and all sky. We mostly find poor prospects to
465 measure a meaningful regional effect on clouds due to the strong influence of natural variability.

For 2050, we obtain an ERF spread of -0.68 to -0.38 Wm^{-2} relative to the pre-industrial, which is less negative than the ERF from the same model for 2005 and 1975 (Fiedler et al., 2019a). These ERF estimates for 2050 fall within the
470 ERF spread for 2095 associated with the emission pathways from CMIP6 and uncertainty in aerosol-cloud interactions (Fiedler et al., 2019b). Interestingly, the stronger green scenario does not yield the smallest anthropogenic aerosol optical depth and least negative forcing, but the moderate green recovery does. This is associated with a relative increase in NH_3 emissions due to intense land-
475 use paired with an energy system primarily relying on renewable sources. Such a pathway implies a slightly stronger warming due to weaker aerosol cooling in the strong green than the moderate green scenario. We expect, however, a stronger reduction in greenhouse gas emissions in the strong green pathway.

Taken together the anthropogenic warming in the strong green scenario is there-
480 fore expected to be the weakest. Regionally, our simulations suggest the largest
differences in the aerosol radiative effects across sub-tropical and tropical re-
gions. An interesting, yet difficult question in this context is to what extent
future changes in emissions, and the associated radiative forcing and climate
response, are truly attributed to influences of the pandemic and how much of it
485 would have occurred for other reasons, e.g., due to policy changes as measure
for improving air quality and mitigating climate change.

Our global ERF estimates for the anthropogenic aerosols fall within the plausi-
ble range of the present-day aerosol ERF (Belloin et al., 2020) underlining
the still large uncertainty in our understanding of aerosol effects compared to
490 our ability to estimate a change in ERF from different emission pathways from
a complex model. We expect that models participating in CovidMIP will show
diversity in their aerosol ERF owing to model-internal variability and model
biases, even when they use the same emissions or MACv2-SP data (e.g., Fiedler
et al., 2019a; Smith et al., 2020). For instance, Weber et al. (2020) point to the
495 small signal compared to the noise in a complex aerosol-climate model simula-
tion for spring 2020. Reasons for the model diversity in aerosol ERF include not
only uncertainties in the aerosol parameterizations, but also the ability of the
host model to accurately simulate important processes influencing the aerosol
life cycle and therefore ERF, e.g., the representation of clouds, precipitation,
500 and circulation. Future research should therefore also address the relative con-
tributions from host model biases to the model diversity in ERF.

5. Acknowledgments

SF acknowledges the funding from the German Research Foundation for the
Collaborative Research Centre 1211 (ID: DFG 268236062, [https://sfb1211.
505 uni-koeln.de](https://sfb1211.uni-koeln.de)) and from the German Federal Ministry for Transportation and
Digital Infrastructure for the Hans-Ertel-Centre for Weather Research (ID:
BMVI/DWD 4818DWDP5A, <https://www.herz.uni-bonn.de>) as well as re-

sources of the Deutsches Klimarechenzentrum (DKRZ) granted by its Scientific Steering Committee (WLA) under project ID bb1198. KW and TvN received support from European Union's Horizon 2020 research and innovation programme under grant agreement No 821205 (FORCeS). The EC-Earth3 simulations were performed on resources provided by the Swedish National Infrastructure for Computing (SNIC) at NSC. We thank the four anonymous reviewers for their comments and the editor for handling our manuscript.

6. Data availability

The MACv2-SP code and input data are freely accessible in the supplement of previous (Stevens et al., 2017; Fiedler et al., 2019a) and this article. The results from the EC-Earth3 simulations used in this study are available from Wyser (2020).

Processed MODIS satellite data for the aerosol optical depth was downloaded from the Giovanni online data system (MOD08_M3 v6.1, <https://giovanni.gsfc.nasa.gov/>, accessed 4 Jan 2021), developed and maintained by the NASA GES DISC. We also acknowledge the MODIS mission scientists and associated NASA personnel for the production of the data shown in this article.

References

References

- Abdul-Razzak, H., Ghan, S.J., 2000. A parameterization of aerosol activation: 2. Multiple aerosol types. *Journal of Geophysical Research: Atmospheres* 105, 6837–6844. doi:<https://doi.org/10.1029/1999JD901161>.
- Acker, J.G., Leptoukh, G., 2007. Online Analysis Enhances Use of NASA Earth Science Data. URL: <https://giovanni.gsfc.nasa.gov>.
- Andrijevic, M., Schleussner, C.F., Gidden, M.J., McCollum, D.L., Rogelj, J., 2020. Covid-19 recovery funds dwarf clean energy investment needs. *Science* 370, 298–300. doi:[10.1126/science.abc9697](https://doi.org/10.1126/science.abc9697).

- 535 Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K.S., Christensen, M., Daniau, A.L., Dufresne, J.L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J.M., Lohmann, U., Malavelle, F., Mauritsen, T., McCoy, D.T., Myhre, G., Mülmenstädt, J., Neubauer, D., Possner, A., Ruge-
540 genstein, M., Sato, Y., Schulz, M., Schwartz, S.E., Sourdeval, O., Storelvmo, T., Toll, V., Winker, D., Stevens, B., 2020. Bounding global aerosol radiative forcing of climate change. *Reviews of Geophysics* 58, e2019RG000660. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019RG000660>, doi:10.1029/2019RG000660.
- 545 Berman, J.D., Ebisu, K., 2020. Changes in u.s. air pollution during the covid-19 pandemic. *Science of The Total Environment* 739, 139864. doi:<https://doi.org/10.1016/j.scitotenv.2020.11.9864>.
- Broomandi, P., Karaca, F., Nik'al, A., Jahanbakhshi, A., Tamjidi, M., Kim, J.R., 2020. Impact of COVID-19 event on the air quality in iran. *Aerosol and*
550 *Air Quality Research* 20, 1793–1804. doi:10.4209/aaqr.2020.05.0205.
- Cherian, R., Quaas, J., 2020. Trends in AOD, Clouds, and Cloud Radiative Effects in Satellite Data and CMIP5 and CMIP6 Model Simulations Over Aerosol Source Regions. *Geophysical Research Letters* 47, e2020GL087132. doi:<https://doi.org/10.1029/2020GL087132>.
- 555 Chu, B., Ma, Q., Liu, J., Ma, J., Zhang, P., Chen, T., Feng, Q., Wang, C., Yang, N., Ma, H., Ma, J., Russell, A.G., He, H., 2020. Air pollutant correlations in china: Secondary air pollutant responses to nox and so2 control. *Environmental Science & Technology Letters* 7, 695–700. doi:10.1021/acs.estlett.0c00403.
- 560 Döscher, R., Acosta, M., Alessandri, A., Anthoni, P., Arneth, A., Arsouze, T., Bergmann, T., Bernadello, R., Bousetta, S., Caron, L.P., Carver, G., Castrillo, M., Catalano, F., Cvijanovic, I., Davini, P., Dekker, E., Doblas-Reyes, F.J., Docquier, D., Echevarria, P., Fladrich, U., Fuentes-Franco, R.,

- Gröger, M., v. Hardenberg, J., Hieronymus, J., Karami, M.P., Keskinen,
 565 J.P., Koenigk, T., Makkonen, R., Massonnet, F., Ménégoz, M., Miller, P.A.,
 Moreno-Chamarro, E., Nieradzik, L., van Noije, T., Nolan, P., O'Donnell, D.,
 Ollinaho, P., van den Oord, G., Ortega, P., Prims, O.T., Ramos, A., Reerink,
 T., Rousset, C., Ruprich-Robert, Y., Le Sager, P., Schmith, T., Schrödner, R.,
 Serva, F., Sicardi, V., Sloth Madsen, M., Smith, B., Tian, T., Tourigny, E.,
 570 Uotila, P., Vancoppenolle, M., Wang, S., Wårlind, D., Witten, U., Wyser, K.,
 Yang, S., Yepes-Arbós, X., Zhang, Q., 2021. The EC-Earth3 Earth System
 Model for the Climate Model Intercomparison Project 3. Geoscientific Model
 Development Discussions 2021, 1–90. doi:10.5194/gmd-2020-446.
- ECMWF, 2010a. IFS DOCUMENTATION – Cy36r1 Operational imple-
 575 mentation 26 January 2010, PART III: DYNAMICS AND NUMERI-
 CAL PROCEDURES. European Centre for Medium Range Weather
 Forecasts. URL: [www.ecmwf.int/sites/default/files/elibrary/2010/
 9232-part-iii-dynamics-and-numerical-procedures.pdf](http://www.ecmwf.int/sites/default/files/elibrary/2010/9232-part-iii-dynamics-and-numerical-procedures.pdf).
- ECMWF, 2010b. IFS DOCUMENTATION – Cy36r1 Operational
 580 implementation 26 January 2010, PART IV: PHYSICAL PRO-
 CESSSES. European Center for Medium Range Weather Fore-
 casts. URL: [www.ecmwf.int/sites/default/files/elibrary/2010/
 9233-part-iv-physical-processes.pdf](http://www.ecmwf.int/sites/default/files/elibrary/2010/9233-part-iv-physical-processes.pdf).
- Eyring, V., Boyle, S., Meehl, G.A., Senior, C.A., Stevens, B., Stouffer, R.J.,
 585 Taylor, K.F., 2016. Overview of the Coupled Model Intercomparison Project
 Phase 6 (CMIP6) experimental design and organization. Geosci. Model Dev.
 9, 1937–1958. doi:10.5194/gmd-9-1937-2016.
- Fiedler, S., Kinne, S., Huang, W.T.K., Räisänen, P., O'Donnell, D., Bellouin,
 N., Stier, P., Merikanto, J., van Noije, T., Carslaw, K., Makkonen, R.,
 590 Lohmann, U., 2019a. Anthropogenic aerosol forcing – insights from multi-
 estimates from aerosol-climate models with reduced complexity. Atmospheric
 Chemistry and Physics 19, 6821–6841. doi:10.5194/acp-19-6821-2019.

- Fiedler, S., Stevens, B., Gidden, M., Smith, S.J., Riahi, K., van Vuuren, D.,
 2019b. First forcing estimates from the future CMIP6 scenarios of anthro-
 595 pogenic aerosol optical properties and an associated Twomey effect. *Geosci-
 entific Model Development* 12, 989–1007. doi:10.5194/gmd-12-989-2019.
- Fiedler, S., Stevens, B., Mauritsen, T., 2017. On the sensitivity of anthropogenic
 aerosol forcing to model-internal variability and parameterizing a Twomey
 effect. *J. Adv. Model. Earth Syst.* 9, n/a–n/a. doi:10.1002/2017MS000932.
- 600 Forster, P.M., Forster, H.I., Evans, M.J., Gidden, M.J., Jones, C.D.,
 Keller, C.A., Lamboll, R.D., Quéré, C.L., Rogelj, P., Rosen, D., Schleuss-
 ner, C.F., Richardson, T.B., Smith, C.J., Tameck, S.T., 2020a. Cur-
 rent and future global climate impacts resulting from covid-19. *Nature
 Climate Change* URL: <https://doi.org/10.1038/s41558-020-0883-0>,
 605 doi:10.1038/s41558-020-0883-0.
- Forster, P.M., Lamboll, R., Rogelj, P., 2020b. Emissions changes in 2020 due to
 Covid19 doi:10.5281/zenodo.3957826.
- Gettelman, A., Lamboll, R., Bardeen, C.G., Forster, P.M., Watson-Parris,
 D., . . . Climate Impacts of COVID-19 Induced Emission Changes. *Geo-
 610 physical Research Letters* , e2020GL091805doi:[https://doi.org/10.1029/
 2020GL091805](https://doi.org/10.1029/2020GL091805). e2020GL091805 2020GL091805.
- Giles, D.M., Slevin, A., Sorokin, M.G., Schafer, J.S., Smirnov, A., Slutsker, I.,
 Eck, T.F., Polben, B.N., Lewis, J.R., Campbell, J.R., Welton, E.J., Korkin,
 S.V., Lyapustin, A.I., 2019. Advancements in the Aerosol Robotic Network
 615 (AERONET) Version 3 database – automated near-real-time quality control
 algorithm with improved cloud screening for Sun photometer aerosol optical
 depth (AOD) measurements. *Atmospheric Measurement Techniques* 12, 169–
 209. doi:10.5194/amt-12-169-2019.
- Gillett, N.P., Shiogama, H., Funke, B., Hegerl, G., Knutti, R., Matthes, K., San-
 620 ter, B.D., Stone, D., Tebaldi, C., 2016. The Detection and Attribution Model

Intercomparison Project (DAMIP v1.0) contribution to CMIP6. *Geoscientific Model Development* 9, 3685–3697. doi:10.5194/gmd-9-3685-2016.

van Heerwaarden, C.C., Mol, W.B., Veerman, M.A., Benedict, I., Heusinkveld, B.G., Knap, W.H., Kazadzis, S., Kouremeti, N., Fiedler, S., 2021. Record
 625 high solar irradiance in western europe during first covid-19 lockdown largely due to unusual weather. *Communications Earth & Environment* 2, 37. URL: <https://doi.org/10.1038/s43247-021-00110-0>, doi:10.1038/s43247-021-00110-0.

Huppmann, D., Rogelj, J., Kriegler, E., Krey, V., Nabel, K., 2018. A new
 630 scenario resource for integrated 1.5 °C research. *Nature Climate Change* 8, 1027–1030. doi:10.1038/s41558-018-0317-4.

IPCC, 2021. Annex II: Models [Gutiérrez, J. M., A-M. Treguiér (eds.)]. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on
 635 Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Cui, J. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press. In Press .

Jones, C.D., Hickman, J.E., Rumbold, S.T., Walton, J., Lamboll, R.D., Skeie,
 640 R.B., Fiedler, S., Forster, P.M., Rogelj, J., Abe, M., Botzet, M., Calvin, K., Cassou, C., Cole, J.N., Davini, P., Deushi, M., Dix, M., Fyfe, J.C., Gillett, N.P., Ilyina, T., Kawamiya, M., Kelley, M., Kharin, S., Koshiro, T., Li, H., Mackallah, C., Müller, W.A., Nabat, P., van Noije, T., Nolan, P., Ohgaito, R., Olivie, D., Oshima, N., Parodi, J., Reerink, T.J., Ren, L., Romanou, A., Séférian, R., Tang, Y., Timmreck, C., Tjiputra, J., Tourigny, E., Tsigaridis,
 645 K., Wang, H., Wu, M., Wyser, K., Yang, S., Yang, Y., Ziehn, T., 2021. The Climate Response to Emissions Reductions Due to COVID-19: Initial Results From CovidMIP. *Geophysical Research Letters* 48, e2020GL091883. doi:<https://doi.org/10.1029/2020GL091883>.

- 650 Lamboll, R.D., Jones, C.D., Skeie, R.B., Fiedler, S., Samset, B.H., Gillett, N.P., Rogelj, J., Forster, P.M., 2021. Modifying emissions scenario projections to account for the effects of covid-19: protocol for covidmip. *Geoscientific Model Development* 14, 3683–3695. URL: <https://gmd.copernicus.org/articles/14/3683/2021/>, doi:10.5194/gmd-14-3683-2021.
- 655 Lamboll, R.D., Nicholls, Z.R.J., Kikstra, J.S., Meinshausen, M., Rogelj, J., 2020b. Silicone v1.0.0: an open-source python package for inferring missing emissions data for climate change research, 1–23doi:<https://doi.org/10.5194/gmd-2020-138>.
- 660 Madec, G., the NEMO team, 2016. NEMO ocean engine. Note du Pôle de modélisation, Institut Pierre-Simon Laplace (IPSL), France, No 27, ISSN No 1288-1619.
- Mauritsen, T., Bader, J., Becker, G., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Colinger, T., Esch, M., Fast, I., Fiedler, S., Fläschner, D., Gayler, V., Giorgetta, M., Goll, D.S., Haak, H., Hagemann, S., Hedemann, C., Hoheneser, C., Ilyina, T., Jahns, T., Jimenez de la Cuesta Otero, D., Jurgens, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lisslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B., Müller, W.A., Nabel, J.E.M.S., Nam, C.C.W., Notz, D., Nyawira, S.S., Paulsen, H., Peters, K., Pincus, R., Polmann, H., Pongratz, J., Popp, M., Raddatz, T., Rast, S., Redler, R., Reick, C.H., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K.D., Stein, L., Stemmler, I., Stevens, B., von Storch, J.S., Tian, F., Voigt, A., de Vrese, P., Wieners, K.H., Wilken-
675 skjeld, S., Winkler, A., Roeckner, E., 2019. Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM 1.2) and its response to increasing CO₂. *Journal of Advances in Modeling Earth Systems* 11, 998–1038. doi:10.1029/2018MS001400.
- van Noije, T., Bergman, T., Le Sager, P., O'Donnell, D., Makkonen, R.,

- 680 Gonçalves-Ageitos, M., Döscher, R., Fladrich, U., von Hardenberg, J., Keskinen, J.P., Korhonen, H., Laakso, A., Myriokefalitakis, S., Ollinaho, P., Pérez García-Pando, C., Reerink, T., Schrödner, R., Wyser, K., Yang, S., 2021. EC-Earth3-AerChem, a global climate model with interactive aerosols and atmospheric chemistry participating in CMIP6. *Geoscientific Model Development Discussions* accepted, 1–46. URL: <https://doi.org/10.5194/gmd-2020-413>.
- 685 Pincus, R., Forster, P.M., Stevens, B., 2016. The Radiative Forcing Model Intercomparison Project (RFMIP): experimental protocol for CMIP6. *Geoscientific Model Development* 9, 3447–3460. doi:10.5194/gmd-9-3447-2016.
- Platnick, S., Hubanks, P., Meyer, K., Knutson, M.D., 2015. MODIS Atmosphere L3 Monthly Product (v8 L3). NASA MODIS Adaptive Processing System, Goddard Space Flight Center. doi:[http://dx.doi.org/10.5067/MODIS/MOD08_M3.006\(Terra\)](http://dx.doi.org/10.5067/MODIS/MOD08_M3.006(Terra)), [http://dx.doi.org/10.5067/MODIS/MYD08_M3.006\(Aqua\)](http://dx.doi.org/10.5067/MODIS/MYD08_M3.006(Aqua)).
- 695 Popp, A., Calvin, K., Fujimori, S., Havlik, P., Humpenöder, F., Stehfest, E., Bodirsky, B.L., Dietrich, J.P., Doelmann, J.C., Gusti, M., Hasegawa, T., Kyle, P., Obersteiner, M., Tabebu, A., Takahashi, K., Valin, H., Waldhoff, S., Weindl, I., Wise, M., Kriegler, E., Lotze-Campen, H., Fricko, O., Riahi, K., Vuuren, D.P.v., 2017. Land-use futures in the shared socio-economic pathways. *Global Environmental Change* 42, 331–345. doi:10.1016/j.gloenvcha.2016.10.002.
- 700 Ranjan, A.K., Patra, A.K., Gorai, A.K., 2020. Effect of lockdown due to sars covid-19 on aerosol optical depth (aod) over urban and mining regions in india. *The Science of the total environment* 745, 141024–141024. doi:10.1016/j.scitotenv.2020.141024.
- Riemer, N., Doherty, O.M., Hameed, S., . on the variability of african dust transport across the atlantic .

- Rogelj, J., Shindell, D., Jiang, K., Fifita, S., Forster, P., Ginzburg, V., Handa, C., Kheshgi, H., Kobayashi, S., Kriegler, E., Mundaca, L., Sefarian, R., Vilarino, M.V., 2018. Mitigation pathways compatible with 1.5°C in the context of sustainable development. IPCC/WMO. p. 93–174. URL: <http://www.ipcc.ch/report/sr15/>.
710
- Sanap, S., 2021. Global and regional variations in aerosol loading during COVID-19 imposed lockdown. *Atmospheric Environment* 246, 118132. doi:<https://doi.org/10.1016/j.atmosenv.2020.118132>.
- 715 Smirnov, A., Holben, B.N., Slutsker, I., Giles, D.M., McClain, C.R., Eck, T.F., Sakerin, S.M., Macke, A., Croot, P., Zibordi, G., Quinn, P.K., Sciare, J., Kinne, S., Harvey, M., Smyth, T.J., Piketh, S., Zielinski, T., Proshutinsky, A., Goes, J.I., Nelson, N.B., Larouche, P., Radionov, V.F., Goloub, P., Krishna Moorthy, K., Matarrese, R., Pobereson, E.J., Jourdin, F., 2009. Maritime aerosol network as a component of aerosol robotic network. *Journal of Geophysical Research: Atmospheres* 114. doi:10.1029/2008JD011257.
720
- Smith, C.J., Kramer, R.J., Muehre, G., Alterskjær, K., Collins, W., Sima, A., Boucher, O., Dufrene, J.L., Nabat, P., Michou, M., Yukimoto, S., Cole, J., Paynter, D., Shiogama, H., O'Connor, F.M., Robertson, E., Wiltshire, A., Andrews, T., Hannay, C., Miller, R., Nazarenko, L., Kirkevåg, A., Olivie, E., Friedler, S., Lewinschal, A., Mackallah, C., Dix, M., Pincus, R., Forster, P.M., 2020. Effective radiative forcing and adjustments in cmip6 models. *Atmospheric Chemistry and Physics* 20, 9591–9618. doi:10.5194/acp-20-9591-2020.
- 730 Society, T.R., 2020. Ammonia: zero-carbon fertiliser, fuel and energy store. 978-1-78252-448-9. URL: <https://royalsociety.org/-/media/policy/projects/green-ammonia/green-ammonia-policy-briefing.pdf>.
- Song, Q., Zhang, Z., Yu, H., Ginoux, P., Shen, J., 2021. Global dust optical depth climatology derived from caliop and modis aerosol retrievals on decadal

- 735 time scales: Regional and interannual variability. *Atmospheric Chemistry and Physics Discussions* 2021, 1–57. doi:10.5194/acp-2021-1.
- Srivastava, A.K., Bhojar, P.D., Kanawade, V.P., Devara, P.C., Thomas, A., Soni, V.K., 2021. Improved air quality during COVID-19 at an urban megacity over the Indo-Gangetic Basin: From stringent to relaxed lockdown phases. *Urban Climate* 36, 100791. URL: <https://www.sciencedirect.com/science/article/pii/S221203521000213>, doi:<https://doi.org/10.1016/j.uclim.2021.100791>.
- 740 Stevens, B., Fiedler, S., Kinne, S., Peters, K., Rast, S., Müsse, J., Smith, S.J., Mauritsen, T., 2017. MACv2-SP: a parameterization of anthropogenic aerosol optical properties and an associated Twomey effect for use in CMIP6. *Geosci. Mod. Dev.* 10, 433–452. doi:10.5194/gmd-10-433-2017.
- Sun, W., Shao, M., Granier, C., Liu, Y., Ye, C.S., Zheng, J.Y., 2018. Long-Term Trends of Anthropogenic SO₂, NO_x, CO, and NMVOCs Emissions in China. *Earth's Future* 6, 1112–1130. doi:<https://doi.org/10.1029/2018EF000822>.
- 750 Weber, J., Shin, Y.M., Staunton-Sykes, J., Archer-Nicholls, S., Abraham, N.L., Archibald, A.T., 2020. Minimal Climate Impacts From Short-Lived Climate Forcers Following Emission Reductions Related to the COVID-19 Pandemic. *Geophysical Research Letters* 47, e2020GL090326. doi:<https://doi.org/10.1029/2020GL090326>.
- 755 Wyser, K., 2021. Effective radiative forcing (ERF) experiments with EC-Earth3. doi:10.5281/zenodo.4106258.
- Wyser, K., Kjellström, E., Koenigk, T., Martins, H., Döscher, R., 2020. Warmer climate projections in EC-Earth3-Veg: the role of changes in the greenhouse gas concentrations from CMIP5 to CMIP6. *Environmental Research Letters* 760 15, 054020.
- Xin, J., Wang, Y., Pan, Y., Ji, D., Liu, Z., Wen, T., Wang, Y., Li, X., Sun, Y., Sun, J., Wang, P., Wang, G., Wang, X., Cong, Z., Song, T., Hu, B.,

- 765 Wang, L., Tang, G., Gao, W., Guo, Y., Miao, H., Tian, S., Wang, L., 2015. The Campaign on Atmospheric Aerosol Research Network of China: CARE-China. *Bulletin of the American Meteorological Society* 96, 1137 – 1155. doi:10.1175/BAMS-D-14-00039.1.
- 770 Yamazaki, D., de Almeida, G.A.M., Bates, P.D., 2013. Improving computational efficiency in global river models by implementing the local inertial flow equation and a vector-based river network map. *Water Resources Research* 49, 7221–7235. doi:<https://doi.org/10.1002/wrcr.12055>.
- Yang, Y., Ren, L., Li, H., Wang, H., Wang, P., Chen, L., Yue, X., Liao, H., 2020. Fast Climate Responses to Aerosol Emission Reductions During the COVID-19 Pandemic. *Geophysical Research Letters* 47, e2020GL089788. doi:<https://doi.org/10.1029/2020GL089788>.
- 775 Zhang, M., Katiyar, A., Zhu, S., Shen, J., Xia, M., Ma, J., Kota, S.H., Wang, P., Zhang, H., 2021. Impact of reduced anthropogenic emissions during COVID-19 on air quality in India. *Atmospheric Chemistry and Physics* 21, 4025–4037. URL: <https://acp.copernicus.org/articles/21/4025/2021/>, doi:10.5194/acp-21-4025-2021.
- 780 Zuidema, P., Redemann, J., Haywood, J., Wood, R., Piketh, S., Hipondoka, M., Formenti, P., 2016. Smoke and Clouds above the Southeast Atlantic: Upcoming Field Campaigns Probe Absorbing Aerosol's Impact on Climate. *Bulletin of the American Meteorological Society* 97, 1131–1135. doi:10.1175/BAMS-D-15-00082.1.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof

Stephanie Fiedler led the study and the writing of the manuscript. She created, analysed, and visualized the MACv2-SP data. Klaus Wyser performed the EC-Earth3 experiments, analysed the forcing in the model output, and created the associated figures. All authors contributed to the writing, reviewing and editing of the article.

Journal Pre-proof

Manuscript: “Radiative effects of reduced aerosol emissions during the COVID-19 pandemic and the future recovery”

Authors: Stephanie Fiedler, Klaus Wyser, Joeri Rogelj, and Twan van Noije

Highlights

- New COVID-19 data to parameterize anthropogenic aerosol properties are released for use in climate studies.
- First MACv2-SP estimate of anthropogenic aerosol radiative forcing for 2020 suggests a change by $+0.04 \text{ Wm}^{-2}$ due to the pandemic.
- Recovery scenarios for 2050 have a spread in anthropogenic aerosol forcing of -0.38 to -0.68 Wm^{-2} relative to pre-industrial.