

# 1 Increasing risks of multiple breadbasket failure under 1.5 and 2°C global 2 warming

3 Franziska Gaupp\*<sup>^</sup>, Jim Hall\*, Dann Mitchell\*\* , Simon Dadson\*

4 \* *Environmental Change Institute, University of Oxford*; <sup>^</sup>*International Institute for Applied Systems Analysis*;  
5 \*\* *University of Bristol*

## 6 *Abstract*

7 The increasingly inter-connected global food system is becoming more vulnerable to production  
8 shocks owing to increasing global mean temperatures and more frequent climate extremes. Little is  
9 known, however, about the actual risks of multiple breadbasket failure due to extreme weather  
10 events. Motivated by the Paris Climate Agreement, this paper quantifies spatial risks to global  
11 agriculture in 1.5 and 2°C warmer worlds. This paper focuses on climate risks posed to three major  
12 crops - wheat, soybean and maize - in five major global food producing areas. Climate data from the  
13 atmosphere-only HadAM3P model as part of the “Half a degree Additional warming, Prognosis and  
14 Projected Impacts” (HAPPI) experiment are used to analyse the risks of climatic extreme events. Using  
15 the copula methodology, the risks of simultaneous crop failure in multiple breadbaskets are  
16 investigated. Projected losses do not scale linearly with global warming increases between 1.5 and 2°C  
17 Global Mean Temperature (GMT). In general, whilst the differences in yield at 1.5 versus 2°C are  
18 significant they are not as large as the difference between 1.5°C and the historical baseline which  
19 corresponds to 0.85°C above pre-industrial GMT. Risks of simultaneous crop failure, however, do  
20 increase disproportionately between 1.5 and 2°C, so surpassing the 1.5°C threshold will represent a  
21 threat to global food security. For maize, risks of multiple breadbasket failures increase the most, from  
22 6% to 40% at 1.5 to 54% at 2°C warming. In relative terms, the highest simultaneous climate risk  
23 increase between the two warming scenarios was found for wheat (40%), followed by maize (35%)  
24 and soybean (23%). Looking at the impacts on agricultural production, we show that limiting global  
25 warming to 1.5°C would avoid production losses of up to 2 753 million (161 000, 265 000) tonnes  
26 maize (wheat, soybean) in the global breadbaskets and would reduce the risk of simultaneous crop  
27 failure by 26%, 28% and 19% respectively.

28  
29 Keywords: Climate Risks, Multiple Breadbasket Failure, Paris Agreement, Copula Methodology

## 30 31 1 Introduction

32 The Paris Agreement in 2015, in which 197 countries agreed to limit the increase of mean global  
33 temperature to 1.5°C rather than 2°C above pre-industrial levels (UNFCCC, 2015), has received  
34 considerable interest from the scientific community (i.e., Mitchell et al. 2016b; Rogelj and Knutti 2016;  
35 Verschuuren 2016; Schleussner et al. 2016; James et al. 2017). However, so far little research has been  
36 done on the impacts of a 1.5°C temperature increase, let alone on the quantification of the differential  
37 impacts of 1.5 versus 2°C global warming (James et al., 2017). Quantitative impacts assessments of

38 the relative benefits of limiting global warming to 1.5°C are required to support such policies and the  
39 scientific community is now encouraged to address research gaps related to a 1.5°C temperature  
40 increase, especially to the different impacts at local and regional scales (Rogelj and Knutti, 2016) and  
41 the impacts on other industries.

42 This paper focuses on the climate change impacts on the agricultural sector. Although agriculture is  
43 not explicitly mentioned in the Paris Climate Agreement, “safeguarding food security” and the  
44 “vulnerabilities of the food production systems to the adverse impacts of climate change” are  
45 recognized (UNFCCC, 2015). Agriculture is one of the sectors that will experience the largest negative  
46 impacts from climatic change (Porter et al., 2014). Climate trends and specifically climate variability  
47 have already negatively impacted agricultural production in many regions (Field and IPCC, 2012; Lobell  
48 et al., 2011). On the other hand, it has been estimated that by 2050, an increase of 40% of global food  
49 production is necessary to meet the growing demand resulting from population growth and rising  
50 calorie intake in developing countries (Verschuuren, 2016). Today, FAO (2014) estimates that 805  
51 million people are undernourished globally, which is one in nine people. In a crisis such as the 2007/08  
52 food price crisis, however, the number of undernourished people increased by 75 million in only four  
53 years owing to food price spikes for major crops (Von Braun, 2008). An increasingly interconnected  
54 global food system (Puma et al., 2015) and the projected fragility of the global food production system  
55 due to climatic change (Fraser et al., 2013) further exacerbate the threats to food security. The  
56 potential impact of simultaneous climate extremes on global food security is in particular need of  
57 further investigation. Crop losses in a single, main crop producing area, termed a breadbasket, can be  
58 offset through trade with other crop-producing regions (Bren d’Amour et al., 2016). If several  
59 breadbaskets suffer from negative climate impacts at the same time, however, global production  
60 losses might lead to price shocks and trigger export restrictions which amplify the threats to global  
61 food security (Puma et al., 2015).

62

63 Research has started to focus on the impacts of multiple, interconnected adverse weather events on  
64 agricultural production and indirect effects on other industries due to supply chain disruptions and  
65 indirect losses such as the financial sector (Lunt et al., 2016; Maynard, 2015). However, more research  
66 and information about climate risk distributions and the connection of extreme weather events across  
67 the world is needed to estimate the likelihood of multiple breadbasket failures (Bailey and Benton,  
68 2015; Schaffnit-Chatterjee et al., 2010). This paper quantifies simultaneous climate risks to agricultural  
69 production in the global breadbaskets under 1.5 and 2°C warming scenarios. Whilst the difference of  
70 half a degree might be considered comparatively “small” on an aggregated global level, regional  
71 changes can be much larger (Seneviratne et al., 2016). Moreover, changes in extreme events and  
72 spatial dependence, which influence global risks such as multiple breadbasket failures, may expose  
73 significant differences between the two global mean temperature increments.

74 This paper uses initial results from the “Half a degree Additional warming, Prognosis and Projected  
75 Impacts” (HAPPI) project (Mitchell et al., 2016a). HAPPI provides a set of climate data specifically  
76 designed to address the Paris Agreement by simulating scenarios that are 1.5 and 2°C warmer than  
77 pre-industrial worlds. It provides a large enough ensemble of climate model runs to enable a thorough  
78 assessment of extreme weather and climate-related risks. Results will provide an important  
79 contribution to current climate policy discussions about differential impacts at specific global warming  
80 levels.

81 Our paper is organized as follows. In Section 2 we explain the HAPPI experiment and the HadAM3P  
82 model which was used in this study. In Section 3 we describe the climate indicators that have been  
83 used to assess agricultural risks and how we bias-corrected the data. We introduce the copula  
84 methodology used for the multivariate climate risk analysis in this paper and explain how we estimate  
85 the impact of climate risks on agricultural production. Section 4 shows the results, which will be  
86 further discussed in Section 5. The paper ends in Section 6, which summarizes our findings and gives  
87 an outlook to possible future work.

88

## 89 2 Data

90

### 91 2.1 *HadAM3P model*

92 Monthly precipitation and maximum temperature data are taken from the global atmosphere only  
93 model, HadAM3P (Massey et al., 2015; Pope et al., 2000). HadAM3P was developed by the UK Met  
94 Office Hadley Centre and is based on the atmosphere component of HadCM3, a coupled ocean-  
95 atmosphere model (Gordon et al., 2000). HadAM3P is run at a global resolution of 1.875° longitude  
96 and 1.25° latitude with a 15 minute time step. The model is run via the *climateprediction.net* volunteer  
97 distributed computed network (Anderson, 2004) and is, owing to its large ensemble size, well suited  
98 to analyse large-scale extreme weather events. Compared to other models from the same modelling  
99 family, HadAM3P has improvements in resolution and model physics (Pope and Stratton, 2002).  
100 Results of the HadAM3P distributed computing simulations are comparable to results of state of the  
101 art global climate model (GCM) simulations (Massey et al., 2015).

### 102 2.2 *HAPPI experiment*

103 HadAM3P is one of several models used for the HAPPI experiment (Mitchell et al., 2016a) which  
104 compares the statistics of extreme weather events simulated for a world which is 1.5 and 2 °C warmer  
105 than in pre-industrial (1861-1880) conditions with those of the present day. Driven by several leading  
106 atmosphere-only GCMs, HAPPI uses an experimental design similar to CMIP5 and is able to produce  
107 large simulation ensembles of high resolution global and regional climate data. Compared to CMIP5  
108 style experiments which use different Representative Concentration Pathways (RCPs) to reach a  
109 certain radiative forcing by 2100 and which contain uncertainty in climate model responses, HAPPI  
110 uses large sets of simulations under steady forcing conditions to calculate risks at certain warming  
111 levels irrespective of the emission pathway. Historical (in this study denoted with HIST) refers to the  
112 2006-2015 decade (which has already experienced a GMT rise of 0.85°C compared to pre-industrial  
113 levels (Fischer and Knutti, 2015)), a time period in which ocean temperatures have been

114 approximately constant but observed Sea Surface Temperatures (SSTs) contain a range of different  
115 patterns including El Nino patterns which were used to force the models. Each of the one-decade-  
116 simulations in the 50 to 100 member ensembles starts with a different initial weather state which  
117 results in 500 to 1000 years of model output. The 1.5°C warming experiment refers to conditions  
118 relevant for the 2106-2115 period and uses anthropogenic radiative forcing conditions from RCP2.6  
119 (Van Vuuren et al., 2011) which coincides with a global average temperature response of ~1.5°C  
120 relative to pre-industrial levels. Natural radiative forcings are used from the 2006-2015 decade. SSTs  
121 in the 1.5°C scenario are calculated by adding the mean projected CMIP5 RCP 2.6 SST changes across  
122 23 models averaged over the 2091-2100 period to observed 2006-2015 SSTs. The 2°C warming  
123 scenario refers to 2106-2115 conditions as well and uses a weighted average between the RCP2.5 and  
124 RCP4.6 scenarios to reach a ~2°C global mean temperature response, exactly 0.5°C warmer than the  
125 1.5°C scenario. Natural forcings again stay at 2006-2015 levels. For more information on the HAPPI  
126 experiment, see (Mitchell et al., 2016a).

127 Using *climateprediction.net*'s large ensemble modelling system, ~150, ~100 and ~120 ensemble  
128 members for the historical, 1.5 and 2°C scenario respectively were obtained. Owing to the large  
129 number of ensemble members with varied initial conditions, the HAPPI HadAM3P results used in this  
130 study are well suited to the analysis of extreme weather events with an improved representation of  
131 internal climate variability. Choosing a one-decade time period allows for assessment of the impacts  
132 of inter-annual climate variability on agricultural production. Note that the number of ensemble  
133 members differs as only ensemble members that were completed on the volunteers' computers could  
134 be included. Reasons for non-completion could be hardware failure or termination of the experiment  
135 by the volunteer (Massey et al., 2015).

136

137

138

139        *2.3 Historical crop yield and climate data*

140        This study focuses on climate risks to agricultural production in major global breadbaskets.  
141        Breadbaskets are important sub-national crop producing regions on a province/state scale in the US,  
142        Argentina, China, India and Australia for wheat and the US, Argentina, Brazil, China and India for maize  
143        and soybean (see details in Supplementary Information). The regions were chosen based on FAO  
144        (2015) country production statistics and official governmental statistics of subnational production. The  
145        highest crop producing provinces and states were selected with the premise that the provinces/states  
146        of a breadbasket have to be adjacent. For the analysis, the provinces/states were then aggregated to  
147        breadbaskets. Europe and Russia/Ukraine were excluded due to a lack of sufficiently long, subnational  
148        time-series data. Sub-national, annual crop yield data for all states/provinces of a breadbasket from  
149        1967 to 2012 (maize and soybean data in Brazil and India from 1977 and Argentina from 1970) were  
150        collected from official governmental databases (Australian Bureau of Statistics, 2015; Conab  
151        (Companhia Nacional De Abastecimento) Brazil, 2015; Ministerio de Agricultura, Ganaderia y Pesca de  
152        Argentina, 2015; Ministry of Agriculture and Farmers Welfare, Govt. of India, 2015; National Bureau  
153        of Statistics of China, n.d.; USDA, 2015). For the analysis, yield data was detrended using a four-  
154        parametric logistic function (Gaupp et al., 2016) which has the advantage that it can take on the form  
155        of a linear, exponential, damped or logistic trend. Detrended yield data and monthly Princeton re-  
156        analysis precipitation and maximum temperature data between 1967 and 2012 (Sheffield et al., 2006)  
157        were used to find region- and crop-specific relationships between climate and agricultural production.  
158        Princeton re-analysis data is a combination of a number of observation-based datasets and  
159        NCEP/NCAR re-analysis data and provides globally consistent, bias-corrected climate data.

160

161

162

163 3 Methods

164

165 *3.1 Climate indicator selection*

166 We identified climate indicators which significantly impact three important crops - wheat, maize and  
167 soybean - in five breadbaskets around the globe. A climate indicator is a crop and region specific  
168 variable based on either monthly maximum temperature or precipitation data which correlates with  
169 crop yields.

170 By concentrating on breadbaskets rather than using the national scale, the region-specific relationship  
171 between climate indicator and detrended yield could be determined. This is particularly relevant in  
172 large countries where crop production is concentrated in only a few regions. In order to find the most  
173 robust climate indicators for each crop and breadbasket, in a first step, an extensive literature review  
174 was carried out. Regional case studies were chosen in locations within or very close to the breadbasket  
175 areas used in this study. Indicators are mainly average maximum temperature or cumulative  
176 precipitation during the crop's growing season (e.g. June to November in India's soybean breadbasket)  
177 but also precipitation during the monsoon season (June to September in India's wheat breadbasket)  
178 which is stored in the soil and influences wheat growth from October to March (a table with detailed  
179 description of climate indicator selection and literature review is in Supplementary Information). In a  
180 second step, the choice of the climate indicator was validated through a correlation analysis between  
181 the climate re-analysis data and the observed, logistically de-trended (Gaupp et al., 2016) subnational  
182 crop yield data on state/province level using the Pearson correlation coefficient, shown in Table 1. The  
183 Pearson correlation coefficient is a widely used method to quantify the crop yield-climate relationship  
184 (Chen et al., 2014; Luo et al., 2005; Magrin et al., 2005; Podestá et al., 2009; Tao et al., 2008).

185 Depending on the value and significance of the correlation coefficient, one or two indicators per crop  
186 and breadbasket were chosen. In exceptional cases, an indicator was selected when Pearson's  $r$   
187 showed a non-significant but strong relationship pointing to the same direction as indicated in the

188 literature if it has been described as significant there. Differences can arise through differences  
189 between re-analysis data and locally observed climate data, different spatial scales or different  
190 statistical methods<sup>1</sup>. Figure 1 shows the indicator selection for each crop and breadbasket as well as  
191 the harvesting dates. For the analysis of climate risks, with a climate risk defined as a climate indicator  
192 exceeding a critical threshold, climate thresholds were set for each crop, breadbasket and indicator.  
193 A simple linear regression between each climate indicator and observed, detrended crop yield was  
194 used to define a temperature or precipitation threshold related to the lower 25% detrended yield  
195 percentile (see figure SF2 in Supplementary Information). We acknowledge that using a simple linear  
196 regression cannot account for the possibility of non-linear relationships between climate indicator and  
197 crop yield or the interaction between precipitation and temperature. Applying a simple linear  
198 regression allows one to identify the most relevant climate indicators for different crop yields (Tao,  
199 2008) which serves the purpose of this paper. Similar to other papers in the field (e.g. Lobell et al.,  
200 2011) this study does not aim to predict actual future yields but to estimate the future impact of  
201 climate on agricultural production. In contrast to process-based models (e.g. Asseng et al., 2015;  
202 Rosenzweig et al., 2014; Schleussner et al., 2016a), which represent key dynamic processes affecting  
203 crop yields, our approach is based on empirical relationships between location- and crop-specific  
204 climate indicators and crop yields. As Lobell and Asseng (2017) have shown, there are no systematic  
205 difference between the predicted sensitivities to warming between the two approaches up to 2°C  
206 warming. Empirical models are able to assess the climate-yield relationship location-specifically.  
207 Process-based models are typically better in understanding the interaction between crop genetics,  
208 management options and climate but might ignore factors that are important to crop growth in some  
209 seasons or specific environments.

---

<sup>1</sup> This is why in reports such as the IPCC reports (Allen et al., 2019; IPCC, 2014), different models are used and compared to give policy recommendations and model inter-comparison projects such as ISIMIP ([www.isimip.org](http://www.isimip.org)) or AgMIP ([www.agmip.org](http://www.agmip.org)) have been conducted. Lobell and Asseng (2017) compared process based and statistical crop models and found no systematic difference between predicted sensitivities to warming between the two model types up to a 2 degree warming.



210 To account for uncertainties in the sample statistics of the HAPPI data, the data were bootstrapped  
 211 1000 times for the threshold exceedance calculation. Results in Figure 2 show the simulation mean. A  
 212 breadbasket is experiencing a climate risks for a crop as soon as one of the temperature or  
 213 precipitation based indicators is exceeding the threshold. The breadbasket-specific relationship  
 214 between temperature and precipitation is accounted for through the copula correlation structure  
 215 explained in Section 3.3.

### 216 3.2 Bias-correction

217 In order to quantify the likelihood of threshold exceedance of different climate indicators, the  
 218 HadAM3P model output has to be comparable to the observed historical climate used for setting these  
 219 thresholds. Therefore, both historical and future experiment results were calibrated using a simple  
 220 bias-correction method (Hawkins et al., 2013; Ho, 2010) which corrects mean and variability biases of  
 221 the climate indicators distributions using the Princeton re-analysis data (Sheffield et al., 2006) as  
 222 calibration dataset:

223

$$I_{BC}(t) = \overline{O_{REF}} + \frac{\sigma_{O,REF}}{\sigma_{I,REF}} (I_{REF}(t) - \overline{I_{REF}}) \quad (1)$$

$$I_{FUT,BC}(t) = \overline{O_{REF}} + \frac{\sigma_{O,REF}}{\sigma_{I,REF}} (I_{FUT}(t) - \overline{I_{REF}}) \quad (2)$$

224

225

226  $I_{BC}$  denotes the HAPPI HadAM3P bias-corrected climate indicator,  $O_{REF}$  and  $I_{REF}$  the observational  
 227 Princeton dataset and HAPPI HadAM3P historical raw climate indicators and  $I_{FUT}$  represents the 1.5 or  
 228 2°C raw climate indicator. This method has the advantage of being simple to calculate and being  
 229 independent of the shape of the climate variable distribution (Hawkins et al., 2013). It is used widely  
 230 in agricultural modelling (Navarro-Racines et al., 2016). Although for precipitation usually a more

231 complicated calibration method has to be applied as it cannot take negative values, in this case it was  
232 possible as we use aggregated precipitation values which never reach zero. HadAM3P generally  
233 overestimated temperature compared to the Princeton dataset with HadAM3P maximum  
234 temperature being between 7 and 57% higher than Princeton in all breadbaskets. Precipitation is  
235 underestimated in the maize and soybean breadbaskets by between 2 and 30%. Precipitation for  
236 wheat, which has a different growing season, is both higher and lower than the reference dataset  
237 (between 40% lower in Australia and 37% higher in the US breadbasket).

238

### 239 *3.3 Regular vine copulas*

240 In this study, climate indicators based on historical Princeton re-analysis data were used to estimate  
241 the spatial dependence structure between the five breadbaskets to avoid biases in inter-regional  
242 correlation in the HadAM3P climate model. As the dependence structure of the HAdAM3P climate  
243 indicators in the different breadbaskets did not change between historical and warming scenarios, we  
244 kept the historical dependence structure constant in the 1.5 and 2°C scenarios. Changes in  
245 simultaneous climate risks between scenarios occur due to changes in mean and variance of the  
246 underlying marginal distributions of the climate indicators based on HadAM3P data.

247 In order to estimate risks of multiple breadbasket failure owing to joint climate extremes in major crop  
248 production areas<sup>2</sup>, the spatial dependence structure of the global breadbasket's climate indicators  
249 was modelled using regular vine (RVine) copulas (Aas et al., 2009; Dißmann et al., 2013; Kurowicka  
250 and Cooke, 2006). RVines are a flexible class of multivariate copulas which are able to model complex  
251 dependencies in larger dimensions. They are based on Sklar's theorem (Sklar, 1959) which states that  
252 any multivariate distribution  $F$  can be written as

---

<sup>2</sup> We acknowledge that heterogeneity is lost with aggregation to breadbaskets. However, we made sure that the relationship between climate indicators and yields were robust between our states/provinces, the aggregated breadbasket scale and local studies taken from the literature.

253

$$F(x_1, \dots, x_n) = C[F_1(x_1), \dots, F_n(x_n)] \quad (3)$$

254 with marginal probability distributions  $F_1(x_1), \dots, F_n(x_n)$  and  $C$  denoting an  $n$ -dimensional copula, a  
255 multivariate distribution on the unit hypercube  $[0,1]^2$  with uniform marginal distributions. Vine  
256 copulas are constructed using conditional and unconditional bivariate pair-copulas from a set of  
257 copula families with distinct dependence structures (Aas et al., 2009; Joe, 1997). A set of linked RVine  
258 trees describes the factorisation of the copula's multivariate density function (Bedford and Cooke,  
259 2002). An  $n$ -dimensional RVine model consists of  $(n-1)$  trees including  $N_i$  nodes and  $E_{i-1}$  edges which  
260 join the nodes. The tree structure is built according to the proximity condition which means that if an  
261 edge connects two nodes in tree  $j+1$ , the corresponding edges in tree  $j$  share a node (Bedford and  
262 Cooke, 2002). The first tree consist of  $n-1$  pairs of variables with directly modelled distributions. The  
263 second tree identifies  $n-2$  variable pairs with a distribution modelled by a pair-copula conditional on a  
264 single variable which is determined in the second tree. Proceeding in this way, the last tree consist of  
265 a single pair of variables with a distribution conditional on all remaining variables, defined by a last  
266 pair-copula (Dißmann et al., 2013). The RVine tree structure, the pair-copula families and the copula  
267 parameters are estimated in an automated way starting with the first tree. The tree is selected using  
268 a maximum spanning tree algorithm and Kendall's tau as edge weights. The best fitting pair-copula  
269 family is chosen using the Akaike Information Criterion (Akaike, 1973) and copula parameters are  
270 estimated using Maximum Likelihood Estimation (MLE). In this study we chose from six different  
271 copula families representing different types of tail dependencies to capture the exact patterns of  
272 dependence between the different climate indicators in the crop breadbaskets: Gaussian, Clayton,  
273 Student-t, Gumbel, Joe and Frank copulas (Nelsen, 2007).

274

275 *3.4 Impact on agricultural production*

276 We analyse events where the climatic conditions in all five breadbaskets are associated with losses in  
 277 agricultural yields. We identify a ‘breadbasket failure’ event as being when the climatic conditions are  
 278 at least as severe as those conditions associated with the 25 percentile of the logistically detrended  
 279 yields (with detrended yields as residuals of the non-linear logistic regression with a residual mean  
 280 equal to zero). The crop production loss for an event of this severity is the 25 percentile of the  
 281 logistically detrended yield multiplied with the 2012 harvested area. Given that we identify climatic  
 282 events that are *at least* as severe as this condition, our estimated loss is the lower bound on the loss,  
 283 i.e. the minimum expected loss. Minimum expected losses are then defined as the sum of crop losses  
 284 in all five breadbaskets multiplied with the joint probability that climate thresholds are exceeded in  
 285 all regions simultaneous as shown in Equation 4:

$$286 \quad \textit{Minimum expected losses} \geq \sum_i^{BB} (|y_{25_i}| \cdot \textit{area}_{i,2012}) \cdot p_5 \quad (4)$$

287 with

$$288 \quad p_5 = P(\textit{Clim}_1 \geq t_{\textit{clim}_1}, \textit{Clim}_2 \geq t_{\textit{clim}_2}, \textit{Clim}_3 \geq t_{\textit{clim}_3}, \textit{Clim}_4 \geq t_{\textit{clim}_4}, \textit{Clim}_5 \geq t_{\textit{clim}_5})$$

$$289 \quad = C[F_1(t_{\textit{clim}_1}), F_2(t_{\textit{clim}_2}), F_3(t_{\textit{clim}_3}), F_4(t_{\textit{clim}_4}), F_5(t_{\textit{clim}_5})]$$

291

292 with  $y_{25_i}$  as the 25 percentile of logistically detrended yields in the breadbasket  $i$  which was used to  
 293 define climate thresholds and which indicates a minimum yield loss,  $\textit{area}_{i,2012}$  as the 2012 harvested  
 294 area in breadbasket  $i$  and with  $p_5$  as the probability of all five breadbaskets exceeding the climate  
 295 thresholds in the same year.  $\textit{Clim}_i$  denotes the temperature or precipitation-based climate indicator,  
 296 associated with the 25 percentile of the detrended yields. In case that a breadbasket has two  
 297 indicators for a crop, the exceedance of at least one of the climate thresholds  $t_{\textit{clim}_i}$  is counted as  
 298 threshold exceedance in the breadbasket.  $C$  denotes the copula.

299

300 4 Results

301

302 *4.1 Changes in climate risks to agriculture under 1.5 and 2°C global warming*

303 The change of climate risks to major crops in the global breadbaskets were examined for each region  
304 and crop separately comparing historical risks with risks associated with a 1.5 and 2°C global warming,  
305 shown in Figure 2. As expected from an increase of global mean temperature, temperature based  
306 climate risks are increasing, but to different extents depending on the region. Precipitation signals  
307 associated with 1.5 and 2°C warming are less clear. While precipitation based climate risks in the US  
308 and Brazil increase in both scenarios for the summer crops maize and soybean, precipitation in  
309 Argentina does not significantly change. Risks in China and India decrease due to an increase in  
310 monsoon precipitation. For wheat, precipitation-based climate risks only increase in Australia.

311 The decrease of precipitation-based climate risks to wheat in the US and China, and the increase in  
312 the Australian breadbaskets for both warming scenarios mostly coincide with findings of a previous  
313 study (Gaupp et al., in review) which examined climate risk trends in the past. In India and China,  
314 wheat is indirectly impacted by the summer monsoon rainfall which provides stored soil moisture for  
315 the “rabi” wheat crop. Although precipitation between June and September in the Chinese  
316 breadbasket showed a decrease in the recent past, in a 1.5 and 2°C warmer world precipitation during  
317 monsoon months in the Chinese breadbasket is projected to increase. This coincides with (Lv et al.,  
318 2013) who project a decrease in precipitation in China during the wheat growing season between the  
319 2000s and 2030s and a consistent precipitation increase from the 2030s to the 2070s. In India, rainfall  
320 during summer monsoon months (June to September) showed a decreasing decadal trend in the  
321 recent past (Guhathakurta et al., 2015) which was reflected in an increasing climate risk for wheat in  
322 India in the past (Gaupp et al., in review). In the future, however, monsoon precipitation is projected  
323 to increase under all RCP scenarios in CMIP5 projections (Jayasankar et al., 2015; Menon et al., 2013)

324 which coincides with decreasing precipitation climate risks to wheat in the Indian breadbasket found  
325 in this study. However, precipitation-based risks in India and China might be underestimated in this  
326 study because of the HAPPI experiment structure which has fixed SSTs driving the model, rather than  
327 a fully couple ocean simulation. This often leads to variability in land-ocean driven cycles not changing  
328 much and thereby to an underestimation of precipitation variability during the monsoon months.  
329 CMIP5 models project both increasing and decreasing standard deviations of monsoon precipitation  
330 in India for RCP 2.6 and 4.5. In Australia, precipitation in the wheat growing season is projected to  
331 decrease following different CMIP5 models under RCP4.5 (Ummenhofer et al., 2015) which our study  
332 confirms through increased precipitation-based climate risks. Temperature risks are increasing in all  
333 temperature sensitive breadbaskets with stronger increases in India and Australia than in Argentina.  
334 Our estimates of climate risks to wheat production coincide with results of crop model experiments  
335 in other studies. Asseng et al. (2015) compared results of 30 wheat crop simulation models in 30 main  
336 wheat producing locations without water stress, focussing only on the effect of temperature. All  
337 models showed yield losses at a 2°C warming, which coincides with our temperature-based climate  
338 risk increases in India, Australia and Argentina. Rosenzweig et al. (2018) and Ruane et al. (2018) used  
339 HAPPI climate data and other climate model experiments from CMIP5 to compare climate impacts on  
340 crops under a 1.5°C and 2°C warming using process-based crop models. They found wheat yield losses  
341 smaller than 5% in the North American Great Plains, but larger losses in Australia and Argentina under  
342 1.5°C warming. In India and China the models showed yield increases in a 1.5°C world. Challinor et al.  
343 (2014) came to similar conclusions in a meta-analysis of crop yield under climate change. He found no  
344 changes in wheat yields under a 1.5°C warming in tropical regions but a slight decrease under 2°C. In  
345 temperate regions, such as the US, China or Argentina, wheat yields are projected to decrease for both  
346 warming levels, when adaptation strategies such as irrigation, planting times of crop varieties are not  
347 considered.  
348

349 For soybean, precipitation-based climate risks in South America increase in Brazil but do not change  
350 notably in Argentina. This coincides with findings from other CMIP5 studies (Barros et al., 2015; IPCC,  
351 2014). In the US, CMIP5 models show a small, not significant increase in annual precipitation (IPCC,  
352 2014)) which can be seen in HadAM3P as well. Precipitation during the soybean growing season, on  
353 the other hand, is projected to decrease in both 1.5 and 2°C scenarios which results in higher climate  
354 risks. In China and India, soybean growing seasons are directly aligned with the summer monsoon.  
355 Hence, precipitation-based soybean climate risks decrease due to the above discussed increase in  
356 monsoon precipitation. Temperature based risks, on the other hand, increase significantly in the US,  
357 Argentina and India. Those temperature and precipitation changes translated into yield changes in  
358 several crop model experiments for rainfed and irrigated soybean. The models show slight yield  
359 decreases over the interior of Northern America but small increases towards the eastern US in a 1.5°C  
360 scenario for rainfed soybean. In Brazil and Argentina, soybean shows both increases and decreases  
361 under a 1.5°C warming and in the Indian breadbasket, soybean yields are projected to increase. In  
362 China, yields are projected to increase in the North, but decrease in the South. Models for irrigated  
363 crop that also include CO<sub>2</sub> benefits, yields are projected to increase (Ruane et al., 2018). Under a 2°C  
364 warming, GCMs revealed yield increases when CO<sub>2</sub> effects were considered as they largely overcome  
365 increased temperature risks (Ruane et al., 2018; Schleussner et al., 2016a).

366

367 For maize, climate risks show very similar patterns to soybean as the two summer crops have similar  
368 growing seasons and indicators. Additional to the soybean climate indicators, maize in the Chinese  
369 breadbasket is sensitive to temperature. Owing to those local precipitation changes and temperature  
370 rise, global crop models (GCMs) have shown declines in maize yields in all five breadbaskets in both a  
371 1.5 and 2°C warmer world (Ruane et al., 2018; Schleussner et al., 2016a). In contrast to soybean, maize  
372 is not able to capture the same level of CO<sub>2</sub> benefits and hence yields decrease further under in a 2°C  
373 world. Those finding coincide with results of the meta-study by Challinor et al. (2014).

374 One of the major concerns in studies of the difference between a 1.5 and 2°C global warming is the  
375 significance of the difference between the temperature increments (James et al., 2017). The  
376 difference between climate risks for 1.5 and 2°C in this study was tested with the student two-sample  
377 Kolmogorov-Smirnov (KS) test which tests the null hypothesis that both distributions of resampled  
378 threshold exceedance are drawn from the same distribution. Results showed significant differences  
379 for all indicators and crops at the 0.001 significance level between the two warming levels. The KS test  
380 allows for robust statements about the difference between climate risks under 1.5 and 2°C warming  
381 even if there is an overlap of uncertainty bands (Schleussner et al., 2016a). Error bars are small  
382 compared to the absolute change in climate threshold exceedance with the exception of precipitation  
383 risks in Argentina for soybean and maize. Figure 2 also compares the difference in changes from  
384 historical climate for both global mean temperature increases. Across all three crops, we found  
385 stronger signals for temperature based risks than for precipitation based risks which show smaller,  
386 both positive and negative signals. Additionally, the difference between the 1.5 and 2°C warming is  
387 more pronounced in temperature based indicators with the largest difference in the Indian soybean  
388 breadbasket (26% points). The difference in precipitation risk changes between the two warming  
389 scenarios lies between 0 and 6% points. What stands out is the difference between 1.5 and 2°C for  
390 precipitation risks in Brazil. In contrast to other climate indicators, precipitation between December  
391 and February and March in Brazil shows a significantly stronger difference from historical data to 1.5°C  
392 than to 2°C.

#### 393 *4.2 Increasing risks of multiple breadbasket failure*

394 Having analysed individual changes of climate risks in the global wheat, soybean and maize  
395 breadbaskets for 1.5 and 2°C enables us to calculate joint climate risks on a global scale. Figure 3 shows  
396 the largest increase in risks of simultaneous crop failure (resulting from climate exceeding a crop- and  
397 region-specific threshold) in the global breadbaskets for maize, followed by soybean and wheat. For  
398 all three crops the likelihoods that none or just one of the breadbaskets experiences climate risks



399 decreases to (nearly) zero. For wheat and soybean, the likelihoods of breadbaskets experiencing  
400 detrimental climate change increases significantly from the historical scenario to 1.5°C and even more  
401 assuming 2°C warming. The figure can be interpreted as a discrete probability distribution with the  
402 sum of all breadbasket threshold exceedances adding up to 1. The shape of the distribution stays  
403 roughly the same across warming scenarios with higher probabilities that parts of the breadbaskets  
404 exceed the thresholds and smaller likelihoods in the extremes. While the historical baseline climate  
405 still shows the probability for zero simultaneous climate risks, for higher temperature scenarios these  
406 likelihoods disappear. The average threshold exceedance increases significantly (measured using the  
407 KS-test), more for soybean than for wheat. For maize, likelihoods of simultaneous climate risks  
408 increase strongly. Under the 2°C scenario the likelihood of all five breadbaskets suffering detrimental  
409 climate is the highest. For wheat, which shows the smallest simultaneous climate risks, the return  
410 period for all five breadbaskets exceeding their climate thresholds decreases from 43 years (or 0.023  
411 annual probability under historical conditions to 21 years (0.047) in a 1.5°C scenario and further down  
412 to around 15 years (0.066) under 2°C. Soybean has a return period of simultaneous climate risks in all  
413 breadbaskets of around 20 years (0.049 today which decreases to 9 (0.116) and 7 years (0.143 in a 1.5  
414 and 2°C warmer world respectively. Maize risks are highest in our study with an initial return period  
415 of 16 years (0.061), decreasing to less than 3 (0.39) and less than 2 years (0.538) under future global  
416 warming. In general, one can say that whilst the differences in yield at 1.5 vs 2°C are significant they  
417 are not as large as the difference between 1.5 and historical. Risk of simultaneous crop failure,  
418 however, do increase disproportionately between 1.5 and 2 degrees and this is important because  
419 correlated risks lead to disproportionately high impacts.

420 To illustrate the effects of simultaneous climate risks in a 1.5 and a 2°C warmer world, we estimated  
421 the impacts on agricultural production. Simultaneous crop failure in all breadbaskets, defined as the  
422 25 percentile of detrended yield, would add up to at least 9.86 million tons of soybean losses, 19.75  
423 million tons of maize losses and 8.59 million tons of wheat losses assuming 2012 agricultural area.  
424 Historical examples of global crop production shocks include 7.2 million tons soybean losses in

425 1988/99 and 55.9 million tons maize losses in 1988 which were mostly caused by low rainfall and high  
426 temperatures during summer growing season in the US (Bailey and Benton, 2015). Historical global  
427 wheat production shocks include 36.6 million tons wheat losses in 2003 mostly caused by heat waves  
428 and drought in spring in Europe and Russia but also by reduced acreage due to drought or winterkill  
429 in Europe, India and China (Bailey and Benton, 2015). Maize and wheat losses in this study are lower  
430 than in historical cases as our breadbaskets only account for 38% and 52% of global production  
431 respectively. Soybean in this study accounts for 80% of global production. Combining absolute losses  
432 with likelihoods of simultaneous climate risks, we calculated expected crop losses following Equation  
433 4. For all three crops, expected crop losses are significantly higher under the 2°C than under the 1.5°C  
434 scenario. Under a scenario of 2°C mean global warming, expected wheat, maize and soybean losses  
435 are projected to be 161 000, 2 753 000 and 265 000 tonnes higher than if global temperature  
436 increases are limited to 1.5°C. This equals total annual maize production in Uganda, the world's 33<sup>rd</sup>  
437 largest maize producer in 2012. The difference of wheat losses is larger than Bolivia's annual total  
438 production in 2012 (145 000 tonnes) and the increase of expected soybean losses is comparable to  
439 Mexico's annual production (248 000 tonnes), the world's 20<sup>th</sup> biggest soybean producer (FAO, 2015).

440 To test for the influence of inter-dependence between the climate indicators in the different  
441 breadbaskets on the results of this analysis, we excluded the correlations between them. We assumed  
442 independence between the breadbaskets, but still accounted for the negative correlation between  
443 temperature and precipitation indices within one breadbasket. Supplementary Figure SF3 illustrates  
444 the difference between independence and correlation. Between the three crops, no consistent  
445 pattern was found between dependent and independent cases. The only crop that shows significant  
446 differences is soybean with smaller likelihoods in the extremes when dependence is excluded. This  
447 means that the likelihood of all five soybean breadbaskets experiencing detrimental climate in one  
448 year is underestimated if correlations between the breadbaskets are not considered in a risk analysis.  
449 Expressed in expected production losses, the losses are up to 190 000 tonnes higher in the dependent

450 case which is more than what the 22<sup>nd</sup> largest soybean producer harvests annually (FAO, 2015). For  
451 wheat and maize, the difference between the dependencies was mostly not significant.

452

## 453 5 Discussion

454 Our results illustrate future climate conditions under two warming scenarios in the global  
455 breadbaskets and investigate simultaneous climate risks affecting three major crops. The study  
456 focused explicitly on the climate impact on crop yields. The effects of other factors such as soil quality,  
457 land management, land use or technology were held constant under future warming scenarios.  
458 Therefore, our estimates of crop production losses have to be interpreted with care. By not explicitly  
459 including CO<sub>2</sub> concentrations, for instance, the CO<sub>2</sub> fertilizer effect which increases productivity in  
460 wheat and soybean and to a certain extent in maize (Schleussner et al., 2016a) was not taken into  
461 account. The effects of climatic change on plant phenology were not considered. In China, for instance,  
462 the flowering date of wheat is projected to advance owing to increased temperatures and the gain-  
463 filling period will shorten which might further reduce yields (Lv et al., 2013). By holding harvested area  
464 constant at 2012 levels, shifts in land use and cropped area in response to projected climatic changes  
465 (Nelson et al., 2014; Schmitz et al., 2014) were not considered. Owing to a lack of subnational historic  
466 time series of irrigated crop yields, irrigation was not specifically taken into account in setting climate  
467 risk thresholds. This was acceptable in this study as, even without considering irrigation, the  
468 correlation coefficients between observed, detrended yields and climate indicators were mostly  
469 significant. A large share of the regions in this study are completely rain-fed. In other regions such as  
470 India or the US, irrigated crops still show correlations with rainfall (Pathak and Wassmann, 2009) or  
471 no significant difference to rain-fed crops at all (Zhang et al., 2015). Results of the analysis of  
472 simultaneous climate risks may vary depending on the climate indicator selection. The two-step  
473 approach of pre-selecting potential indicators in a literature review and verification through the  
474 correlation analysis with re-analysis climate data and observed historical yield data represents a

475 robust way of indicator selection. However, including different climate variables such as number of  
476 days above a crop dependent heat threshold (Schlenker and Roberts, 2009; Tack et al., 2015; Zhang  
477 et al., 2015) or dry spell length (Hernandez et al., 2015; Ramteke et al., 2015; Schleussner et al., 2016a)  
478 might lead to different results. So far, the HAPPI project only provides monthly data which limited the  
479 climate variable choice. In order to reduce uncertainties, we bootstrapped the climate indicators and  
480 repeatedly simulated the copula models. However, results from 1.5 and 2°C warming scenarios vary  
481 between different GCMs (Schleussner et al., 2016a). A comparison with additional climate models  
482 from the HAPPI project will further improve the robustness of the results.

483

## 484 6 Conclusion

485 This study found disproportionally increasing future risks of simultaneous crop failure in the global  
486 wheat, maize and soybean breadbaskets in a 1.5 and 2°C warmer world using results of the HadAM3P  
487 atmospheric model as part of the HAPPI experiment. Increases in temperature-based climate risks  
488 were found to be stronger than precipitation-based risks which showed different signals depending  
489 on crop and region. Using the copula methodology, it was possible to capture dependence structures  
490 between regions and to calculate joint climate risks in the major crop producing areas. Additionally,  
491 the copula analysis accounted for the region-specific relationships between temperature and  
492 precipitation. Strongest increases in simultaneous climate risks were found for maize where return  
493 periods of simultaneous crop failure decrease from 16 years in the past to less than 3 and less than 2  
494 years under 1.5 and 2°C warming. In percentage terms, the largest increase of simultaneous climate  
495 threshold exceedance in all five breadbaskets between the two warming scenarios was found for  
496 wheat (40%), followed by maize (35%) and soybean (23%). Looking at the impacts on crop production,  
497 the study showed that limiting global warming to 1.5°C would avoid production losses of up to  
498 2 753 million (161 000, 265 000) tonnes maize (wheat, soybean) in the main production regions.

499 Our study represents an important first step in the analysis of differential temperature increases of  
500 1.5 and 2°C and their impacts on agricultural production. Compared to climate studies which often  
501 focus on average annual values, this study focused on crop growth periods which may show opposite  
502 signals to annual means – as shown here for soybean in the US - and therefore added valuable  
503 information to existing studies.

504 Results are based on HadAM3P, the first model in the HAPPI experiment set up. Including outputs  
505 from additional climate models will give more robust information on future climate risks. Additionally,  
506 further analysis of the ability of climate models to accurately model spatial dependence between  
507 regions is needed. This study used historical dependence to avoid biases in spatial correlation and kept  
508 dependence constant under future scenarios. Some literature, however, suggests that teleconnection  
509 patterns might change, i.e. owing to changes in El Niño Southern Oscillation (ENSO) (Cai et al., 2014;  
510 Power et al., 2013), which could then alter the spatial climate dependence structure in the  
511 breadbaskets. Future work (under preparation) will look into climate risks under different ENSO  
512 phases.

513 This paper provides insights into risks of multiple breadbasket failure under 1.5 and 2°C warming  
514 which can contribute to current climate policy discussions and potentially provides useful information  
515 for the Intergovernmental Panel on Climate Change (IPCC) Special Report on the impact of 1.5°C global  
516 warming commissioned by the UN-FCCC after the Paris Agreement.

517

## 518 **References**

- 519 Aas, K., Czado, C., Frigessi, A., Bakken, H., 2009. Pair-copula constructions of multiple dependence.  
520 *Insur. Math. Econ.* 44, 182–198. <https://doi.org/10.1016/j.insmatheco.2007.02.001>
- 521 Akaike, H., 1973. Information theory and an extension of the maximum likelihood principle. Á In:  
522 Petran, BN and Csáki, F, in: *International Symposium on Information Theory, Second Edition.*  
523 *Akadémiai Kiadi, Budapest, Hungary, Pp. 267Á281.*
- 524 Allen, M., Antwi-Agyei, P., Aragon-Durand, F., Babiker, M., Bertoldi, P., Bind, M., Brown, S.,  
525 Buckeridge, M., Camilloni, I., Cartwright, A., 2019. Technical Summary: Global warming of  
526 1.5° C. An IPCC Special Report on the impacts of global warming of 1.5° C above pre-  
527 industrial levels and related global greenhouse gas emission pathways, in the context of

528 strengthening the global response to the threat of climate change, sustainable development,  
529 and efforts to eradicate poverty.

530 Anderson, D.P., 2004. Boinc: A system for public-resource computing and storage, in: Grid  
531 Computing, 2004. Proceedings. Fifth IEEE/ACM International Workshop On. IEEE, pp. 4–10.

532 Asseng, S., Ewert, F., Martre, P., Rötter, R.P., Lobell, D.B., Cammarano, D., Kimball, B.A., Ottman,  
533 M.J., Wall, G.W., White, J.W., Reynolds, M.P., Alderman, P.D., Prasad, P.V.V., Aggarwal, P.K.,  
534 Anothai, J., Basso, B., Biernath, C., Challinor, A.J., De Sanctis, G., Doltra, J., Fereres, E., Garcia-  
535 Vila, M., Gayler, S., Hoogenboom, G., Hunt, L.A., Izaurralde, R.C., Jabloun, M., Jones, C.D.,  
536 Kersebaum, K.C., Koehler, A.-K., Müller, C., Naresh Kumar, S., Nendel, C., O’Leary, G., Olesen,  
537 J.E., Palosuo, T., Priesack, E., Eyshi Rezaei, E., Ruane, A.C., Semenov, M.A., Shcherbak, I.,  
538 Stöckle, C., Stratonovitch, P., Streck, T., Supit, I., Tao, F., Thorburn, P.J., Waha, K., Wang, E.,  
539 Wallach, D., Wolf, J., Zhao, Z., Zhu, Y., 2015. Rising temperatures reduce global wheat  
540 production. *Nat. Clim. Change* 5, 143–147. <https://doi.org/10.1038/nclimate2470>

541 Australian Bureau of Statistics, 2015. Historical Selected Agriculture Commodities. URL  
542 <http://www.abs.gov.au/>

543 Bailey, R., Benton, T., 2015. Extreme weather and resilience of the global food system. Final Project  
544 Report from the UK-US Taskforce on Extreme Weather and Global Food System Resilience,  
545 The Global Food Security programme, UK.

546 Barros, V.R., Boninsegna, J.A., Camilloni, I.A., Chidiak, M., Magrín, G.O., Rusticucci, M., 2015. Climate  
547 change in Argentina: trends, projections, impacts and adaptation: Climate change in  
548 Argentina. *Wiley Interdiscip. Rev. Clim. Change* 6, 151–169. <https://doi.org/10.1002/wcc.316>

549 Bedford, T., Cooke, R.M., 2002. Vines: A new graphical model for dependent random variables. *Ann.*  
550 *Stat.* 1031–1068.

551 Bren d’Amour, C., Wenz, L., Kalkuhl, M., Christoph Steckel, J., Creutzig, F., 2016. Teleconnected food  
552 supply shocks. *Environ. Res. Lett.* 11, 035007. [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/11/3/035007)  
553 [9326/11/3/035007](https://doi.org/10.1088/1748-9326/11/3/035007)

554 Cai, W., Borlace, S., Lengaigne, M., Van Rensch, P., Collins, M., Vecchi, G., Timmermann, A., Santoso,  
555 A., McPhaden, M.J., Wu, L., others, 2014. Increasing frequency of extreme El Niño events  
556 due to greenhouse warming. *Nat. Clim. Change* 4, 111–116.

557 Challinor, A.J., Watson, J., Lobell, D.B., Howden, S.M., Smith, D.R., Chhetri, N., 2014. A meta-analysis  
558 of crop yield under climate change and adaptation. *Nat. Clim. Change* 4, 287.

559 Chen, H., Wang, J., Huang, J., 2014. Policy support, social capital, and farmers’ adaptation to drought  
560 in China. *Glob. Environ. Change* 24, 193–202.  
561 <https://doi.org/10.1016/j.gloenvcha.2013.11.010>

562 Conab (Companhia Nacional De Abastecimento) Brazil, 2015. Séries históricas. URL  
563 <http://www.conab.gov.br>

564 Dißmann, J., Brechmann, E.C., Czado, C., Kurowicka, D., 2013. Selecting and estimating regular vine  
565 copulae and application to financial returns. *Comput. Stat. Data Anal.* 59, 52–69.  
566 <https://doi.org/10.1016/j.csda.2012.08.010>

567 FAO, 2015. Statistical database.

568 FAO (Ed.), 2014. Strengthening the enabling environment for food security and nutrition, The state  
569 of food insecurity in the world. FAO, Rome.

570 Field, C.B., IPCC (Eds.), 2012. Managing the risks of extreme events and disasters to advance climate  
571 change adaption: special report of the Intergovernmental Panel on Climate Change.  
572 Cambridge University Press, New York, NY.

573 Fischer, E.M., Knutti, R., 2015. Anthropogenic contribution to global occurrence of heavy-  
574 precipitation and high-temperature extremes. *Nat. Clim. Change* 5, 560–564.  
575 <https://doi.org/10.1038/nclimate2617>

576 Fraser, E.D.G., Simelton, E., Termansen, M., Gosling, S.N., South, A., 2013. “Vulnerability hotspots”:  
577 Integrating socio-economic and hydrological models to identify where cereal production

578 may decline in the future due to climate change induced drought. *Agric. For. Meteorol.* 170,  
579 195–205. <https://doi.org/10.1016/j.agrformet.2012.04.008>

580 Gaupp, F., Pflug, G., Hochrainer-Stigler, S., Hall, J., Dadson, S., 2016. Dependency of Crop Production  
581 between Global Breadbaskets: A Copula Approach for the Assessment of Global and  
582 Regional Risk Pools. *Risk Anal.*

583 Gordon, C., Cooper, C., Senior, C.A., Banks, H., Gregory, J.M., Johns, T.C., Mitchell, J.F., Wood, R.A.,  
584 2000. The simulation of SST, sea ice extents and ocean heat transports in a version of the  
585 Hadley Centre coupled model without flux adjustments. *Clim. Dyn.* 16, 147–168.

586 Guhathakurta, P., Rajeevan, M., Sikka, D.R., Tyagi, A., 2015. Observed changes in southwest  
587 monsoon rainfall over India during 1901-2011: TREND IN SOUTHWEST MONSOON RAINFALL  
588 OVER INDIA. *Int. J. Climatol.* 35, 1881–1898. <https://doi.org/10.1002/joc.4095>

589 Hawkins, E., Osborne, T.M., Ho, C.K., Challinor, A.J., 2013. Calibration and bias correction of climate  
590 projections for crop modelling: An idealised case study over Europe. *Agric. For. Meteorol.*  
591 170, 19–31. <https://doi.org/10.1016/j.agrformet.2012.04.007>

592 Hernandez, V., Moron, V., Riglos, F.F., Muzi, E., 2015. Confronting farmers' perceptions of climatic  
593 vulnerability with observed relationships between yields and climate variability in Central  
594 Argentina. *Weather Clim. Soc.* 7, 39–59.

595 Ho, C.K., 2010. Projecting extreme heat-related mortality in Europe under climate change.

596 IPCC, 2014. *Climate Change 2014—Impacts, Adaptation and Vulnerability: Regional Aspects.*  
597 Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental  
598 Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New  
599 York, NY, USA.

600 James, R., Washington, R., Schleussner, C.-F., Rogelj, J., Conway, D., 2017. Characterizing half-a-  
601 degree difference: a review of methods for identifying regional climate responses to global  
602 warming targets: Characterizing half-a-degree difference. *Wiley Interdiscip. Rev. Clim.*  
603 *Change* e457. <https://doi.org/10.1002/wcc.457>

604 Jayasankar, C.B., Surendran, S., Rajendran, K., 2015. Robust signals of future projections of Indian  
605 summer monsoon rainfall by IPCC AR5 climate models: Role of seasonal cycle and  
606 interannual variability: FUTURE PROJECTIONS OF ISMR. *Geophys. Res. Lett.* 42, 3513–3520.  
607 <https://doi.org/10.1002/2015GL063659>

608 Joe, H., 1997. *Multivariate models and multivariate dependence concepts.* CRC Press.

609 Kurowicka, D., Cooke, R.M., 2006. *Uncertainty analysis with high dimensional dependence*  
610 *modelling.* John Wiley & Sons.

611 Lobell, D.B., Asseng, S., 2017. Comparing estimates of climate change impacts from process-based  
612 and statistical crop models. *Environ. Res. Lett.* 12, 015001.

613 Lobell, D.B., Schlenker, W., Costa-Roberts, J., 2011. Climate trends and global crop production since  
614 1980. *Science* 333, 616–620.

615 Lunt, T., Jones, A.W., Mulhern, W.S., Lezaks, D.P.M., Jahn, M.M., 2016. Vulnerabilities to agricultural  
616 production shocks: An extreme, plausible scenario for assessment of risk for the insurance  
617 sector. *Clim. Risk Manag.* 13, 1–9. <https://doi.org/10.1016/j.crm.2016.05.001>

618 Luo, Q., Bellotti, W., Williams, M., Bryan, B., 2005. Potential impact of climate change on wheat yield  
619 in South Australia. *Agric. For. Meteorol.* 132, 273–285.  
620 <https://doi.org/10.1016/j.agrformet.2005.08.003>

621 Lv, Z., Liu, X., Cao, W., Zhu, Y., 2013. Climate change impacts on regional winter wheat production in  
622 main wheat production regions of China. *Agric. For. Meteorol.* 171, 234–248.

623 Magrin, G.O., Travasso, M.I., Rodríguez, G.R., 2005. Changes in climate and crop production during  
624 the 20th century in Argentina. *Clim. Change* 72, 229–249.

625 Massey, N., Jones, R., Otto, F.E.L., Aina, T., Wilson, S., Murphy, J.M., Hassell, D., Yamazaki, Y.H., Allen,  
626 M.R., 2015. weather@home-development and validation of a very large ensemble modelling  
627 system for probabilistic event attribution: weather@home. *Q. J. R. Meteorol. Soc.* 141,  
628 1528–1545. <https://doi.org/10.1002/qj.2455>

629 Maynard, T., 2015. Food System Shock: The Insurance Impacts of Acute Disruption to Global Food  
630 Supply. Lloyd's of London., London, UK.

631 Menon, A., Levermann, A., Schewe, J., Lehmann, J., Frieler, K., 2013. Consistent increase in Indian  
632 monsoon rainfall and its variability across CMIP-5 models. *Earth Syst. Dyn.* 4, 287–300.  
633 <https://doi.org/10.5194/esd-4-287-2013>

634 Ministerio de Agricultura, Ganaderia y Pesca de Argentina, 2015. Statistical database. URL  
635 <http://www.siaa.gov.ar/>

636 Ministry of Agriculture and Farmers Welfare, Govt. of India, 2015. Crop Production Statistics. URL  
637 <http://eands.dacnet.nic.in/>

638 Mitchell, D., AchutaRao, K., Allen, M., Bethke, I., Forster, P., Fuglestedt, J., Gillett, N., Haustein, K.,  
639 Iverson, T., Massey, N., Schleussner, C.-F., Scinocca, J., Seland, Ø., Shiogama, H., Shuckburgh,  
640 E., Sparrow, S., Stone, D., Wallom, D., Wehner, M., Zaaboul, R., 2016a. Half a degree  
641 Additional warming, Projections, Prognosis and Impacts (HAPPI): Background and  
642 Experimental Design. *Geosci. Model Dev. Discuss.* 1–17. <https://doi.org/10.5194/gmd-2016-203>

644 Mitchell, D., James, R., Forster, P.M., Betts, R.A., Shiogama, H., Allen, M., 2016b. Realizing the  
645 impacts of a 1.5 [deg] C warmer world. *Nat. Clim. Change*.

646 National Bureau of Statistics of China, n.d. Regional data. URL <http://data.stats.gov.cn/>

647 Navarro-Racines, C.E., Tarapues-Montenegro, J.E., Ramírez-Villegas, J.A., 2016. BIAS-CORRECTION IN  
648 THE CCAFS-CLIMATE PORTAL: A DESCRIPTION OF MEHOTODOLOGIES.

649 Nelsen, R.B., 2007. An introduction to copulas. Springer Science & Business Media.

650 Nelson, G.C., Valin, H., Sands, R.D., Havlík, P., Ahammad, H., Deryng, D., Elliott, J., Fujimori, S.,  
651 Hasegawa, T., Heyhoe, E., others, 2014. Climate change effects on agriculture: Economic  
652 responses to biophysical shocks. *Proc. Natl. Acad. Sci.* 111, 3274–3279.

653 Pathak, H., Wassmann, R., 2009. Quantitative evaluation of climatic variability and risks for wheat  
654 yield in India. *Clim. Change* 93, 157–175. <https://doi.org/10.1007/s10584-008-9463-4>

655 Podestá, G., Herrera, N., Veiga, H., Pujol, G., Skansi, M. de los M., Rovere, S., 2009. Towards a  
656 regional drought monitoring and warning system in southern South America: an assessment  
657 of various drought indices for monitoring the 2007-2009 drought in the Argentine Pampas.

658 Pope, V., Stratton, R., 2002. The processes governing horizontal resolution sensitivity in a climate  
659 model. *Clim. Dyn.* 19, 211–236. <https://doi.org/10.1007/s00382-001-0222-8>

660 Pope, V.D., Gallani, M.L., Rowntree, P.R., Stratton, R.A., 2000. The impact of new physical  
661 parameterizations in the Hadley Center coupled model without flux adjustments. *Clim. Dyn.*  
662 17, 61–81.

663 Porter, J.R., Xie, L., Challinor, A.J., Cochrane, K., Howden, S.M., Iqbal, M.M., Lobell, D.B., Travasso,  
664 M.I., 2014. Food security and food production systems. In: *Climate Change 2014: Impacts,*  
665 *Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working*  
666 *Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*  
667 *[Field, C.B., V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee,*  
668 *K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R.*  
669 *Mastrandrea, and L.L.White (eds.)]. Cambridge University Press, Cambridge, United Kingdom*  
670 *and New York, NY, USA.*

671 Power, S., Delage, F., Chung, C., Kociuba, G., Keay, K., 2013. Robust twenty-first-century projections  
672 of El Niño and related precipitation variability. *Nature* 502, 541–545.  
673 <https://doi.org/10.1038/nature12580>

674 Puma, M.J., Bose, S., Chon, S.Y., Cook, B.I., 2015. Assessing the evolving fragility of the global food  
675 system. *Environ. Res. Lett.* 10, 024007. <https://doi.org/10.1088/1748-9326/10/2/024007>

676 Ramteke, R., Gupta, G.K., Singh, D.V., 2015. Growth and Yield Responses of Soybean to Climate  
677 Change. *Agric. Res.* 4, 319–323. <https://doi.org/10.1007/s40003-015-0167-5>

678 Rogelj, J., Knutti, R., 2016. Geosciences after Paris. *Nat. Geosci.* 9, 187–189.



679 Rosenzweig, C., Elliott, J., Deryng, D., Ruane, A.C., Müller, C., Arneth, A., Boote, K.J., Folberth, C.,  
680 Glotter, M., Khabarov, N., others, 2014. Assessing agricultural risks of climate change in the  
681 21st century in a global gridded crop model intercomparison. *Proc. Natl. Acad. Sci.* 111,  
682 3268–3273.

683 Rosenzweig, C., Ruane, A.C., Antle, J., Elliott, J., Ashfaq, M., Chatta, A.A., Ewert, F., Folberth, C.,  
684 Hathie, I., Havlik, P., Hoogenboom, G., Lotze-Campen, H., MacCarthy, D.S., Mason-D’Croz, D.,  
685 Contreras, E.M., Mller, C., Perez-Dominguez, I., Phillips, M., Porter, C., Raymundo, R.M.,  
686 Sands, R.D., Schleussner, C.-F., Valdivia, R.O., Valin, H., Wiebe, K., 2018. Coordinating AgMIP  
687 data and models across global and regional scales for 1.5C and 2.0C assessments. *Philos.*  
688 *Trans. R. Soc. Math. Phys. Eng. Sci.*

689 Ruane, A.C., Antle, J., Elliott, J., Folberth, C., Hoogenboom, G., Croz, D.M.-D., Müller, C., Porter, C.,  
690 Phillips, M.M., Raymundo, R.M., 2018. Biophysical and economic implications for agriculture  
691 of+ 1.5 and+ 2.0 C global warming using AgMIP Coordinated Global and Regional  
692 Assessments. *Clim. Res.* 76, 17–39.

693 Schaffnit-Chatterjee, C., Schneider, S., Peter, M., Mayer, T., 2010. Risk management in agriculture.  
694 *Dtsch. Bank Reseach Sept.*

695 Schlenker, W., Roberts, M.J., 2009. Nonlinear temperature effects indicate severe damages to US  
696 crop yields under climate change. *Proc. Natl. Acad. Sci.* 106, 15594–15598.

697 Schleussner, C.-F., Lissner, T.K., Fischer, E.M., Wohland, J., Perrette, M., Golly, A., Rogelj, J., Childers,  
698 K., Schewe, J., Frieler, K., Mengel, M., Hare, W., Schaeffer, M., 2016a. Differential climate  
699 impacts for policy-relevant limits to global warming: the case of 1.5 &deg;C and 2  
700 &deg;C. *Earth Syst. Dyn.* 7, 327–351. <https://doi.org/10.5194/esd-7-327-2016>

701 Schleussner, C.-F., Rogelj, J., Schaeffer, M., Lissner, T., Licker, R., Fischer, E.M., Knutti, R., Levermann,  
702 A., Frieler, K., Hare, W., 2016b. Science and policy characteristics of the Paris Agreement  
703 temperature goal. *Nat. Clim. Change* 6, 827–835. <https://doi.org/10.1038/nclimate3096>

704 Schmitz, C., van Meijl, H., Kyle, P., Nelson, G.C., Fujimori, S., Gurgel, A., Havlik, P., Heyhoe, E., d’Croz,  
705 D.M., Popp, A., Sands, R., Tabeau, A., van der Mensbrugghe, D., von Lampe, M., Wise, M.,  
706 Blanc, E., Hasegawa, T., Kavallari, A., Valin, H., 2014. Land-use change trajectories up to  
707 2050: insights from a global agro-economic model comparison. *Agric. Econ.* 45, 69–84.  
708 <https://doi.org/10.1111/agec.12090>

709 Seneviratne, S.I., Donat, M.G., Pitman, A.J., Knutti, R., Wilby, R.L., 2016. Allowable CO2 emissions  
710 based on regional and impact-related climate targets. *Nature* 529, 477–483.  
711 <https://doi.org/10.1038/nature16542>

712 Sheffield, J., Goteti, G., Wood, E.F., 2006. Development of a 50-year high-resolution global dataset of  
713 meteorological forcings for land surface modeling. *J. Clim.* 19, 3088–3111.

714 Sklar, M., 1959. Fonctions de répartition à n dimensions et leurs marges. Université Paris 8.

715 Tack, J., Barkley, A., Nalley, L.L., 2015. Effect of warming temperatures on US wheat yields. *Proc.*  
716 *Natl. Acad. Sci.* 112, 6931–6936. <https://doi.org/10.1073/pnas.1415181112>

717 Tao, F., Yokozawa, M., Liu, J., Zhang, Z., 2008. Climate-crop yield relationships at provincial scales in  
718 China and the impacts of recent climate trends. *Clim. Res.* 38, 83–94.

719 Ummenhofer, C.C., Xu, H., Twine, T.E., Girvetz, E.H., McCarthy, H.R., Chhetri, N., Nicholas, K.A., 2015.  
720 How Climate Change Affects Extremes in Maize and Wheat Yield in Two Cropping Regions. *J.*  
721 *Clim.* 28, 4653–4687. <https://doi.org/10.1175/JCLI-D-13-00326.1>

722 UNFCCC, 2015. Adoption of the Paris Agreement. FCCC/ CP/2015/10/Add.1. Paris, France. 1–32.

723 USDA, 2015. Economics, Statistics ad Market Information System.

724 Van Vuuren, D.P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G.C., Kram, T.,  
725 Krey, V., Lamarque, J.-F., others, 2011. The representative concentration pathways: an  
726 overview. *Clim. Change* 109, 5.

727 Verschuuren, J., 2016. The Paris agreement on climate change: Agriculture and food security. *Eur J*  
728 *Risk Reg* 7, 54.

729 Von Braun, J., 2008. The food crisis isn’t over. *Nature* 456, 701–701.

730 Zhang, T., Lin, X., Sassenrath, G.F., 2015. Current irrigation practices in the central United States  
731 reduce drought and extreme heat impacts for maize and soybean, but not for wheat. *Sci.*  
732 *Total Environ.* 508, 331–342.  
733  
734

735

736

737

738

739

740

741

742

743

744

745

746

747

748

749

750

751


752


753 **Tables and Figures**

754 **Table 1.** Pearson correlation coefficient between Princeton re-analysis climatological data and detrended,  
 755 observed historical subnational crop yield data. \*\*\*, \*\* and \* indicate  $p < 0.01$ ,  $p < 0.05$ , and  $p < 0.20$ ,  
 756 respectively. Bold values indicate those properties that have been chosen as climate indicators in this paper.

	<b>Wheat</b>		<b>Maize</b>		<b>Soybean</b>	
	Maximum temperature	Precipitation	Maximum temperature	Precipitation	Maximum temperature	Precipitation
<b>Argentina</b>	<b>-0.493***</b>	-0.140	<b>-0.602***</b>	<b>0.645***</b>	<b>-0.490***</b>	<b>0.675***</b>
<b>Australia</b>	<b>-0.356**</b>	<b>0.825***</b>				
<b>Brazil</b>			-0.023	<b>0.260*</b>	0.041	<b>0.392**</b>
<b>China</b>	0.237	<b>0.147</b>	<b>-0.157</b>	<b>0.335**</b>	-0.032	<b>0.137</b>
<b>India</b>	<b>-0.406***</b>	<b>-0.195*</b>	<b>-0.232*</b>	<b>0.335**</b>	<b>-0.334**</b>	<b>0.533***</b>
<b>USA</b>	-0.035	<b>0.309**</b>	<b>-0.293**</b>	<b>0.420***</b>	<b>-0.208*</b>	<b>0.330**</b>

757





758

759

760

761

762

**Figure 1.** Climate indicators and harvesting periods for the global breadbaskets: Argentina (AR), Australia (AU), Brazil (BR), China (CN), India (IN) and the USA (US). Temperature-based indicators (continuous line) are monthly maximum temperature averaged over the crop growth relevant period. Precipitation-based indicators are cumulative precipitation over selected time periods (dashed line).

763

764

765

766

767

768

769

770

771

772

773

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789



790

791


792

793

a)




b)




794


795

796  
 797  
 798  
 799  
 800  
 801  
 802  
 803  
 804  
 805  
 806  
 807  
 808  
 809  
 810  
 811  
 812  
 813  
 814  
 815  
 816



**Figure 2.** Changes in climate threshold exceedance between historical and 1.5 or 2 °C warming scenarios (in percentage points) using temperature and rainfall based indicators. A) shows the global breadbaskets for wheat, maize and soybean, b) summarizes the risk changes for the two warming scenarios. The error bar indicates the standard error between the 1000 iterations of threshold exceedance using resampled climate data. Using the KS test, all differences between the 1.5°C and 2°C scenarios are significant at a 0.001 significance level.





817

818

819

**Figure 3.** Risks of multiple breadbasket failure under 1.5 and 2°C warming. Error bars reflect the sampling error as well as the copula simulation error which was determined in 1000 iterations.