

Predictability and transferability of local biodiversity environment relationships

Martin Jung

Biodiversity and Natural Resources Program, International Institute for Applied Systems Analysis (IIASA), Laxenburg, Lower Austria, Austria

ABSTRACT

Background: Biodiversity varies in space and time, and often in response to environmental heterogeneity. Indicators in the form of local biodiversity measures—such as species richness or abundance—are common tools to capture this variation. The rise of readily available remote sensing data has enabled the characterization of environmental heterogeneity in a globally robust and replicable manner. Based on the assumption that differences in biodiversity measures are generally related to differences in environmental heterogeneity, these data have enabled projections and extrapolations of biodiversity in space and time. However so far little work has been done on quantitatively evaluating if and how accurately local biodiversity measures can be predicted.

Methods: Here I combine estimates of biodiversity measures from terrestrial local biodiversity surveys with remotely-sensed data on environmental heterogeneity globally. I then determine through a cross-validation framework how accurately local biodiversity measures can be predicted within (“predictability”) and across similar (“transferability”) biodiversity surveys.

Results: I found that prediction errors can be substantial, with error magnitudes varying between different biodiversity measures, taxonomic groups, sampling techniques and types of environmental heterogeneity characterizations. And although errors associated with model predictability were in many cases relatively low, these results question—particular for transferability—our capability to accurately predict and project local biodiversity measures based on environmental heterogeneity. I make the case that future predictions should be evaluated based on their accuracy and inherent uncertainty, and ecological theories be tested against whether we are able to make accurate predictions from local biodiversity data.

Subjects Biodiversity, Conservation Biology, Ecology, Data Science, Spatial and Geographic Information Science

Keywords Spectral-diversity, Biodiversity-productivity, Transferability, Remote-sensing, PREDICTS, Extrapolation, Biodiversity indicators, Prediction uncertainty

INTRODUCTION

Local biodiversity on land is known to vary with environmental heterogeneity (*Hillebrand, 2004; Stein & Kreft, 2015; Holt et al., 2017*), often quantified as difference in availability and variability of resources available to a species. These resources, such as nutrients, water, energy characterize the suitable habitat where the population of a species can persist and commonly include the key availability and diversity of components of ecosystem

Submitted 4 May 2022
Accepted 19 July 2022
Published 23 August 2022

Corresponding author
Martin Jung, jung@iiasa.ac.at

Academic editor
Eric Ward

Additional Information and
Declarations can be found on
page 15

DOI 10.7717/peerj.13872

© Copyright
2022 Jung

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

functioning (Stein & Kreft, 2015; Regos et al., 2022). Several theories have been postulated as possible source of the relationship of environmental heterogeneity with local biodiversity. These include, among others, the widely tested species-energy (Hurlbert, 2004; Evans, Warren & Gaston, 2005; Duncan et al., 2015), the species spectral-heterogeneity (Oldeland et al., 2010; Rocchini et al., 2010) or the species-geodiversity hypotheses (Alahuhta, Toivanen & Hjort, 2020). The assumption is that habitats with greater heterogeneity provide more niches for diverse sets of species and species population expansions. However, despite a number of global meta-analyses on the relationship between environmental heterogeneity and local biodiversity for plant, bird and mammal species (Stein, Gerstner & Kreft, 2014; Duncan et al., 2015), it has rarely been comprehensively investigated how predictable and transferable these relationships are, especially across taxonomic groups and different biodiversity measures more generally.

Predictions made by statistical models are key for our understanding of the living world and for the creation of outputs relevant for conservation management (Miller et al., 2004; Houlahan et al., 2017). Because of the evermore increasing demand for scenarios and spatial maps by policy makers and land managers, biodiversity modellers often need to rely on inter- and extrapolations of model predictions across space and time (Miller et al., 2004). These predictions need to be precise and accurate enough for the context and decisions they are meant to inform (Santini et al., 2021). Thus model predictions should be investigated for their predictability, e.g., a model's ability to accurately predict correlative relationships within the same spatial and/or temporal context by withholding some parts of the data (as in cross-validation procedures), and transferability, e.g., the capacity to produce accurate predictions for conditions dissimilar to those of the data for which a model was trained by withholding data outside the range used for calibration or making use of independently collected data (Petchey et al., 2015; Jung et al., 2017; Yates et al., 2018; Tredennick et al., 2021). And yet, model predictability and transferability is rarely consistently assessed and, when studied in more detail, results rarely look promising.

There is increasing evidence that models using variables of environmental heterogeneity, in the context of this work defined as characterizations of local habitats by vegetation and land-surface conditions, often fail to accurately predict and transfer local biodiversity measures. Studies have found that the predictability of local biodiversity as function of a difference in environmental heterogeneity are highly variable between geographic regions (Phillips, Newbold & Purvis, 2017) and local contexts (Duncan et al., 2015; Jung et al., 2017). Similarly, transferability of model predictions to spatial or temporally distinct regions has long been recognized as key issue for species distribution models (Zurell, Elith & Schröder, 2012; Mesgaran, Cousens & Webber, 2014; Regos et al., 2019) or models using local and regional biodiversity measures (Parmentier et al., 2011; Schmidtlein & Fassnacht, 2017). Despite the development of techniques for assessing the novel parameter space of a model (Zurell, Elith & Schröder, 2012; Meyer & Pebesma, 2021), the limited uptake of modellers to evaluate and present model uncertainty can hinder the application and affect trust in biodiversity model predictions (Rapacciuolo, 2019). Here, one key reason and shortcoming in doing so in macroecological studies are the various and

context-specific ways in which environmental heterogeneity is quantified (Stein & Kreft, 2015), preventing assessments of predictability and transferability.

Recent advances in remote sensing and cloud-processing have enabled the robust quantification of environmental heterogeneity at high spatial and temporal resolution (Gorelick et al., 2017; Randin et al., 2020; Regos et al., 2022). Through repeated satellite observations, measures of environmental heterogeneity, such as differences in photosynthetic activity or spectral variability as proxies for vegetation cover, vegetation condition and structure and overall variability of land surfaces (Rocchini et al., 2010; Radeloff et al., 2019), can be robustly quantified. On their own they can be considered continuous representations of contrasts in land cover and land use (Hansen et al., 2000; Jung, Scharlemann & Rowhani, 2020) and ecosystem functioning (Regos et al., 2022), while also being related to key species population processes (Pettorelli et al., 2005). Although such remotely-sensed measures do not cover all types of environmental heterogeneity (Stein, Gerstner & Kreft, 2014), they can be exogenously quantified in space and time and have often been incorporated in statistical models for the prediction of species distributions (Cord et al., 2013; He et al., 2015; Regos et al., 2022) or to infer differences in local biodiversity measures (Oldeland et al., 2010; Goetz et al., 2014; Rocchini, Hernández-Stefanoni & He, 2015; Jung et al., 2019; Jung, Scharlemann & Rowhani, 2020). Remote sensing measures of environmental heterogeneity can therefore—opposed to study-specific predictors commonly included in ecological meta-analysis—serve as a globally consistent predictor for evaluating biodiversity environment relationships (Duncan et al., 2015). With the availability of new global databases on local biodiversity *in-situ* observations (Hudson et al., 2017), it has become possible to comparably investigate the predictability and transferability of biodiversity environment relationships across taxonomic groups and biodiversity measures.

There are a number of shortcomings in previous analyses on the predictability and transferability of local biodiversity environment relationships. Most studies have (a) focussed on effect sizes among studies (e.g., strength of inference), rather than the predictability and transferability of this relationships (Tredennick et al., 2021), (b) tended to focus mostly on species richness (Stein, Gerstner & Kreft, 2014), thus ignoring other biodiversity measures such as abundance or differences in species assemblage composition, (c) used variables of varying origin to capture effects of changes in environmental heterogeneity on biodiversity (Supp & Ernest, 2014; Shackelford et al., 2017) or have (d) focussed only on regional extents and single taxonomic groups such as birds, butterflies or plants (Kerr, Southwood & Cihlar, 2001; Oldeland et al., 2010; Goetz et al., 2014; Schmidlein & Fassnacht, 2017). Quantitatively addressing these issues is key, if we are to understand in which cases spatial and/or temporal predictions of local biodiversity measures are reliable and accurate.

In this study I investigate the predictability and transferability of model-based predictions on local biodiversity environment relationships. The expectation is that (i) predictability is generally stronger than transferability, (ii) transferability of species-environment relationships affects some biodiversity measures and taxonomic groups more than others, and that (iii) unexplained variation in the tested relationships is




Figure 1 Schematic of the analysis framework showing the distribution of two hypothetical studies and their sites at which a biodiversity measure and environmental predictor has been calculated. (A) Hypothetical studies are coloured in orange and red and the Normalized Difference Vegetation Index (NDVI) is shown as example of a remotely sensed environmental predictor. A simplified procedure for investigating the (B) predictability and (C) transferability of local biodiversity-environment relationships is shown. For (B) ‘testing’ sites within a studies are removed at random, regressions refitted and the within-study prediction error quantified in relation to study properties. In contrast, in (C) regression fits from one study (orange) are used to predict permuted biodiversity estimates in another study (red) that have been removed (beige), with the prediction error quantified in relation to study properties (*i.e.*, scale, sampling length, indicated by different icons).

Full-size DOI: [10.7717/peerj.13872/fig-1](https://doi.org/10.7717/peerj.13872/fig-1)

predominantly linked to differences in study design, *e.g.*, spatial scale and sampling duration. To test this, I combine local biodiversity data from globally distributed surveys with remotely-sensed environmental predictors quantifying photosynthetic activity (Evans, Warren & Gaston, 2005; Stein, Gerstner & Kreft, 2014; Duncan *et al.*, 2015) and spectral variability (Rocchini *et al.*, 2010); predictors that represent a continuous characterization of the availability of resources and land surface modifications. Using variations of generalized linear and additive models, I assess the predictability, quantified as overall and within-study reduction in prediction error, and transferability, quantified as reduction in prediction error between different studies of comparable design but identical taxonomic groups (Fig. 1). The aim of this work is thus to comparatively assess the strength and generality of local biodiversity-environment relationships at a global scale, which hopefully stimulates a debate on whether predicted local biodiversity measures, such as total site-based abundance or richness, can accurately be predicted or transferred to unsampled regions.

MATERIALS AND METHODS

Biodiversity data preparation

For data on biodiversity I took species assemblage data from the global Projecting Responses of Ecological Diversity In Changing Terrestrial Systems (PREDICTS) database (Hudson *et al.*, 2017), which contains records of species occurrence and abundance at spatial-explicit sites ‘sites’ as reported in published ‘studies’. PREDICTS includes only studies which differ in ‘land-use’ and/or ‘land-use intensity’ and have spatial and temporal information associated with them, *e.g.*, sampling extent and date of sampling (Hudson *et al.*, 2014). Studies in the PREDICTS database vary widely in study properties, notably in

taxonomic coverage (studies contain data on terrestrial species of invertebrates, plants, birds, mammals, reptiles and amphibians), spatial grain (0.05–39,150 m, median = 70 m \pm 97 MAD), sampling start (1984–2013), sampling effort (>0–4,382 days, median = 91 days) and methodology (flight traps, transects,...). Owing to these differences, a hierarchical modelling framework is usually necessary when analysing biodiversity estimates from databases such as PREDICTS ([Purvis et al., 2018](#)).

For each study j and site i in the PREDICTS database, I calculated four different site-based measures of local biodiversity: total Species richness (S_i), total log-transformed abundance ($\log_{10}A_i$), the arcsine square root transformed probability of interspecific encounter as measure of assemblage evenness ($\sin^{-1}\sqrt{PIE_i}$) and the logit transformed pairwise Sørensen similarity index as measure of difference in assemblage composition ($\text{logit } SIM_{i-i_n}$). Similar to previous studies I assumed that, in the few cases where within-study study effort differs among sites, the abundance of species individuals increases linearly with sampling effort ([Newbold et al., 2015](#)). In cases where the sampling extent of a site is missing in the PREDICTS database, I approximated the mean sampling extent using a heuristic that fills missing estimates with the average used within studies of the same sampling method and/or taxonomic group. Earlier work has shown that this approximation can accurately fill missing sampling extents ([Jung et al., 2019](#)). Lastly, I created, based on the taxonomic group and sampling method attributed to a study in the PREDICTS database, a new factor variable that groups studies of comparable method, unit and broad taxonomic grouping ([Table S1](#)), such as for instance studies involving bird individuals that were counted using point counts. I realize that not all differences in sampling techniques can be attributed to this new contrast between sites and therefore post-hoc analyse the contribution of differing sampling methods in explaining the cross-validated model error (see statistical analysis). In total I used data from 564 different studies and 25,849 sites, with the median number of sites per study being 18 (IQR = 35, see also [Fig. S6](#)).

Environmental predictors

In this work I exclusively used remotely-sensed environmental predictors, namely photosynthetic activity and spectral variability, which are (1) available at adequate spatial resolution, (2) consistently quantified at global extent in comparable units, (3) temporally explicit, often differing between years, (4) correlate with differences in local biodiversity ([Duncan et al., 2015](#); [Jung et al., 2019](#)) and land use ([Mueller et al., 2014](#); [Yin et al., 2014](#)). These predictors can be considered generic proxies of resources available to species ([Pettorelli et al., 2005](#)) as well as characterizing differences in land surface conditions on a continuous scale ([Rocchini et al., 2010](#); [Jung et al., 2019](#); [Randin et al., 2020](#)). It should be noted that the aim of this work is not to identify best possible predictors of local biodiversity, but rather to evaluate most commonly used ones for their predictability and transferability. Other environmental predictors quantifiable from remote sensing exist can provide a better characterization of processes related to the water cycle (e.g., NDWI ([Gao, 1996](#)) or energy balance such as LST ([Albright et al., 2011](#))). A more detailed

discussion on the potential of remote sensing in predicting local biodiversity is provided in the discussion.

For each site in the PREDICTS databases, I calculated two different remotely sensed predictors that reflect environmental heterogeneity. First, 16-day time series of atmospherically corrected spectral observations (MCD43A v006, (Schaaf *et al.*, 2002)) from the Moderate Resolution Imaging Spectroradiometer (MODIS) sensor on board the Terra and Aqua satellites. These data products are available at a 500 m spatial resolution globally and were downloaded and extracted for each PREDICTS site from Google Earth Engine (Gorelick *et al.*, 2017). Time series of remotely sensed spectral observations often have data gaps caused by clouds or sensor errors. To reduce the number of data gaps, I first aggregated (arithmetic mean) the obtained time series to monthly estimates for each spectral observation (band 1 to 7). The overall proportion of missing data in the aggregated time series was low (mean: $5.9\% \pm 10.5$ SD), nevertheless I subjected the aggregated time series to a missing value imputation using a Kalman smoother on the whole time series (Hyndman & Khandakar, 2008) as implemented in the 'imputeTS' R package (Moritz & Bartz-Beielstein, 2017). Whenever the imputation did not converge, a linear interpolation was used to impute missing observations among years. Only data gaps smaller than 5 months were filled in that manner and sites with six or more missing months were excluded from subsequent analyses. From the full time series, I then selected for each site the first year (12 months) of data preceding biodiversity sampling as representation of environmental heterogeneity (Jung *et al.*, 2019).

Second, I calculated from the remaining time series of spectral observations, as proxy of overall photosynthetic activity, the arithmetic mean of the two-band Enhanced Vegetation Index (EVI, Jiang *et al.*, 2008). Photosynthetic activity approximates the condition, structure and availability of plant biomass. Variations in photosynthetic activity have previously been shown to reflect continuous gradients in land cover (Huete *et al.*, 2002; Radeloff *et al.*, 2019) and directly influence local biodiversity measures and life history (Pettorelli *et al.*, 2005; He, Zhang & Zhang, 2009; Oldeland *et al.*, 2010; Jung *et al.*, 2019; Jung, Rowhani & Scharlemann, 2019). Furthermore, I also calculated a measure of overall spectral variability from the satellite sensor data (Rocchini *et al.*, 2010; Rocchini, Hernández-Stefanoni & He, 2015; Randin *et al.*, 2020). Spectral variability is expected to give a more nuanced view on land surface conditions than any single vegetation index, given that it utilizes not two but all spectral bands of the satellite (Rocchini *et al.*, 2010) and thus being able to capture variation among most of spectral scales covered by the MODIS sensor. To capture spectral variability, I first calculated a principal component analysis of all spectral observations (bands 1–7) and then calculated from the first two axes, which on average explained $93\% \pm 5.92$ SD of all variation, the centroid of the resulting bivariate scatter plot. Spectral variability per site was then summarized as the mean Euclidean distance to this centroid. Both environmental predictors, photosynthetic activity and spectral variability are only weakly correlated (Pearson's $r = -0.21$, Fig. S1). In total, 21,821 sites had suitable remote sensing data for subsequent analyses, with the remainder (4,028 sites) being sampled either too long ago for sufficient remote sensing coverage from MODIS (2,000 onwards) or having too many data gaps.

Statistical analysis

In the context of this work, ‘predictability’ is defined as the ability to accurately infer a biodiversity measure y_{ij} based on the environmental covariates x_{ij} among the sites i of a PREDICTS study j (Fig. 1B), and ‘transferability’ as the ability to accurately predict y_i based on the environmental covariates x_i across studies of the same sampling methodology and taxonomic group (Fig. 1C).

In both predictability and transferability variants prediction accuracy is assessed by calculating for each study the symmetric mean absolute percentage error

$$\left(\text{sMAPE}_j = \frac{100}{n} \sum_{i=1}^I \left| \frac{Y_{\text{predicted}} - Y_{\text{observed}}}{(|Y_{\text{observed}}| + |Y_{\text{predicted}}|)} \right| \right)$$

between the observed biodiversity measures (y_{observed}) and the ones predicted by the model ($y_{\text{predicted}}$) for a given site i .

The sMAPE quantifies the percentage error in a model prediction and is bounded between 0% and 100%. Alternative metrics to quantify prediction precision and accuracy exists, however in this case the sMAPE is preferable for PREDICTS style data owing to its simplicity and inter-comparability between studies that use biodiversity measures of different units and value ranges.

I constructed separate models for each study j and biodiversity measure y in site i , by assuming that $y_i = \alpha_i + \beta_i x_i + \epsilon$, where α is the study specific intercept, β a slope coefficient, x the environmental predictor and ϵ an error term. Models of S_i were assumed to have Poisson distributed errors and a log-link function ($\log y$), while models of A_i , PIE_i and SIM_{i-i_n} were assumed to have Gaussian distributed errors. Pairwise similarities in species composition (Sorensen Index) were related to differences in environmental predictors x in addition to pairwise distance between sites, calculated as $\log_{10}(x + 0.05 \text{ km})$ from great circle distances between sites. Here I calculated pairwise absolute difference in mean photosynthetic activity or between spectral centroids of each site (see environmental predictors). For each constructed full model I furthermore calculate an R^2 measure as indication of overall variance explained.

To evaluate the predictability and transferability of local biodiversity environment relationships, I constructed in total ten permutation sets, in each of which sites were split into testing (33%) and training (66%) datasets. For evaluating predictability, I removed one third of sites (33%) at random (Fig. 1B), but weighted them by the mean distance to the study centroid, therefore placing extra weight on sites that are less likely to be in close proximity (Roberts *et al.*, 2017). For transferability, instead of individual sites within the same study, I instead sampled and removed 33% for each set sites at random from a different study of comparable methodology in the PREDICTS database (Fig. 1C, methods above). Thus predictability permutations contained samples of training and testing sites for the same study, while transferability permutations contained training sites from one study and withheld testing sites from another comparable one. For studies with sufficient number of sites (>4) I furthermore tested whether considering non-linear relationships (estimated using generalized additive models) resulted in prediction with lower sMAPE, however found that linear models throughout resulted in predictions with smaller errors. However across all ten permutation sets, I iteratively weighted (0–1) this sampling by

whether a given study has been sampled before, therefore ensuring that each study is part of both testing and training dataset at least once.

For each respective permutation set, predictability and transferability was then evaluated by using the remaining training data to estimate the regression specified above for each study or group of comparable methodology. I excluded combinations of taxonomic groups, sampling method and sampling unit for which fewer than two studies were available. In total 77.3% of all studies had a matching study of comparable methodology and unit for the same taxonomic group. A table with all recategorized combinations (43) can be found in the [Supplemental Materials \(Table S1\)](#). Using the fitted models I predicted y for the excluded 'hold-out' 33% sites and then calculated the average sMAPE for each study in the permutation sets.

Lastly, I explored possible correlates of why sMAPE for some studies is larger than for others for each of the four considered biodiversity measures. I considered a series of variables commonly related to differences in sampling design, species and individual detectability and errors in remotely-sensed environmental predictors. Specifically, I calculated for each study in the permutation sets, the median sampling extent (m) as measure of sample grain, the median sampling duration (days) of the study, the number of sites with a study as measure of effort for the whole study, the average number of samples across sites as effort for area-based sampling effort or the average time sampled (hours) for time-based sampling effort, average accessibility to sites in the study (distance to nearest city in meters) from [Weiss et al. \(2018\)](#), and finally factors related to possible errors in remotely-sensed environmental variables, including the amount of missing data (before gap filling) and the average topographic ruggedness per study using data from [Amatulli et al. \(2018\)](#). To make comparisons across these different units and scales, I standardized all variables before model fitting by subtracting the mean and dividing by one standard deviation.

I fitted linear models allowing partial pooling among studies j ([Harrison et al., 2018](#)) by adding a random intercept α_k in addition to the overall intercept, e.g., $SMAP E_j = \alpha + \alpha_k + \beta_j x_j + \epsilon$. These kind of models can borrow strength among studies by shrinking individual estimates towards an overall population-wide average ([Purvis et al., 2018](#); [Harrison et al., 2018](#)). As random intercept k I used the methodology specific grouping (see methods and [Table S1](#)), thus pooling possible correlates among studies of similar methodology. I fitted all possible combinations between the above mentioned variables, including an interaction between sampling extent and sampling effort, finally constructing an average ensemble model of the 5% best performing models. Models were fitted in `lme4` ([Bates et al., 2015](#)) using the 'MuMIn' package in R for model averaging ([Bartoń, 2015](#)).

RESULTS

The explanatory power of environmental predictors—photosynthetic activity and spectral variability—in explaining differences in biodiversity varied across biodiversity measures and individual studies. Models fitted with photosynthetic activity explained on average slightly more variance than models fitted with spectral variability, the former having an average R^2 of 0.21 (± 0.285 SD) compared to an average R^2 of 0.19 (± 0.284 SD) in the latter.




Figure 2 Explained variance (R^2) calculated from models fitted between different biodiversity measures and either photosynthetic activity or spectral variability. Each point is an individual study in the PREDICTS database with point size indicating the number of sites per study and the colour being a visual indication of density in the plot. A map of the average R^2 per study and biodiversity measure can be found in Fig. S2. [Full-size !\[\]\(b345a1c4255362eec3746050dd71ccac_img.jpg\) DOI: 10.7717/peerj.13872/fig-2](https://doi.org/10.7717/peerj.13872/fig-2)

There was considerable variation of R^2 values across studies and biodiversity measures (Fig. 2), with species richness on average being best explained by photosynthetic activity ($R^2 = 0.246 \pm 0.311$ SD) or spectral variability ($R^2 = 0.22 \pm 0.306$ SD). Notably, correlations with species abundance were particularly low, with the R^2 being close to 0 ($R^2 < 0.001$) for more than a quarter of all studies (Fig. 2). Meanwhile the difference in explained variance between models using photosynthetic activity compared to spectral variability was lowest for differences in assemble composition (Pearson's $R = 0.922$). There were no obvious spatial (Fig. S2) or directional patterns (Fig. S3) in the average explained variance, although some studies notably had high explanatory power regardless of the considered biodiversity measure (Fig. S2).

When applying local biodiversity models to known ('Predictability') or different ('Transferability') contexts, the main issue is how accurately such models can predict local biodiversity measures in unknown situations based on the covariates of interest (Fig. 3). Regardless of whether remotely-sensed photosynthetic activity or spectral variability was used as covariate, linear models were reasonably accurate for known contexts in inferring species richness (sMAPE of 19.1%), abundance (11.8%) and evenness (10.3%), but less so when inferring differences in species assemblages (49.3%). Errors in predicting local biodiversity to different contexts were expectedly larger (Fig. 3), whereas particular species richness could be extrapolated relatively poorly (relative error 43.3%) similarly to differences in species assemblages (67.9%), compared to abundance (25.4%) or evenness (14.3%). Notably, when local biodiversity models are used to extrapolate richness to different contexts, the sMAPE was larger than 50% in 35% of all studies, compared to 8.1% and 4.7% for abundance and evenness (Fig. 3).




Figure 3 Distribution of the symmetric mean absolute percentage error (sMAPE) of biodiversity measures calculated from models using photosynthetic activity or spectral variability. Larger values (range 0 to 100) indicate a larger prediction error. Colours differentiate between models that evaluate Predictability and Transferability (see Methods). Point error ranges show the arithmetic mean and standard deviation of the sMAPE. [Full-size !\[\]\(fcc3264021d438d9732560e78099f674_img.jpg\) DOI: 10.7717/peerj.13872/fig-3](https://doi.org/10.7717/peerj.13872/fig-3)

There were also considerable differences in prediction error, as quantified by the sMAPE, among taxonomic groups. Across taxonomic groups and biodiversity measures the sMAPE was larger when predictions were extrapolated to novel contexts compared to predictability, particularly so for reptiles (Δ sMAPE = 21.3%) and mammals (Δ sMAPE = 20.8%), with the greatest difference being for reptile species richness (Δ sMAPE = 33%) and abundance (Δ sMAPE = 28%). The transferability of fungi (sMAPE = 7.5%), and bird (sMAPE = 9.1%) assemblage evenness was overall the lowest, while predictability was best for evenness and abundance of fungi (sMAPE = 5.11%) and plants (sMAPE = 9.65%). Fungi and Plants had across biodiversity measures the lowest sMAPE in predictability and transferability (Fig. 4). Overall, assemblage composition of vertebrates was the most poorly predicted with sMAPE estimates well over 50% throughout (Fig. 4).

I also explored across studies which factors helped explain differences in prediction error, as quantified by the sMAPE (Fig. 5). Across biodiversity measures, having a greater number of samples per site most effectively reduced the sMAPE ($\Delta\beta = -3.14$) for transferability, and so did sample duration but to a lesser degree ($\Delta\beta = -0.98$). Meanwhile a greater number of sites per study on average increased the sMAPE ($\Delta\beta = 2.23$). Patterns of comparison results were broadly similar between transferability (Fig. 5) and predictability (Fig. S5), although notably a study being more accessible resulted in an average larger reduction in the sMAPE ($\Delta\beta = -1.02$) for predictability (Fig. S5). Overall variance




Figure 4 Average error (sMAPE) across models for predictability and transferability. Errors were averaged (lines indicating standard deviation) across models with different biodiversity measures (shapes) and taxonomic group (colours). Shown only for models using photosynthetic activity as predictor as spectral variability results were broadly comparable in overall patterns (Fig. S4).

Full-size [DOI: 10.7717/peerj.13872/fig-4](https://doi.org/10.7717/peerj.13872/fig-4)

explained by these factors in the average model was relatively low ($R_{marginal}^2 = 0.08$, $R_{conditional}^2 = 0.14$).

DISCUSSION

In this work I make use of a large database of local biodiversity survey records and remote sensing data to evaluate the predictability and transferability of biodiversity-environment relationships, *e.g.*, the ability of models to infer local biodiversity measures in known and novel contexts. Particular emphasis is placed on differences among biodiversity measures, taxonomic groups and sampling circumstances. I found that the explanatory power of biodiversity-environment was relatively low for most studies (Fig. 2). This aligns with a previous meta-analysis that found that relationships between biodiversity measures and photosynthetic activity cannot always be established (Duncan *et al.*, 2015). I also discovered that prediction errors are on average lowest for evenness and abundance, and, maybe unsurprisingly, generally larger when models predictions are transferred to novel contexts (Fig. 3). Biodiversity measures of sessile organisms were on average more precisely predicted (Fig. 4), although not by much with predictions errors generally larger than 25% compared to observed values, particularly so for differences in species




Figure 5 Averaged and standardized model coefficients of variables that best explain differences in sMAPE. Standardized coefficients smaller than zero indicate that increases in a given variable reduce study-specific prediction errors, while coefficients greater than zero increase the error. Shapes distinguish different biodiversity measures (as in Fig. 3). Standardized coefficients shown for transferability permutations only as predictability results follow similar patterns (Fig. S5).

Full-size DOI: [10.7717/peerj.13872/fig-5](https://doi.org/10.7717/peerj.13872/fig-5)

assemblage composition. Overall these results shed some doubts on the predictability and transferability of biodiversity measures, although they have to be interpreted in the context of the individual studies (Fig. 5) and ultimately in what is an acceptable accuracy to achieve with such predictions.

Indeed, it is not formerly defined what makes a prediction better or worse based on quantitative measures such as the cross-validated error metrics used in this study. According to *Yates et al. (2018)* ‘transferability’ is broadly defined as the capacity of a model to produce predictions for a new set of predictor values that differ from those on which the model was trained. Similarly predictability can be understood as the capacity of a model to infer held-out observations (Fig. 1). In this context, a good precision could be understood as a model that demonstrates transferability errors smaller or comparable to errors inherent in model inferences or that don’t exceed an a priori set threshold. I found that the predictability of local biodiversity measures was overall reasonable good with errors being smaller than 25% in most cases (Fig. 3), although particularly differences in assemblage composition were poorly predicted. This might indicate that photosynthetic activity and spectral variability are useful predictors for quantifying differences in local

biodiversity measures, although the variance explained varied considerably across studies (Fig. 2). In contrast I found that errors associated with transferability of biodiversity measures can be considerable, exceeding 50% relative to the original measure for species richness and differences in assemblage composition in many studies (Fig. 3). This is especially relevant, since a number of studies spatially extrapolated local biodiversity estimates, e.g., species richness or abundance, to unsampled areas based on environmental predictors (König, Weigelt & Kreft, 2017; Phillips et al., 2019; van den Hoogen et al., 2019). These approaches assume that local biodiversity-environment relationships are transferable to new, unsampled environments and the results by this work indicate that this often entails considerable errors. Ideally models are evaluated on their ability to accurately reproduce their data in novel contexts (Jung et al., 2017), quantify the uncertainty in doing so, or alternatively limit predictions to areas within the models applicability (Mesgaran, Cousens & Webber, 2014; Meyer & Pebesma, 2021), and the results of this work further highlight that prediction errors should be reported.

Biodiversity measures for certain taxonomic groups might be easier to predict than others owing to the dynamics, drivers and mechanisms underlying them (Magurran, 2004). Indeed previous studies have found species abundance to be stronger correlated with photosynthetic activity than other measures (Oldeland et al., 2010; Duncan et al., 2015). Similarly, I found that abundance-based biodiversity measures—e.g., abundance and evenness—had overall lowest precision errors (Fig. 3). A potential mechanism could be that a greater photosynthetic activity or spectral variability is indicative of resources available to species populations, facilitating population growth (Hurlbert, 2004; Pettoelli et al., 2006). While species richness had the largest average explained variance compared to other biodiversity measures, it performed considerably poorer when evaluated in predictions (Fig. 3). Possibly, the processes underlying patterns of local species richness, such as colonization and extinction, might cause simple predictions to fail (Chase, 2003), unless the spatial-temporal dynamics of environmental predictors are taken into account (Fernández, Román & Delibes, 2016). Similarly, the fact that both predictability and transferability errors were on average lowest for more sessile organisms such as Fungi and Plants (Fig. 4), likely indicates that similar important processes mediate biodiversity-environment relationships. Overall this study highlights the benefit of comparing relationships across a range of studies and biodiversity measures (Stein, Gerstner & Kreft, 2014; Duncan et al., 2015), revealing that biodiversity-environment relationships are not universally strong.

Investigating as to what factors best explain prediction errors can help to improve future monitoring and modelling efforts. Among the most important factors that resulted in overall smaller prediction errors was the average number of samples per sites (Fig. 5), which can be considered a simplified metric of sampling completeness. Given that errors were smaller for sites with many samples, it could be that many species communities in the PREDICTS database have not been comprehensively sampled, if one assumes that biodiversity-environment relationships are strongest in equilibrium. There are ways to

account for detectability and observation biases (Royle, Nichols & Kéry, 2005), which however was not feasible for the studies in the PREDICTS database given the heterogeneity of sampling information. Thus better standards for sampling techniques and monitoring are advisable to enable better comparability (Montgomery *et al.*, 2021).

Interestingly, and in contrast to previous studies (Chase & Knight, 2013), differences in sample grain, *e.g.*, the linear scale of sampling, did not help to explain why biodiversity measures could be better predicted in some studies. A likely explanation is that the contrasts between sampling extents are relatively small (most studies in the PREDICTS database were sampled at scales between ~1 and 4,000 m). Scale-dependent effects might only become apparent at spatial scales that go beyond the local scale. A spatial mismatch at the lower end, *e.g.*, that the grain of the used MODIS data is too coarse to be matched to the extent of sampling in PREDICTS studies, could be another explanation, however previous studies that used very-high resolution satellite imagery (<10 m) did not find much more accurate predictions than presented here (Dalmayne *et al.*, 2013; Hofmann *et al.*, 2017). Other, non-explored factors could further explain differences in prediction error, such as for instance preceding changes in environmental predictors (Jung *et al.*, 2019; Jung, Rowhani & Scharlemann, 2019) or a better accounting of differences in species traits (Duncan *et al.*, 2015; Regos *et al.*, 2019). Future efforts could evaluate if inter- and intra-specific variability of species traits can be more precisely linked to differences in environmental heterogeneity.

In this work I used photosynthetic activity and spectral variability as measures of environmental heterogeneity, acknowledging that other characterizations of environmental heterogeneity (*e.g.*, soil, micro-climate) could be more important (Stein & Kreft, 2015). The rationale behind focusing on photosynthetic activity and spectral variability in this work was the assumption that they can serve as general broad predictive measures across taxa and regions. There are, however, also a number of other remote-sensing measures, such as for example land-surface temperature or remotely-sensed moisture (Albright *et al.*, 2011; Regos *et al.*, 2022), as well as different summary statistics of such measures (maximum, minimum, variation), which for some local contexts might capture important dynamics and lead to an improved prediction with lower errors. The finding that prediction errors were lowest for plants and fungi could be related to the fact that photosynthetic activity is more closely related to the abundance of these taxa, than for other taxonomic groups, where only indirect correlations (resources for herbivores, differences in land cover) could be the most likely explanation and local factors might have a deciding influence on explaining these differences (Jung *et al.*, 2017). Yet, focussing solely on remotely-sensed variables ensures global consistency and is frequently used to predict local biodiversity measures (Dalmayne *et al.*, 2013; Hobi *et al.*, 2017; Hofmann *et al.*, 2017; Randin *et al.*, 2020).

Another key limitation is that environmental heterogeneity is not necessarily related to differences in land use and land-use intensity, for which the PREDICTS database was explicitly designed (Purvis *et al.*, 2018). Indeed, it could be that the potential of remotely sensed environmental heterogeneity in predicting local biodiversity measures has been exaggerated, and better and more direct characterizations of land use and its management

from remote sensing have to be developed. Previous studies using PREDICTS data found that even using the local land-use categorizations explained few differences in biodiversity measures, with most variance being ‘soaked up’ by study-specific differences (*Newbold et al., 2015, 2018; De Palma et al., 2015*). Further, given the complexities of local species community assembly (*Chase, 2003; Leibold et al., 2004*), any claim of a direct prediction of ‘biodiversity’ through remotely-sensed proxies should thus be interpreted with caution and only under consideration of prediction uncertainties. Remote sensing measures are at best able to capture changes in habitat extent or condition; and those changes do not necessarily correlate strongly with changes in biodiversity measures. Future work should ideally focus on the principal mechanisms of species community assembly, their practical incorporation into models and how remote sensing can assist in capturing relevant predictors.

CONCLUSIONS

The findings presented in this study have particular implications for spatial projections of local biodiversity-environment relationships. Ecological models can and should be used for predictions (*Houlahan et al., 2017; Tredennick et al., 2021*); however, caveats and limitations should be better identified, communicated and hopefully build upon. We need to create models that enable biodiversity-environment relationships to be more predictable across scales and novel contexts, especially when applied to conservation contexts (*Santini et al., 2021*). Remote sensing data can be used for global consistent characterizations of environmental heterogeneity, but given the considerable drops in precision for transferability, this work recommends that prediction errors in projecting local biodiversity measures are clearly communicated and quantified. To improve future biodiversity predictions I further propose that models (a) should be evaluated comprehensively based on their ability to create accurate predictions, (b) account better for underlying hierarchies and sampling effects, (c) ensure that environmental predictors are quantified in a globally replicable and transparent way. Quantitative correlative models might not be the most precise in many situations, but that does not invalidate their use if shortcomings are appropriately communicated.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

The author received no funding for this work.

Competing Interests

The author declares that there are no competing interests.

Author Contributions

- Martin Jung conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

The raw biodiversity data is available at the Natural History Museum:

Lawrence Hudson; Tim Newbold; Sara Contu; Samantha L. L. Hill et al. (2016).

The 2016 release of the PREDICTS database [Data set]. Natural History Museum. <https://doi.org/10.5519/0066354>.

All intermediate analysis results and code are available at GitHub:

<https://github.com/Martin-Jung/TransferabilityPREDICTS>.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.13872#supplemental-information>.

REFERENCES

- Alahuhta J, Toivanen M, Hjort J. 2020.** Geodiversity-biodiversity relationship needs more empirical evidence. *Nature Ecology & Evolution* **4**(1):2–3 DOI [10.1038/s41559-019-1051-7](https://doi.org/10.1038/s41559-019-1051-7).
- Albright TP, Pidgeon AM, Rittenhouse CD, Clayton MK, Flather CH, Culbert PD, Radeloff VC. 2011.** Heat waves measured with MODIS land surface temperature data predict changes in avian community structure. *Remote Sensing of Environment* **115**(1):245–254 DOI [10.1016/j.rse.2010.08.024](https://doi.org/10.1016/j.rse.2010.08.024).
- Amatulli G, Domisch S, Tuanmu M-N, Parmentier B, Ranipeta A, Malczyk J, Jetz W. 2018.** A suite of global, cross-scale topographic variables for environmental and biodiversity modeling. *Scientific Data* **5**(1):180040 DOI [10.1038/sdata.2018.40](https://doi.org/10.1038/sdata.2018.40).
- Bartón K. 2015.** MuMIn: multi-model inference. Available at https://r-forge.r-project.org/R/?group_id=346.
- Bates DM, Maechler M, Bolker B, Walker S. 2015.** lme4: linear mixed-effects models using Eigen and Eigenpack. *Journal of Statistical Software* **67**(1):1–48 DOI [10.18637/jss.v067.i01](https://doi.org/10.18637/jss.v067.i01).
- Chase JM. 2003.** Community assembly: when should history matter? *Oecologia* **136**(4):489–498 DOI [10.1007/s00442-003-1311-7](https://doi.org/10.1007/s00442-003-1311-7).
- Chase JM, Knight TM. 2013.** Scale-dependent effect sizes of ecological drivers on biodiversity: why standardised sampling is not enough. *Ecology Letters* **16**:17–26 DOI [10.1111/ele.12112](https://doi.org/10.1111/ele.12112).
- Cord AF, Meentemeyer RK, Leitão PJ, Václavík T. 2013.** Modelling species distributions with remote sensing data: bridging disciplinary perspectives. *Journal of Biogeography* **40**(12):2226–2227 DOI [10.1111/jbi.12199](https://doi.org/10.1111/jbi.12199).
- Dalmayne J, Möckel T, Prentice HC, Schmid BC, Hall K. 2013.** Assessment of fine-scale plant species beta diversity using WorldView-2 satellite spectral dissimilarity. *Ecological Informatics* **18**(243):1–9 DOI [10.1016/j.ecoinf.2013.05.004](https://doi.org/10.1016/j.ecoinf.2013.05.004).
- De Palma A, Kuhlmann M, Roberts SPM, Potts SG, Börger L, Hudson LN, Lysenko I, Newbold T, Purvis A. 2015.** Ecological traits affect the sensitivity of bees to land-use pressures in European agricultural landscapes. *Journal of Applied Ecology* **52**(6):1567–1577 DOI [10.1111/1365-2664.12524](https://doi.org/10.1111/1365-2664.12524).
- Duncan C, Chauvenet ALM, Brown ME, Pettorelli N. 2015.** Energy availability, spatio-temporal variability and implications for animal ecology. *Diversity and Distributions* **21**(3):290–301 DOI [10.1111/ddi.12270](https://doi.org/10.1111/ddi.12270).

- Evans KL, Warren PH, Gaston KJ. 2005. Species-energy relationships at the macroecological scale: a review of the mechanisms. *Biological Reviews* **80**(1):1–25
DOI [10.1017/S1464793104006517](https://doi.org/10.1017/S1464793104006517).
- Fernández N, Román J, Delibes M. 2016. Variability in primary productivity determines metapopulation dynamics. *Proceedings of the Royal Society B: Biological Sciences* **283**(1828):20152998 DOI [10.1098/rspb.2015.2998](https://doi.org/10.1098/rspb.2015.2998).
- Gao B. 1996. NDWI—a normalized difference water index for remote sensing of vegetation liquid water from space. *Remote Sensing of Environment* **58**(3):257–266
DOI [10.1016/S0034-4257\(96\)00067-3](https://doi.org/10.1016/S0034-4257(96)00067-3).
- Goetz SJ, Sun M, Zolkos S, Hansen A, Dubayah R. 2014. The relative importance of climate and vegetation properties on patterns of North American breeding bird species richness. *Environmental Research Letters* **9**(3):34013 DOI [10.1088/1748-9326/9/3/034013](https://doi.org/10.1088/1748-9326/9/3/034013).
- Gorelick N, Hancher M, Dixon M, Ilyushchenko S, Thau D, Moore R. 2017. Google Earth Engine: planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment* **202**(3):18–27 DOI [10.1016/j.rse.2017.06.031](https://doi.org/10.1016/j.rse.2017.06.031).
- Hansen MC, Defries RS, Townshend JRG, Sohlberg R. 2000. Global land cover classification at 1 km spatial resolution using a classification tree approach. *International Journal of Remote Sensing* **21**(6–7):1331–1364 DOI [10.1080/014311600210209](https://doi.org/10.1080/014311600210209).
- Harrison XA, Donaldson L, Correa-Cano ME, Evans J, Fisher DN, Goodwin CED, Robinson BS, Hodgson DJ, Inger R. 2018. A brief introduction to mixed effects modelling and multi-model inference in ecology. *PeerJ* **6**(1):e4794 DOI [10.7717/peerj.4794](https://doi.org/10.7717/peerj.4794).
- He KS, Bradley BA, Cord AF, Rocchini D, Tuanmu M-N, Schmidtlein S, Turner W, Wegmann M, Pettorelli N. 2015. Will remote sensing shape the next generation of species distribution models? *Remote Sensing in Ecology and Conservation* **1**(1):4–18 DOI [10.1002/rse2.7](https://doi.org/10.1002/rse2.7).
- He KS, Zhang J, Zhang Q. 2009. Linking variability in species composition and MODIS NDVI based on beta diversity measurements. *Acta Oecologica* **35**(1):14–21
DOI [10.1016/j.actao.2008.07.006](https://doi.org/10.1016/j.actao.2008.07.006).
- Hillebrand H. 2004. On the generality of the latitudinal diversity gradient. *The American Naturalist* **163**(2):192–211 DOI [10.1086/381004](https://doi.org/10.1086/381004).
- Hobi ML, Dubinin M, Graham CH, Coops NC, Clayton MK, Pidgeon AM, Radeloff VC. 2017. A comparison of Dynamic Habitat Indices derived from different MODIS products as predictors of avian species richness. *Remote Sensing of Environment* **195**:142–152
DOI [10.1016/j.rse.2017.04.018](https://doi.org/10.1016/j.rse.2017.04.018).
- Hofmann S, Everaars J, Schweiger O, Frenzel M, Bannehr L, Cord AF. 2017. Modelling patterns of pollinator species richness and diversity using satellite image texture. *PLOS ONE* **12**(10):e0185591 DOI [10.1371/journal.pone.0185591](https://doi.org/10.1371/journal.pone.0185591).
- Holt BG, Costa GC, Penone C, Lessard J-P, Brooks TM, Davidson AD, Blair Hedges S, Radeloff VC, Rahbek C, Rondinini C, Graham CH. 2017. Environmental variation is a major predictor of global trait turnover in mammals. *Journal of Biogeography* **45**(1):1–13
DOI [10.1111/jbi.13091](https://doi.org/10.1111/jbi.13091).
- Houlahan JE, McKinney ST, Anderson TM, McGill BJ. 2017. The priority of prediction in ecological understanding. *Oikos* **126**(1):1–7 DOI [10.1111/oik.03726](https://doi.org/10.1111/oik.03726).
- Hudson LN, Newbold T, Contu S, Hill SLL, Lysenko I, De Palma A, Phillips HRP, Alhuseini TI, Bedford FE, Bennett DJ, Booth H, Burton VJ, Chng CWT, Choimes A, Correia DLP, Day J, Echeverría-Londoño S, Emerson SR, Gao D, Garon M, Harrison MLK, Ingram DJ, Jung M, Kemp V, Kirkpatrick L, Martin CD, Pan Y, Pask-Hale GD, Pynegar EL, Robinson AN, Sanchez-Ortiz K, Senior RA, Simmons BI, White HJ, Zhang H, Aben J,

Abrahamczyk S, Adum GB, Aguilar-Barquero V, Aizen MA, Albertos B, Alcalá EL, del Mar Alguacil M, Alignier A, Ancrenaz M, Andersen AN, Arbeláez-Cortés E, Armbrrecht I, Arroyo-Rodríguez Víctor, Aumann T, Axmacher JC, Azhar B, Azpiroz AB, Baeten L, Bakayoko A, Báldi Aás, Banks JE, Baral SK, Barlow J, Barratt BIP, Barrico L, Bartolommei P, Barton DM, Basset Y, Batáry P, Bates AJ, Baur B, Bayne EM, Beja P, Benedick S, Berg Åke, Bernard H, Berry NJ, Bhatt D, Bicknell JE, Bihn JH, Blake RJ, Bobo KS, Bóçon R, Boekhout T, Böhning-Gaese K, Bonham KJ, Borges PAV, Borges SH, Boutin C, Bouyer J, Bragagnolo C, Brandt JS, Brearley FQ, Brito I, Bros Vç, Brunet J, Buczkowski G, Buddle CM, Bugter R, Buscardo E, Buse J, Cabra-García J, Cáceres NC, Cagle NL, Calviño-Cancela Mía, Cameron SA, Canello EM, Caparrós R, Cardoso P, Carpenter D, Carrijo TF, Carvalho AL, Cassano CR, Castro H, Castro-Luna AA, Rolando CB, Cerezo A, Chapman KA, Chauvat M, Christensen M, Clarke FM, Cleary DFR, Colombo G, Connop SP, Craig MD, Cruz-López L, Cunningham SA, D’Aniello B, D’Cruze N, da Silva PG, Dallimer M, Danquah E, Darvill B, Dauber J, Davis ALV, Dawson J, de Sassi C, de Thoisy B, Deheuvels O, Dejean A, Devineau J-L, Diekötter T, Dolia JV, Domínguez E, Dominguez-Haydar Y, Dorn S, Draper I, Dreber N, Dumont B, Dures SG, Dynesius M, Edenius L, Eggleton P, Eigenbrod F, Elek Zán, Entling MH, Esler KJ, de Lima RF, Faruk A, Farwig N, Fayle TM, Felicioli A, Felton AM, Fensham RJ, Fernandez IC, Ferreira CC, Ficotola GF, Fiera C, Filgueiras BKC, Fırıncioğlu HK, Flaspohler D, Floren A, Fonte SJ, Fournier A, Fowler RE, Franzén M, Fraser LH, Fredriksson GM, Freire GB Jr, Frizzo TLM, Fukuda D, Furlani D, Gaigher Ré, Ganzhorn Jörg U, García KP, Garcia-R JC, Garden JG, Garilleti R, Ge B-M, Gendreau-Berthiaume B, Gerard PJ, Gheler-Costa C, Gilbert B, Giordani P, Giordano S, Golodets C, Gomes LGL, Gould RK, Goulson D, Gove AD, Granjon L, Grass I, Gray CL, Grogan J, et al. 2017. The database of the PREDICTS (Projecting Responses of Ecological Diversity In Changing Terrestrial Systems) project. *Ecology and Evolution* 7(1):145–188 DOI 10.1002/ece3.2579.

Hudson LN, Newbold T, Contu S, Hill SLL, Lysenko I, De Palma A, Phillips HRP, Senior RA, Bennett DJ, Booth H, Choimes A, Correia DLP, Day J, Echeverría-Londoño S, Garon M, Harrison MLK, Ingram DJ, Jung M, Kemp V, Kirkpatrick L, Martin CD, Pan Y, White HJ, Aben J, Abrahamczyk S, Adum GB, Aguilar-Barquero V, Aizen MA, Ancrenaz M, Arbeláez-Cortés E, Armbrrecht I, Azhar B, Azpiroz AB, Baeten L, Báldi A, Banks JE, Barlow J, Batáry P, Bates AJ, Bayne EM, Beja P, Berg Å, Berry NJ, Bicknell JE, Bihn JH, Böhning-Gaese K, Boekhout T, Boutin C, Bouyer J, Brearley FQ, Brito I, Brunet J, Buczkowski G, Buscardo E, Cabra-García J, Calviño-Cancela M, Cameron SA, Canello EM, Carrijo TF, Carvalho AL, Castro H, Castro-Luna AA, Cerda R, Cerezo A, Chauvat M, Clarke FM, Cleary DFR, Connop SP, D’Aniello B, da Silva PG, Darvill B, Dauber J, Dejean A, Diekötter T, Dominguez-Haydar Y, Dormann CF, Dumont B, Dures SG, Dynesius M, Edenius L, Elek Z, Entling MH, Farwig N, Fayle TM, Felicioli A, Felton AM, Ficotola GF, Filgueiras BKC, Fonte SJ, Fraser LH, Fukuda D, Furlani D, Ganzhorn JU, Garden JG, Gheler-Costa C, Giordani P, Giordano S, Gottschalk MS, Goulson D, Gove AD, Grogan J, Hanley ME, Hanson T, Hashim NR, Hawes JE, Hébert C, Helden AJ, Henden J-A, Hernández L, Herzog F, Higuera-Diaz D, Hilje B, Horgan FG, Horváth R, Hylander K, Isaacs-Cubides P, Ishitani M, Jacobs CT, Jaramillo VJ, Jauker B, Jonsell M, Jung TS, Kapoor V, Kati V, Katovai E, Kessler M, Knop E, Kolb A, Kőrösi Á, Lachat T, Lantschner V, Le Féon V, LeBuhn G, Légaré J-P, Letcher SG, Littlewood NA, López-Quintero CA, Louhaichi M, Lövei GL, Lucas-Borja ME, Luja VH, Maeto K, Magura T, Mallari NA, Marin-Spiotta E, Marshall EJP, Martínez E, Mayfield MM, Mikusinski G, Milder JC, Miller JR, Morales CL, Muchane MN, Muchane M, Naidoo R, Nakamura A, Naoe S,

- Nates-Parra G, Navarrete Gutierrez DA, Neuschulz EL, Noreika N, Norfolk O, Noriega JA, Nöske NM, O’Dea N, Oduro W, Ofori-Boateng C, Oke CO, Osgathorpe LM, Paritsis J, Parra-H A, Pelegrin N, Peres CA, Persson AS, Petanidou T, Phalan B, Philips TK, Poveda K, Power EF, Presley SJ, Proença Vânia, Quaranta M, Quintero C, Redpath-Downing NA, Reid JL, Reis YT, Ribeiro DB, Richardson BA, Richardson MJ, Robles CA, Römbke J, Romero-Duque LP, Rosselli L, Rossiter SJ, Roulston TH, Rousseau L, Sadler JP, Sáfián S, Saldaña-Vázquez RA, Samnegård U, et al. 2014. The PREDICTS database: a global database of how local terrestrial biodiversity responds to human impacts. *Ecology and Evolution* 4(24):4701–4735 DOI 10.1002/ece3.1303.
- Huete A, Didan K, Miura T, Rodriguez EP, Gao X, Ferreira LG. 2002. Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sensing of Environment* 83(1–2):195–213 DOI 10.1016/S0034-4257(02)00096-2.
- Hurlbert AH. 2004. Species-energy relationships and habitat complexity in bird communities. *Ecology Letters* 7(8):714–720 DOI 10.1111/j.1461-0248.2004.00630.x.
- Hyndman RJ, Khandakar Y. 2008. Automatic time series forecasting: the forecast Package for R. *Journal of Statistical Software* 27(3):1–22 DOI 10.18637/jss.v027.i03.
- Jiang Z, Huete A, Didan K, Miura T. 2008. Development of a two-band enhanced vegetation index without a blue band. *Remote Sensing of Environment* 112(10):3833–3845 DOI 10.1016/j.rse.2008.06.006.
- Jung M, Hill SLL, Platts PJ, Marchant R, Siebert S, Fournier A, Munyekenye FB, Purvis A, Burgess ND, Newbold T. 2017. Local factors mediate the response of biodiversity to land use on two African mountains. *Animal Conservation* 20(4):370–381 DOI 10.1111/acv.12327.
- Jung M, Rowhani P, Newbold T, Bentley L, Purvis A, Scharlemann JPW. 2019. Local species assemblages are influenced more by past than current dissimilarities in photosynthetic activity. *Ecography* 42(4):670–682 DOI 10.1111/ecog.04031.
- Jung M, Rowhani P, Scharlemann JPW. 2019. Impacts of past abrupt land change on local biodiversity globally. *Nature Communications* 10(1):5474 DOI 10.1038/s41467-019-13452-3.
- Jung M, Scharlemann JPW, Rowhani P. 2020. Landscape-wide changes in land use and land cover correlate with, but rarely explain local biodiversity change. *Landscape Ecology* 35(10):2255–2273 DOI 10.1007/s10980-020-01109-2.
- Kerr JT, Southwood TRE, Cihlar J. 2001. Remotely sensed habitat diversity predicts butterfly species richness and community similarity in Canada. *Proceedings of the National Academy of Sciences of the United States of America* 98(20):11365–11370 DOI 10.1073/pnas.201398398.
- König C, Weigelt P, Kreft H. 2017. Dissecting global turnover in vascular plants. *Global Ecology and Biogeography* 26(2):228–242 DOI 10.1111/geb.12536.
- Leibold MA, Holyoak M, Mouquet N, Amarasekare P, Chase JM, Hoopes MF, Holt RD, Shurin JB, Law R, Tilman D, Loreau M, Gonzalez A. 2004. The metacommunity concept: a framework for multi-scale community ecology. *Ecology Letters* 7(7):601–613 DOI 10.1111/j.1461-0248.2004.00608.x.
- Magurran AE. 2004. *Measuring biological diversity*. Oxford: Blackwell publishing company.
- Mesgaran MB, Cousens RD, Webber BL. 2014. Here be dragons: a tool for quantifying novelty due to covariate range and correlation change when projecting species distribution models. *Diversity and Distributions* 20(10):1147–1159 DOI 10.1111/ddi.12209.
- Meyer H, Pebesma E. 2021. Predicting into unknown space? Estimating the area of applicability of spatial prediction models. *Methods in Ecology and Evolution* 12(9):1620–1633 DOI 10.1111/2041-210X.13650.

- Miller JR, Turner MG, Smithwick EAH, Dent CL, Stanley EH. 2004. Spatial extrapolation: the science of predicting ecological patterns and processes. *BioScience* 54(4):310 DOI 10.1641/0006-3568(2004)054[0310:SETSOP]2.0.CO;2.
- Montgomery GA, Belitz MW, Guralnick RP, Tingley MW. 2021. Standards and best practices for monitoring and benchmarking insects. *Frontiers in Ecology and Evolution* 8:282 DOI 10.3389/fevo.2020.579193.
- Moritz S, Bartz-Beielstein T. 2017. imputeTS: time series missing value imputation in R. *The R Journal* 9(1):207 DOI 10.32614/RJ-2017-009.
- Mueller T, Dressler G, Tucker C, Pinzon J, Leimgruber P, Dubayah R, Hurtt G, Böhning-Gaese K, Fagan W. 2014. Human land-use practices lead to global long-term increases in photosynthetic capacity. *Remote Sensing* 6(6):5717–5731 DOI 10.3390/rs6065717.
- Newbold T, Hudson LN, Contu S, Hill SLL, Beck J, Liu Y, Meyer C, Phillips HRP, Scharlemann JPW, Purvis A. 2018. Widespread winners and narrow-ranged losers: land use homogenizes biodiversity in local assemblages worldwide. *PLOS Biology* 16(12):e2006841 DOI 10.1371/journal.pbio.2006841.
- Newbold T, Hudson LN, Hill SLL, Contu S, Lysenko I, Senior RA, Börger L, Bennett DJ, Choimes A, Collen B, Day J, De Palma A, Díaz S, Echeverria-Londoño S, Edgar MJ, Feldman A, Garon M, Harrison MLK, Alhousseini T, Ingram DJ, Itescu Y, Kattge J, Kemp V, Kirkpatrick L, Kleyer M, Correia DLP, Martin CD, Meiri S, Novosolov M, Pan Y, Phillips HRP, Purves DW, Robinson A, Simpson J, Tuck SL, Weiher E, White HJ, Ewers RM, Mace GM, Scharlemann JPW, Purvis A. 2015. Global effects of land use on local terrestrial biodiversity. *Nature* 520(7545):45–50 DOI 10.1038/nature14324.
- Oldeland J, Wesuls D, Rocchini D, Schmidt M, Jürgens N. 2010. Does using species abundance data improve estimates of species diversity from remotely sensed spectral heterogeneity? *Ecological Indicators* 10(2):390–396 DOI 10.1016/j.ecolind.2009.07.012.
- Parmentier I, Harrigan RJ, Buermann W, Mitchard ETA, Saatchi S, Malhi Y, Bongers F, Hawthorne WD, Leal ME, Lewis SL, Nusbaumer L, Sheil D, Sosef MSM, Affum-Baffoe K, Bakayoko A, Chuyong GB, Chatelain C, Comiskey JA, Dauby G, Doucet J-L, Fauset S, Gautier L, Gillet J-F, Kenfack D, Kouamé FN, Kouassi EK, Kouka LA, Parren MPE, Peh KSH, Reitsma JM, Senterre B, Sonké B, Sunderland TCH, Swaine MD, Tchouto MGP, Thomas D, Van Valkenburg JLCH, Hardy OJ. 2011. Predicting alpha diversity of African rain forests: models based on climate and satellite-derived data do not perform better than a purely spatial model. *Journal of Biogeography* 38(6):1164–1176 DOI 10.1111/j.1365-2699.2010.02467.x.
- Petchey OL, Pontarp M, Massie TM, Kéfi S, Ozgul A, Weilenmann M, Palamara GM, Altermatt F, Matthews B, Levine JM, Childs DZ, McGill BJ, Schaepman ME, Schmid B, Spaak P, Beckerman AP, Pennekamp F, Pearse IS. 2015. The ecological forecast horizon, and examples of its uses and determinants. *Ecology Letters* 18(7):597–611 DOI 10.1111/ele.12443.
- Pettorelli N, Gaillard J-M, Mysterud A, Duncan P, Chr. Stenseth N, Delorme D, Van Laere G, Toïgo C, Klein F. 2006. Using a proxy of plant productivity (NDVI) to find key periods for animal performance: the case of roe deer. *Oikos* 112(3):565–572 DOI 10.1111/j.0030-1299.2006.14447.x.
- Pettorelli N, Vik JO, Mysterud A, Gaillard J-M, Tucker CJ, Stenseth NC. 2005. Using the satellite-derived NDVI to assess ecological responses to environmental change. *Trends in Ecology & Evolution* 20(9):503–510 DOI 10.1016/j.tree.2005.05.011.
- Phillips HRP, Guerra CA, Bartz MLC, Briones MJI, Brown G, Crowther TW, Ferlian O, Gongalsky KB, van den Hoogen J, Krebs J, Orgiazzi A, Routh D, Schwarz B, Bach EM, Bennett JM, Brose U, Decaëns T, König-Ries B, Loreau M, Mathieu J, Mulder C,

van der Putten WH, Ramirez KS, Rillig MC, Russell D, Rutgers M, Thakur MP, de Vries FT, Wall DH, Wardle DA, Arai M, Ayuke FO, Baker GH, Beauséjour R, Bedano Jé C, Birkhofer K, Blanchart E, Blossey B, Bolger T, Bradley RL, Callaham MA, Capowiez Y, Caulfield ME, Choi A, Crotty FV, Crumsey JM, Dávalos A, Diaz Cosin DJ, Dominguez Aí, Duhour AE, van Eekeren N, Emmerling C, Falco LB, Fernández R, Fonte SJ, Fragoso C, Franco ALC, Fugère M, Fusilero AT, Gholami S, Gundale MJ, López MG, Hackenberger DK, Hernández LM, Hishi T, Holdsworth AR, Holmstrup M, Hopfensperger KN, Lwanga EH, Huhta V, Hurisso TT, Iannone BV III, Iordache M, Joschko M, Kaneko N, Kanianska R, Keith AM, Kelly CA, Kernecker ML, Klaminder J, Koné AW, Kooch Y, Kukkonen ST, Lalthanzara H, Lammel DR, Lebedev IM, Li Y, Jesus Lidon JB, Lincoln NK, Loss SR, Marichal R, Matula R, Moos JH, Moreno G, Morón-Ríos A, Muys B, Neiryneck J, Norgrove L, Novo M, Nuutinen V, Nuzzo V, Pansu J, Paudel S, Pérès G, Pérez-Camacho L, Piñeiro R, Ponge J-F, Rashid MI, Rebollo S, Rodeiro-Iglesias J, Rodríguez MÁ, Roth AM, Rousseau GX, Rozen A, Sayad E, van Schaik L, Scharenbroch BC, Schirrmann M, Schmidt O, Schröder B, Seeber J, Shashkov MP, Singh J, Smith SM, Steinwandter M, Talavera JA, Trigo D, Tsukamoto J, de Valença AW, Vanek SJ, Virto I, Wackett AA, Warren MW, Wehr NH, Whalen JK, Wironen MB, Wolters V, Zenkova IV, Zhang W, Cameron EK, Eisenhauer N. 2019. Global distribution of earthworm diversity. *Science* 366(6464):480–485
DOI 10.1126/science.aax4851.

Phillips HRP, Newbold T, Purvis A. 2017. Land-use effects on local biodiversity in tropical forests vary between continents. *Biodiversity and Conservation* 26(9):2251–2270
DOI 10.1007/s10531-017-1356-2.

Purvis A, Newbold T, De Palma A, Contu S, Hill SLL, Sanchez-Ortiz K, Phillips HRP, Hudson LN, Lysenko I, Börger L, Scharlemann JPW. 2018. Modelling and projecting the response of local terrestrial biodiversity worldwide to land use and related pressures: the PREDICTS project. *Advances in Ecological Research* 58:201–241
DOI 10.1016/bs.aecr.2017.12.003.

Radeloff VC, Dubinin M, Coops NC, Allen AM, Brooks TM, Clayton MK, Costa GC, Graham CH, Helmers DP, Ives AR, Kolesov D, Pidgeon AM, Rapacciuolo G, Razenkova E, Suttidate N, Young BE, Zhu L, Hobi ML. 2019. The dynamic habitat indices (DHIs) from MODIS and global biodiversity. *Remote Sensing of Environment* 222:204–214
DOI 10.1016/j.rse.2018.12.009.

Randin CF, Ashcroft MB, Bolliger J, Cavender-Bares J, Coops NC, Dullinger S, Dirnböck T, Eckert S, Ellis E, Fernández N, Giuliani G, Guisan A, Jetz W, Joost S, Karger D, Lembrechts J, Lenoir J, Luoto M, Morin X, Price B, Rocchini D, Schaeppman M, Schmid B, Verburg P, Wilson A, Woodcock P, Yoccoz N, Payne D. 2020. Monitoring biodiversity in the Anthropocene using remote sensing in species distribution models. *Remote Sensing of Environment* 239(9):111626 DOI 10.1016/j.rse.2019.111626.

Rapacciuolo G. 2019. Strengthening the contribution of macroecological models to conservation practice. *Global Ecology and Biogeography* 28(1):54–60 DOI 10.1111/geb.12848.

Regos A, Gagne L, Alcaraz-Segura D, Honrado JP, Domínguez J. 2019. Effects of species traits and environmental predictors on performance and transferability of ecological niche models. *Scientific Reports* 9(1):4221 DOI 10.1038/s41598-019-40766-5.

Regos A, Gonçalves J, Arenas-Castro S, Alcaraz-Segura D, Guisan A, Honrado JP. 2022. Mainstreaming remotely sensed ecosystem functioning in ecological niche models. *Remote Sensing in Ecology and Conservation* 8:431–447 DOI 10.1002/rse2.255.

Roberts DR, Bahn V, Ciuti S, Boyce MS, Elith J, Guillera-Aroita G, Hauenstein S, Lahoz-Monfort JJ, Schröder B, Thuiller W, Warton DI, Wintle BA, Hartig F, Dormann CF.

2017. Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic structure. *Ecography* **40**(8):913–929 DOI [10.1111/ecog.02881](https://doi.org/10.1111/ecog.02881).
- Rocchini D, Balkenhol N, Carter GA, Foody GM, Gillespie TW, He KS, Kark S, Levin N, Lucas K, Luoto M, Nagendra H, Oldeland J, Ricotta C, Southworth J, Neteler M. 2010. Remotely sensed spectral heterogeneity as a proxy of species diversity: recent advances and open challenges. *Ecological Informatics* **5**(5):318–329 DOI [10.1016/j.ecoinf.2010.06.001](https://doi.org/10.1016/j.ecoinf.2010.06.001).
- Rocchini D, Hernández-Stefanoni JL, He KS. 2015. Advancing species diversity estimate by remotely sensed proxies: a conceptual review. *Ecological Informatics* **25**:22–28 DOI [10.1016/j.ecoinf.2014.10.006](https://doi.org/10.1016/j.ecoinf.2014.10.006).
- Royle JA, Nichols JD, Kéry M. 2005. Modelling occurrence and abundance of species when detection is imperfect. *Oikos* **110**(2):353–359 DOI [10.1111/j.0030-1299.2005.13534.x](https://doi.org/10.1111/j.0030-1299.2005.13534.x).
- Santini L, Antão LH, Jung M, Benítez-López A, Rapacciuolo G, Di Marco M, Jones FAM, Haghkerdar JM, González-Suárez M. 2021. The interface between Macroecology and Conservation: existing links and untapped opportunities. *Frontiers of Biogeography* **13**(4):e53025 DOI [10.21425/F5FBG53025](https://doi.org/10.21425/F5FBG53025).
- Schaaf CB, Gao F, Strahler AH, Lucht W, Li X, Tsang T, Strugnell NC, Zhang X, Jin Y, Muller J-P, Lewis P, Barnsley M, Hobson P, Disney M, Roberts G, Dunderdale M, Doll C, D'Entremont RP, Hu B, Liang S, Privette JL, Roy D. 2002. First operational BRDF, albedo nadir reflectance products from MODIS. *Remote Sensing of Environment* **83**(1–2):135–148 DOI [10.1016/S0034-4257\(02\)00091-3](https://doi.org/10.1016/S0034-4257(02)00091-3).
- Schmidtlein S, Fassnacht FE. 2017. The spectral variability hypothesis does not hold across landscapes. *Remote Sensing of Environment* **192**(18):114–125 DOI [10.1016/j.rse.2017.01.036](https://doi.org/10.1016/j.rse.2017.01.036).
- Shackelford N, Starzomski BM, Banning NC, Battaglia LL, Becker A, Bellingham PJ, Bestelmeyer B, Catford JA, Dwyer JM, Dynesius M, Gilmour J, Hallett LM, Hobbs RJ, Price J, Sasaki T, Tanner EVJ, Standish RJ. 2017. Isolation predicts compositional change after discrete disturbances in a global meta-study. *Ecography* **40**(11):1256–1266 DOI [10.1111/ecog.02383](https://doi.org/10.1111/ecog.02383).
- Stein A, Gerstner K, Kreft H. 2014. Environmental heterogeneity as a universal driver of species richness across taxa, biomes and spatial scales. *Ecology Letters* **17**(7):866–880 DOI [10.1111/ele.12277](https://doi.org/10.1111/ele.12277).
- Stein A, Kreft H. 2015. Terminology and quantification of environmental heterogeneity in species-richness research. *Biological Reviews* **90**(3):815–836 DOI [10.1111/brv.12135](https://doi.org/10.1111/brv.12135).
- Supp SR, Ernest SKM. 2014. Species-level and community-level responses to disturbance: a cross-community analysis. *Ecology* **95**(7):1717–1723 DOI [10.1890/13-2250.1](https://doi.org/10.1890/13-2250.1).
- Tredennick AT, Hooker G, Ellner SP, Adler PB. 2021. A practical guide to selecting models for exploration, inference, and prediction in ecology. *Ecology* **102**(6):267 DOI [10.1002/ecy.3336](https://doi.org/10.1002/ecy.3336).
- van den Hoogen J, Geisen S, Routh D, Ferris H, Traunspurger W, Wardle DA, de Goede RGM, Adams BJ, Ahmad W, Andriuzzi WS, Bardgett RD, Bonkowski M, Campos-Herrera R, Cares JE, Caruso T, de Brito Caixeta L, Chen X, Costa SR, Creamer R, Mauro da Cunha Castro Jé, Dam M, Djigal D, Escuer M, Griffiths BS, Gutiérrez C, Hohberg K, Kalinkina D, Kardol P, Kergunteuil A, Korthals G, Krashevska V, Kudrin AA, Li Q, Liang W, Magilton M, Marais M, Martín JAR, Matveeva E, Mayad EH, Mulder C, Mullin P, Neilson R, Nguyen TAD, Nielsen UN, Okada H, Rius JEP, Pan K, Peneva V, Pellissier Lic, Carlos Pereira da Silva J, Pitteloud C, Powers TO, Powers K, Quist CW, Rasmann S, Moreno SS, Scheu S, Setälä H, Sushchuk A, Tiunov AV, Trap J, van der Putten W, Vestergård M, Villenave C, Waeyenberge L, Wall DH, Wilschut R, Wright DG, Yang J-I, Crowther TW.

2019. Soil nematode abundance and functional group composition at a global scale. *Nature* 572(7768):194–198 DOI [10.1038/s41586-019-1418-6](https://doi.org/10.1038/s41586-019-1418-6).

- Weiss DJ, Nelson A, Gibson HS, Temperley W, Peedell S, Lieber A, Hancher M, Poyart E, Belchior S, Fullman N, Mappin B, Dalrymple U, Rozier J, Lucas TCD, Howes RE, Tusting LS, Kang SY, Cameron E, Bisanzio D, Battle KE, Bhatt S, Gething PW. 2018. A global map of travel time to cities to assess inequalities in accessibility in 2015. *Nature* 553(7688):333–336 DOI [10.1038/nature25181](https://doi.org/10.1038/nature25181).
- Yates KL, Bouchet PJ, Caley MJ, Mengersen K, Randin CF, Parnell S, Fielding AH, Bamford AJ, Ban S, Barbosa AM, Dormann CF, Elith J, Embling CB, Ervin GN, Fisher R, Gould S, Graf RF, Gregr EJ, Halpin PN, Heikkinen RK, Heinänen S, Jones AR, Krishnakumar PK, Lauria V, Lozano-Montes H, Mannocci L, Mellin C, Mesgaran MB, Moreno-Amat E, Mormede S, Novaczek E, Opiel S, Ortuño Crespo G, Peterson AT, Rapacciuolo G, Roberts JJ, Ross RE, Scales KL, Schoeman D, Snelgrove P, Sundblad G, Thuiller W, Torres LG, Verbruggen H, Wang L, Wenger S, Whittingham MJ, Zharikov Y, Zurell D, Sequeira AMM. 2018. Outstanding challenges in the transferability of ecological models. *Trends in Ecology & Evolution* 33(10):790–802 DOI [10.1016/j.tree.2018.08.001](https://doi.org/10.1016/j.tree.2018.08.001).
- Yin H, Pflugmacher D, Kennedy RE, Sulla-Menashe D, Hostert P. 2014. Mapping annual land use and land cover changes using MODIS time series. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* 7(8):3421–3427 DOI [10.1109/JSTARS.2014.2348411](https://doi.org/10.1109/JSTARS.2014.2348411).
- Zurell D, Elith J, Schröder B. 2012. Predicting to new environments: tools for visualizing model behaviour and impacts on mapped distributions. *Diversity and Distributions* 18(6):628–634 DOI [10.1111/j.1472-4642.2012.00887.x](https://doi.org/10.1111/j.1472-4642.2012.00887.x).