



International Institute for  
Applied Systems Analysis  
[www.iiasa.ac.at](http://www.iiasa.ac.at)

# **A Model for Estimating Nitrogen Oxide Emissions in Europe**

**Luebkert, B.**

**IIASA Working Paper**

**WP-87-122**

**December 1987**



Luebkert B (1987). A Model for Estimating Nitrogen Oxide Emissions in Europe. IIASA Working Paper. IIASA, Laxenburg, Austria: WP-87-122 Copyright © 1987 by the author(s). <http://pure.iiasa.ac.at/id/eprint/2930/>

**Working Papers** on work of the International Institute for Applied Systems Analysis receive only limited review. Views or opinions expressed herein do not necessarily represent those of the Institute, its National Member Organizations, or other organizations supporting the work. All rights reserved. Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage. All copies must bear this notice and the full citation on the first page. For other purposes, to republish, to post on servers or to redistribute to lists, permission must be sought by contacting [repository@iiasa.ac.at](mailto:repository@iiasa.ac.at)

# ***WORKING PAPER***

## A MODEL FOR ESTIMATING NITROGEN OXIDE EMISSIONS IN EUROPE

Barbara Lübkert

December 1987

WP-87-122

# **A MODEL FOR ESTIMATING NITROGEN OXIDE EMISSIONS IN EUROPE**

*Barbara Lübkert*

December 1987

WP-87-122

*Working Papers* are interim reports on work of the International Institute for Applied Systems Analysis and have received only limited review. Views or opinions expressed herein do not necessarily represent those of the Institute or of its National Member Organizations.

INTERNATIONAL INSTITUTE FOR APPLIED SYSTEMS ANALYSIS  
A-2361 Laxenburg, Austria

## **PREFACE**

IIASA's RAINS model have so far dealt with the emissions and effects of sulphur dioxide. However, acidification is caused by other pollutants as well. In particular, oxides of nitrogen play an important role. Barbara Lübkert joined the Acid Rain Project during the summer of 1986 as a participant in the Young Scientists' Summer Program and developed a method to estimate the emissions of nitrogen oxides. This paper describes the method she used and presents the results.

Her work is very important in that it makes the first step toward a RAINS model that covers nitrogen as well as sulphur.

Roderick W. Shaw  
Leader  
Acid Rain Project

## ACKNOWLEDGEMENTS

The work presented in this paper was supported by the Organisation for Economic Co-operation and Development (OECD) and carried out by the author during several visits to IIASA; she is grateful to Erik Lykke and Pierre Lieben at the OECD Environment Directorate as well as to Leen Hordijk, who made this series of visits possible. The author further thanks Soizick de Tilly, Peter Lübkert, and Steven Beatson of the OECD Pollution Control Division and her colleagues in OECD Member countries who helped in compiling and revising emission data used in this study; thanks are also due to Markus Amann, Stuart Batterman and Wolfgang Schöpp of IIASA's Acid Rain Project for their constructive ideas and Joseph Alcamo for his critical review of this paper.

## ABSTRACT

This paper presents a methodology to calculate nitrogen oxide ( $NO_x$ ) emissions in a consistent way for all European countries. First, the principles of  $NO_x$  formation are reviewed, and results of a feasibility study carried out to develop an  $NO_x$  emission submodel for inclusion into IIASA's RAINS model to quantify acid deposition in Europe are summarized. It is shown that  $NO_x$  emissions are most closely correlated to fossil fuel consumption and that this consumption is, therefore, the best single statistical parameter to estimate a country's  $NO_x$  emissions from the various source sectors. Existing  $NO_x$  emission inventories and emission factors are analyzed, and a set of comparable data from the OECD emission inventory for major air pollutants (OECD, 1987) is identified which is appropriate for statistical analysis in order to estimate sector- and fuel-specific emission coefficients. Such aggregate coefficients are determined using ordinary least squares (OLS) regression analysis for the sector/fuel combinations presently contained in RAINS to calculate sulfur emissions. Fuel data used in the regression analysis are from the International Energy Agency (IEA, 1987a,b).

Emission coefficients determined in this way are entered into RAINS and  $NO_x$  emissions are determined for all 27 European countries. Comparison of the results with reported national totals as well as reported traffic  $NO_x$  shows generally good agreement, i.e., within 20 percent. If  $NO_x$  emissions calculated by RAINS are compared to EMEP estimates (United Nations, 1987) on a total European scale (USSR excluded), RAINS overestimates the EMEP total by only four percent. Relative contributions as calculated by RAINS for the transportation sector also reflect numbers reported by countries, being on the average 54 percent of total  $NO_x$  for OECD Europe and 16 percent for non-OECD Europe (excluding the USSR). A qualitative discussion about the uncertainty of estimated emission coefficients and resulting emission rates is also included. The approach described in this paper is, thus, promising and shows a way in which one can use a set of comparable data in a "top-down" approach to extrapolate the data to other countries where less detailed data are available. The method also allows for testing future emission scenarios with and without assumed emission reduction policies.

## TABLE OF CONTENTS

|                                                       |    |
|-------------------------------------------------------|----|
| 1. Introduction                                       | 1  |
| 2. Goals and Limitations of the $NO_x$ Emission Model | 1  |
| 3. Principles of $NO_x$ Formation                     | 3  |
| 3.1. Fuel $NO_x$                                      | 3  |
| 3.2. Thermal $NO_x$                                   | 5  |
| 3.3. "Prompt" $NO_x$                                  | 5  |
| 4. Approaches to $NO_x$ Emission Calculations         | 5  |
| 4.1. Theoretical $NO_x$ Emission Calculations         | 5  |
| 4.2. Definition of $NO_x$ Emitting Source Categories  | 6  |
| 4.3. Use of Emission Factors                          | 6  |
| 4.4. Existing Emission Inventories                    | 7  |
| 4.5. Analysis of the OECD $NO_x$ Emission Inventory   | 9  |
| 4.6. Review of $SO_2$ Emission Calculations in RAINS  | 10 |
| 5. A Method for Calculating $NO_x$ Emissions          | 10 |
| 6. Results                                            | 12 |
| 7. Discussion                                         | 17 |
| 8. Summary and Conclusions                            | 18 |
| References                                            | 19 |



# A MODEL FOR ESTIMATING NITROGEN OXIDE EMISSIONS IN EUROPE

Barbara Lübkert

## 1. INTRODUCTION

The Acid Rain Project at IIASA has developed a set of interactive, computer-based models – RAINS (Regional Acidification Information and Simulation) – to assess long-term acidification in Europe on a regional scale. To date, the RAINS model estimates current and future levels of acid deposition and their impacts on the environment due to sulfur compounds only. The available submodels first estimate emissions of sulfur due to a chosen energy forecast (Economic Commission for Europe (ECE), 1987; International Energy Agency (IEA), 1983), then relate these emissions to atmospheric deposition via a source-receptor matrix (Eliassen and Saltbones, 1983), and lastly predict impacts on forest soils, trees, lakes and groundwater (Kauppi *et al.*, 1986; Mäkelä *et al.*, 1987; Kämäri and Posch, 1987; Holmberg *et al.*, 1987). The model further has the option of evaluating the effect of various emission reduction scenarios and their associated costs (Amann and Kornai, 1987). Since the model is designed to be especially useful to decision makers, particular emphasis has been put on easy-to-use, interactive computer software and good, comprehensible graphical representation of results. The reader is referred to Alcamo *et al.* (1987) for a more detailed description of RAINS and Hordijk (1986) for some policy applications of the model.

To date, we know that acidification of the environment is the result of a combined effect of dry and wet deposition of sulfur and nitrogen compounds. In order to make RAINS more comprehensive it has, therefore, become desirable to also include nitrogen oxides ( $NO_x$ ) and ammonia ( $NH_3$ ) into the model. This paper presents the results of a feasibility study carried out to develop an  $NO_x$  emission submodel, describes the methodology developed, presents  $NO_x$  emissions obtained by this method, and gives a comparison of these results with existing reported nitrogen oxide emission data. Results from this paper can be used to construct an  $NO_x$  emissions submodel which will be part of an expanded version of RAINS, as depicted in *Figure 1*.

## 2. GOALS AND LIMITATIONS OF THE $NO_x$ EMISSION MODEL

The goals of the  $NO_x$  emission model are:

1. To provide a consistent method for calculating  $NO_x$  emissions in all European countries.
2. To base these calculations on statistical data which are comparable and readily available for all countries.




Figure 1. Schematic of the expanded IIASA RAINS model.

3. To be able to calculate emissions for different years in the past, present and future by making use of forecasts of the main variables influencing these emissions.
4. To be able to calculate emissions under different emission reduction scenarios.

Given these objectives a model must be developed by which emissions can be estimated in a relatively simple and aggregated way and therefore one should keep the number of variables that influence predictions to a minimum. Further, these main influencing parameters should be readily available from statistics, preferably collected on an international level which guarantee some compatibility.

The  $NO_x$  emission model presented in this paper is not intended to replace any of the detailed "bottom-up" approaches that are used in different countries to inventory past and present  $NO_x$  emissions, but is meant to show a way in which one can use a set of comparable data in a "top-down" approach to extrapolate these data to other countries and years.

### 3. PRINCIPLES OF $NO_x$ FORMATION

Nitrogen oxides are emitted from man-made and natural sources. Man-made emissions are generated during fuel combustion and in industrial processes; they are emitted in the form of nitric oxide ( $NO$ ) and nitrogen dioxide ( $NO_2$ ). For most emission source categories, the  $NO_2$  fraction of total  $NO_x$  ( $NO$  and  $NO_2$  together) emissions is lower than 10 percent with the exception of plants producing nitric acid and/or ammonia (40 percent) and for gas turbines (15 percent) (Bakkum and Veldt, 1986a). Natural nitrogen oxide emissions are predominantly in the non-reactive form of nitrous oxide ( $N_2O$ ) and are, thus, not believed to contribute to the acid-forming potential of the environment. Of all direct anthropogenic  $NO_x$  emissions, those from combustion processes in stationary and mobile sources constitute the overwhelming majority.

Nitrogen oxides are formed during combustion due to three principal mechanisms which result in the so-called (1) fuel  $NO_x$ , (2) thermal  $NO_x$ , and (3) "prompt"  $NO_x$ . Emissions of nitrogen oxides from industrial processes result from processing raw materials that contain nitrogen either as a necessary constituent or as a contaminant. Such emissions depend on the nitrogen content of the raw material and on its volatility. It is shown later that these process emissions are, however, of secondary importance.

#### 3.1. Fuel $NO_x$

Fuel  $NO_x$  formation is a function of the fuel's nitrogen content as well as burner type and firing mode used in the combustion. Fuel nitrogen contents vary typically between 0.5 and 2.0 percent (by weight) for coal and shale, and between 0.5 and 1.5 percent for synthetic fuels derived from coal. The nitrogen content is less than 1 percent for oil and gas, and negligible for natural gas (US Department of Energy, 1983). Fuel  $NO_x$  is, however, also dependent on the burner type for which no simple relationship exists. *Figure 2* illustrates this complicated relationship.






Figure 2. Effects of firing pattern on  $NO_x$  emissions (Thompson, 1979; cited in Morrison, 1980).

### 3.2. Thermal $NO_x$

Thermal  $NO_x$  formation is due to the mechanism discovered by Zeldovich (1946), in which nitrogen and oxygen from the air combine to form  $NO_x$  under high temperatures:



This formation process becomes important usually at temperatures above 1400 °C (Rentz, 1984), a temperature which is generally exceeded in most combustion processes. Thermal  $NO_x$  formation increases exponentially at higher temperatures, and is also a function of residence time of air in the combustion chamber and of the stoichiometry, i.e., the excess air availability.

To overall  $NO_x$  emissions, thermal  $NO_x$  contributes around 20 percent in coal combustion, about 50 percent in heavy oil combustion, and the largest portion in light oil and gas combustion (Mason and Herther, 1982).

### 3.3. "Prompt" $NO_x$

In the flame reaction zone, significantly higher amounts of  $NO_x$  are produced than can be explained by equilibrium calculations. This  $NO_x$  is termed "prompt"  $NO_x$ , but is generally an order of magnitude lower than thermal and fuel  $NO_x$  and is therefore considered negligible in this paper.

## 4. APPROACHES TO $NO_x$ EMISSION CALCULATIONS

### 4.1. Theoretical $NO_x$ Emission Calculations

As explained above, total  $NO_x$  emission from combustion is the sum of thermal, fuel and "prompt"  $NO_x$  emissions:

$$e(NO_x) = e(fuel-N) + e(thermal) + e(prompt) \quad [1]$$

in which

|               |                                                             |
|---------------|-------------------------------------------------------------|
| $e(NO_x)$     | = total of the $NO_x$ emissions [tonnes],                   |
| $e(fuel-N)$   | = f(fuel, firing mode),                                     |
| $e(thermal)$  | = f(combustion temperature, residence time, stoichiometry), |
| $e("prompt")$ | = 0.                                                        |

Assuming different combustion temperatures, burner types and firing modes in different economic sectors and for different fuels, emissions should, therefore, also depend on the sector in which they are emitted and on the fuel type used:

$$e(NO_x) = \sum_{i,j} e_{ij}(NO_x) \quad , \quad [2]$$

where  $e_{ij}(NO_x)$  is the emission rate per sector per fuel, and

$$e_{ij}(NO_x) = n_j * E_{ij} / h v_j * (1 - a_i) + \alpha * E_{ij} * (1 - a_i) \quad [3]$$

in which

|          |   |                                                                                                                                    |
|----------|---|------------------------------------------------------------------------------------------------------------------------------------|
| $n_j$    | : | nitrogen content of fuel [weight %],                                                                                               |
| $E_{ij}$ | : | energy consumed per sector per fuel [PJ/yr],                                                                                       |
| $h v_j$  | : | heat content per fuel [PJ/tonne],                                                                                                  |
| $a_i$    | : | nitrogen removed by flue gas cleaning in each sector [fraction],                                                                   |
| $\alpha$ | : | emission factor due to the thermal $NO_x$ ; a function of combustion temperature, burner type and configuration, etc. [tonnes/PJ], |
| $i$      | : | emission sector,                                                                                                                   |
| $j$      | : | fuel type.                                                                                                                         |

In practice, it is difficult to use Eq. [3] because the emission factor due to thermal  $NO_x$ ,  $\alpha$ , varies significantly as a function of combustion parameters which are not fuel specific, i.e.,  $\alpha$  can vary within one sector and fuel as much as between sectors and fuels.

Due to the high variability of  $NO_x$  emissions with combustion conditions, it would be necessary to subdivide the emission source categories into various equipment and fuel combinations having distinct emission characteristics. As described by Mason and Herthier (1982), such an approach has resulted in over 150 of these equipment/fuel combinations in preparing an  $NO_x$  emission inventory for the USA for 1980. That many source categories seem presently infeasible in an  $NO_x$  emission calculation model on a European level, in which input data should be comparable between countries.

## 4.2. Definition of $NO_x$ Emitting Source Categories

Nitrogen oxide emitting sources can be grouped according to economic activities and/or  $NO_x$  generating processes; refineries are an example of an economic sector whereas furnaces are an example of the latter category. In preparing emission inventories as well as in making forecasts, it is desirable to aggregate source categories in a way that allows, (1) emissions to be calculated in a uniform manner for the entire source category and (2) emission reduction scenarios to be evaluated. The latter means that one needs to be able to account for control scenarios that appear feasible for a specific economic sector or a particular emission generating process. Broad categories emitting  $NO_x$  are (1) mobile sources, including on- and off-road traffic, (2) power plants, (3) industrial and (4) non-industrial combustion, (5) industrial processes, in particular, ammonia and nitric acid plants, (6) agricultural sources, such as open burning, and (7) waste incineration. Of these seven categories, the last three are generally of minor importance at the level of national emissions.

## 4.3. Use of Emission Factors

Since in almost all countries, emission measurements are not carried out on a routine basis, emission rates are usually based on mass balance calculations and specific point source measurements. This knowledge is translated into an emission factor which represents an extrapolation of point-specific data to an entire emission source category. Therefore, physical verification is not possible of these emission factors nor of the subsequently calculated total emission rates.

It has been stated that, nowadays, knowledge about emission factors for major air pollutants, including  $NO_x$ , has accumulated to a level where, for the most important emission source categories, reasonable estimates can be made for total emission rates in these sectors (Bouscaren *et al.*, 1986). Reviewing recent publications and other available lists of

$NO_x$  emission factors shows, however, that ranges of  $NO_x$  factors are still so large that their use requires some specific knowledge of the sector and its structure in order to give reasonable estimates. *Tables 1 and 2* (OECD-Air Management Policy Group, 1984; Löblich, 1985) give an overview of such ranges of emission factors for  $NO_x$ . It should be noted that these tables are already the result of some extensive literature research trying to present emission factor ranges that are applicable at least to the western European situation.

*Table 1.*  $NO_x$  emission factor ranges found in the literature (OECD - Air Management Policy Group, 1984; Löblich, 1985).

| <i>Sector</i>  | <i>Hard Coal</i> | <i>Brown Coal</i> | <i>Derived Coal</i> | <i>Gas</i> |
|----------------|------------------|-------------------|---------------------|------------|
| Transportation | —                | —                 | —                   | —          |
| Power Plants   | 280–630          | 180–290           | —                   | 50–140     |
| Domestic       | 45–100           | 50–300            | 70–100              | 30–120     |
| Industry       | 200–460          | 200–300           | 180–540             | 40–170     |

| <i>Sector</i>  | <i>Light Fraction of Oil</i>    | <i>Middle Distillate</i>        | <i>Heavy Oil</i> | <i>Other Solid Fuels</i> |
|----------------|---------------------------------|---------------------------------|------------------|--------------------------|
| Transportation | 1050<br>[=44 g/kg] <sup>1</sup> | 1175<br>[=50 g/kg] <sup>2</sup> | —                | —                        |
| Power Plants   | —                               | 60–150                          | 150–245          | 120–300                  |
| Domestic       | 50–100                          | 50–160                          | 140–240          | 50–220                   |
| Industry       | 45–60                           | 60–190                          | 140–240          | 80–300                   |

Units: tonnes  $NO_2$ /PJ input.

Notes:

(1) for gasoline automobiles (see *Table 2*).

(2) for diesel trucks (see *Table 2*).

N.B.: All emission factors proposed by OECD were elaborated in cooperation with TNO (The Netherlands) as average best estimates.

The literature as well as *Tables 1 and 2* show that currently used aggregate  $NO_x$  emission factors are fuel-specific (coal, oil, gas, and other solid fuels) and distinguish between different types of coal and oil. Industrial process emissions are generally related to the amount of product manufactured. The use of such emission factors, therefore, requires detailed statistical data on fuel use and production rates in each of the economic sectors. Usually, these data are not readily available from international statistics. When available from national statistics, they are often not comparable between countries.

#### 4.4. Existing Emission Inventories

Some countries have a significant experience in gathering emission data for major air pollutants. These data are usually in the form of emission inventories for particular years and, in many cases, present information about emissions per source category according to the various economic sectors (e.g., power plants, refineries, non-ferrous metal industry, dry cleaning, agriculture, etc.), and/or by type of emission generating process (e.g., combustion, raw material and product storage, industrial processes, etc.). As noted above, a serious problem, however, is the usual lack of comparability or compatibility

Table 2. Emission factors for the transportation sector (road traffic).

| Vehicle Type      | Fuel Type |           |          | Literature Reference |
|-------------------|-----------|-----------|----------|----------------------|
|                   | Gasoline  | Diesel    | LPG      |                      |
| AUTOMOBILE        |           |           |          |                      |
| [g/km]            | 2.54–2.56 | 0.62–0.74 |          | (1)                  |
|                   | 2.17      | 1.06      |          | (2)                  |
|                   | 1.3–3.8   | 0.9–1.8   | 1.25–2.5 | (3)                  |
| [g/kg fuel]       | 44        | 20        | 33       | (1)                  |
| LIGHT-DUTY TRUCKS |           |           |          |                      |
| [g/km]            | 5.17      | 0.62–0.74 |          | (1)                  |
|                   | 2.0–11.0  | 1.1–3.8   | 1.6–3.2  | (3)                  |
| [g/kg fuel]       | 65        | 22        |          | (1)                  |
| HEAVY-DUTY TRUCKS |           |           |          |                      |
| [g/km]            | 4.1       | 11.5–23.1 |          | (1)                  |
|                   | 6.0–14.0  | 2.1–16.0  |          | (3)                  |
|                   | 3.03      | 13.98     |          | (2)                  |
| [g/kg fuel]       |           | 50        |          | (1)                  |
| MOTORCYCLES       |           |           |          |                      |
| [g/km]            | 0.3       |           |          | (1)                  |
| [g/kg fuel]       | 3.0       |           |          | (1)                  |

- (1) OECD - Air Management Policy Group (1984). Note: All emission factors proposed by the OECD were elaborated in cooperation with TNO as average best estimates.  
 (2) Energiebericht und Energie Konzept der österreichischen Bundesregierung, Austria (1984).  
 (3) Handbook of Emission Factors, The Netherlands (1980).

between data from different countries.

The Organisation for Economic Co-operation and Development (OECD) together with the Commission of the European Communities (CEC) is currently preparing a coherent, and therefore as far as possible, comparable emission inventory for major air pollutants, i.e.,  $SO_2$ ,  $NO_x$ , and volatile organic compounds (VOC's), for OECD-Europe. Together with experts from member countries, the OECD developed a set of general guidelines (OECD-Air Management Policy Group, 1984; Lübker, 1987) to allow countries to prepare national inventories in a comparable form. Individual member countries have followed these prescribed guidelines and, in early 1987, national emission inventories were available for 12 countries. These inventories have undergone several rounds of verification to assure their compatibility and results have been described by Lübker and de Tilly (1987a, 1987b).

In general, it can be assumed that these inventories were calculated in a detailed "bottom-up" way, following a set of similar, if not identical, assumptions and definitions, and by making use of the most detailed information available in each country. Major emission sectors - such as those used in RAINS - include the same sub-categories, and are therefore comparable.



#### 4.5. Analysis of the OECD NO<sub>x</sub> Emission Inventory

The OECD inventory was compiled for the base year 1980. The 12 countries included at the time of this study were: Austria, Denmark, Finland, France, FRG, Italy, the Netherlands, Norway, Portugal, Sweden, Switzerland, and the UK.

The total national emissions as reported by countries have been correlated with general statistical parameters, including the number of inhabitants, the surface area per country, the gross national product, and the fossil fuel energy consumed. As was expected, NO<sub>x</sub> emissions are most closely related to total fossil fuel used (coefficient of determination  $r^2 = 0.98$ ,  $t = 36.09$ , therefore, confidence level  $\geq 99.9$  percent), and energy use is, therefore, the best single statistical parameter to correlate NO<sub>x</sub> emissions with.

OECD emission data have also been split into the same source sectors contained in the current RAINS SO<sub>2</sub> emission model: (1) transportation, (2) power plants, (3) domestic, (4) industrial combustion, (5) fuel conversion, and (6) industrial processes. The "fuel conversion" sector in RAINS represents refineries and other fuel conversion plants; the only NO<sub>x</sub> emissions from this sector are from refinery furnaces and such, resulting from stationary, indirect fuel combustion. The mechanism of generation is, thus, almost identical to industrial combustion, and emissions from the conversion sector in the OECD inventory have been included in "industrial combustion" since they cannot be separated for all 12 countries. The two sectors have, therefore, been added together. Industrial process emissions are not calculated in the sulfur emissions submodel since they are not energy-related, but are added as constants if known, or neglected. In the case of nitrogen oxides, process emissions are generally minor (see Table 3). OECD emission data aggregated into the IIASA source sectors are contained in Annex 1. The relative contributions by these sectors to total national NO<sub>x</sub> emissions in the 12 OECD countries are shown in Table 3. The four main, energy-related emission sectors, which are responsible for almost all NO<sub>x</sub> emissions are: (1) transportation, (2) power plants, (3) domestic and (4) industrial combustion.

Table 3. Overall emission contribution by sectors to total NO<sub>x</sub> emissions in OECD-Europe in 1980 (Lübker and de Tilly, 1987b).

| Country        | Contribution to Total National NO <sub>x</sub> Emissions [%] |                 |                        |                          |                         |
|----------------|--------------------------------------------------------------|-----------------|------------------------|--------------------------|-------------------------|
|                | Trans-<br>portation                                          | Power<br>Plants | Domestic<br>Combustion | Industrial<br>Combustion | Industrial<br>Processes |
| AUSTRIA        | 68                                                           | 9               | 5                      | 14                       | 5                       |
| DENMARK        | 33                                                           | 50              | 9                      | 7                        | 0                       |
| FINLAND        | 53                                                           | 37              | -                      | 4                        | 3                       |
| FRANCE         | 54                                                           | 14              | 11                     | 17                       | 3                       |
| GERMANY        | 55                                                           | 26              | 6                      | 9                        | 5                       |
| ITALY          | 61                                                           | 18              | 5                      | 12                       | 3                       |
| NETHERLANDS    | 58                                                           | 15              | 7                      | 8                        | 11                      |
| NORWAY         | 75                                                           | 0               | 4                      | 12                       | 9                       |
| PORTUGAL       | 64                                                           | 12              | -                      | 8                        | 16                      |
| SWEDEN         | 61                                                           | 3               | 15                     | 11                       | 10                      |
| SWITZERLAND    | 70                                                           | 0               | 4                      | 25                       | 0                       |
| UNITED KINGDOM | 37                                                           | 44              | 3                      | 13                       | 0                       |
| AVERAGE        | 53                                                           | 24              | 6                      | 12                       | 4                       |

Until now, only very limited information is available from the countries concerning emissions per fuel type in each sector; on the other hand, internationally comparable information on total energy consumption in 1980 per sector per fuel is available from the OECD IEA statistics.

#### 4.6. Review of SO<sub>2</sub> Emission Calculations in RAINS

The sulfur emission submodel in RAINS calculates SO<sub>2</sub> emissions for each of the above mentioned economic sectors and for each of the following eight fuel types: (1) hard coal, (2) brown coal, (3) derived coal, (4) light fraction of oil (including gasoline), (5) middle distillate (including diesel), (6) heavy oil, (7) gas, and (8) other solid fuels (including wood, etc.). Future emission estimates are therefore dependent on the increased or decreased use of a particular fuel in each sector. If RAINS is to predict the overall impact of sulfur and nitrogen deposition on the environment, it would be advantageous to have emission estimates for both pollutants based on the same statistical variables, i.e., energy forecasts. The advantage of this would be that any mistake in energy predictions would only affect the absolute total emissions and resulting deposition, but would not change the relative contributions to total deposition by the two pollutants.

RAINS contains energy balances for all 27 European countries included in the model; these are based on UN ECE and OECD IEA energy statistics and balances (ECE, 1987; IEA, 1983; IEA, 1984) and form part of the sulfur emission submodel. Generally, the IEA statistics are more detailed, but give data only for OECD-Europe, whereas the ECE statistics give comparable data for all European countries. The total fuel consumption figures are, therefore, taken from the ECE statistics, whereas the allocation to specific coal and oil types (e.g., hard coal, brown coal, and derived coal) is mainly based on IEA information. Unfortunately, the total fuel consumption figures per country are not necessarily identical in the two sets of energy statistics.

In order to use the RAINS energy balances to also predict NO<sub>x</sub> emissions, representative, average emission factors for each of these 32 sector/fuel combinations would have to be found. The NO<sub>x</sub> emission factor ranges from the literature are, however, too broad, and the proper choice would require too detailed information about each sector in each country.

### 5. A METHOD FOR CALCULATING NO<sub>x</sub> EMISSIONS

From the analysis of the OECD NO<sub>x</sub> emission inventory, it can be concluded that this set of data, although small, is appropriate for statistical analysis. The next objective in this study was, therefore, to find an appropriate set of aggregate emission factors by statistical regression analysis for the emission source sectors and different fossil fuels currently used in RAINS.

Nitrogen oxide emissions per sector are the sum of emissions due to the combustion of each of the eight fuels in this sector:

$$e_i = \sum \beta_{ij} E_{ij} \quad [4]$$

in which  $e_i$  : NO<sub>x</sub> emissions per sector,  $i = 1, \dots, 4$ ;  
 $\beta_{ij}$  : emission factor per fuel per sector,  $j = 1, \dots, 8$ ;  
 $E_{ij}$  : energy consumed by each of the eight fuels per sector.

In order to get a set of equations which could be estimated for all  $\beta_{ij}$ , it was assumed that emission factors per fuel per sector are the same in each of the 12 European OECD countries. This results in the following equation:

$$e_{ik} = \sum \beta_{ij} E_{ijk} \quad [5]$$

in which  $k$  is the country index;  $k = 1, \dots, 12$ .

Energy use data  $E_{ijk}$  were taken from the OECD IEA energy statistics for 1980 (IEA, 1987a) and the OECD IEA energy balances for "other solid fuels" (IEA, 1987b), which are not included in the energy statistics; total  $NO_x$  emission rates per sector  $e_{ik}$  were taken from the 1980 OECD emission inventory for  $NO_x$  (Lübker and de Tilly, 1987b). It was decided to estimate the resulting eight unknown parameters by utilizing ordinary least squares (OLS) regression analysis:

$$e_{ik} = \sum \beta_{ij} E_{ijk} + \epsilon_{ij} \quad [6]$$

in which  $\epsilon$  is the disturbance term. Before doing so, the absolute amounts of fuel use per country, which are quite different between countries, were converted into their relative shares; i.e., the matrix of independent variables was normalized in order to avoid heteroscedasticity of the disturbance term.

When attempting to solve for eight unknowns with only a maximum of 12 observations, this usually gives implausible results, such as negative coefficients. Therefore, several techniques were used to increase the degrees of freedom in this set of equations. First, all coefficients were eliminated if their corresponding fuel use in a sector was zero in all countries. This reduced the total number of coefficients from 32 to 26. Since most coefficients could be eliminated in one sector (transportation) and, thus, did not help in increasing the degrees of freedom in other sectors, seemingly unrelated regression (SUR) analysis (Zellner, 1962) was tried. This still resulted in some negative emission coefficients.

In the next step, various combinations of aggregations were carried out in the following way:

1. Different sectors were taken together if emission factor ranges shown in *Table 1* for all fuels in these sectors were very similar. As stated earlier,  $NO_x$  emissions depend on a variety of combustion parameters and are therefore, in some cases, not necessarily sector-specific but rather fuel-related. In doing so, the number of observations was doubled for the same number of coefficients to be estimated.
2. Different fuel types within one sector were aggregated into the same group if emission factor ranges as shown in *Table 1* were identical or very similar for these fuels. This reduced the number of coefficients to be estimated, and it appears justified because some  $NO_x$  emission factors seem to depend more on combustion conditions, such as temperature, etc., than on fuel type.
3. Fuel types were eliminated from regression analysis, also to decrease the number of coefficients to be estimated. This was, however, only done if the remaining fuel mix still covered, on average for all observations, at least 90 percent of all fossil fuel used in the specific sector. In this way, it was tried to keep the error introduced by eliminating explanatory variables from the regression small.

In order to obtain plausible results, the number of independent variables was reduced to a maximum of three if the sample size was 12 or less, and to four if the sample size was larger than 12. In this way, the following sectors or combinations of sectors and

fuel types or groups of fuels were selected for final use in the OLS regression:

1.    Transportation       Light oil (i.e., gasoline) and middle distillate (i.e., diesel) as two individual fuel types. These cover together on average in all 12 countries 99 percent of total fossil fuel consumption by this sector, varying from 97 to 100 percent for individual countries.
2.    Power Plants and Industry as a combined sector:       Middle distillate and heavy oil as one group, hard coal, brown coal, and gas as individual fuels. Together these five fuels cover 92 percent on average of total fossil fuel use in these two sectors in all 12 countries. For power plants alone, these fuels cover on average 99 percent in the 10 countries that have emissions from this sector, ranging from a low of 84 percent in Finland to a high of 100 percent in Denmark and the UK. For industry alone, these fuels cover 86 percent on average in all 12 countries, ranging from as low as 60 percent in Sweden to 96 percent in Denmark.
3.    Domestic:           Hard coal, light oil, and gas as one group, and middle distillate and heavy oil as two individual fuels, together covering on average 95 percent of total fossil fuel use in this sector in the 10 countries that reported emissions. For individual countries, these fuels cover between 73 (Austria) and 100 percent (Netherlands).

The sample size was 12 for "transportation", 22 for "power plants and industry" (i.e., 10 countries for "power plants" and 12 countries for "industry"), and 10 for "domestic". In the case of "power plants", two countries (Norway and Switzerland) do not have any major fossil fuel power plants and, thus, no emissions from this sector. In the case of "domestic", Finland and Portugal have not reported emissions from this sector. Emission data used in the "industry" sector include those originating from combustion in industry and fuel conversion such as refineries.

The constant in the regression analysis was always set to zero, i.e., forcing the regression line through the origin. This was based on the assumption that emissions only occur if some fuel is burned.

It should be noted that "other solid fuels" were not included in any final regression even though their use makes up a relatively significant fraction of total fossil fuels in some sectors and some countries, in particular in the "domestic" and "industry" sectors, particularly in Scandinavia. However, no plausible emission coefficients could be derived, probably reflecting the weakness of the energy data. It should be recalled that "other solid fuel" consumption for each country is taken as the difference between all "solid fuels" per sector in the IEA Energy Balances (1987b) and the sum of all coals per sector in the IEA Energy Statistics (1987a); this may be a significant source of error. Further, emission data included in the OECD inventory for "other solids" are most likely also relatively weak; and only few aggregate emission factors are available from the literature for comparison.

## 6. RESULTS

Emission coefficients as calculated by OLS regression analysis for the fuel types and fuel type aggregations as described above are given per sector in *Table 4*; standard errors and probability values (t-statistics) for each estimated coefficient as well as the

coefficients of determination ( $r^2$ ) for the overall regressions are also presented. Comparison of the calculated coefficients with emission factor ranges of *Table 1* shows that they generally fall within the same ranges; only for gas, the light fraction of oil, and hard coal in the domestic sector, the coefficient tends to be low.

*Table 4.* Aggregate emission coefficients calculated by ordinary least squares (OLS) method.

| Sector<br>[std. error/<br>(t-statistic)] | Fuels                |                       |    |                     |                       |                        |                      |    | $R^2$ |
|------------------------------------------|----------------------|-----------------------|----|---------------------|-----------------------|------------------------|----------------------|----|-------|
|                                          | HC                   | BC                    | DC | GAS                 | LF                    | MD                     | HO                   | OS |       |
| Transportation                           | -                    | -                     | -  | -                   | 455<br>[195]<br>(2.3) | 1580<br>[395]<br>(3.9) | -                    | -  | 0.29  |
| Power Plants<br>and Industry             | 460<br>[65]<br>(7.3) | 195<br>[245]<br>(0.8) | -  | 70<br>[80]<br>(0.9) | -                     | 185<br>[45]<br>(3.9)   | 185<br>[45]<br>(3.9) | -  | 0.57  |
| Domestic                                 | 35<br>[30]<br>(1.2)  | -                     | -  | 35<br>[30]<br>(1.2) | 35<br>[30]<br>(1.2)   | 90<br>[30]<br>(3.0)    | 180<br>[95]<br>(1.9) | -  | 0.30  |

Units: tonnes  $NO_2$ /PJ input.

LEGEND: HC: hard coal  
BC: brown coal  
DC: derived coal  
LF: light fraction of oil (incl. gasoline)  
MD: middle distillate (incl. diesel)  
HO: heavy oil  
OS: other solid fuels (e.g., wood)

As might be anticipated from the small sample of only 12 countries, the statistical parameters indicate that coefficients as calculated in these regressions are not always significant (i.e., the confidence level is less than 95, or even 90 percent). Since many different sector combinations and fuel aggregations were tried, some further indication about the reliability and thus, the robustness of estimated coefficients was obtained. In those cases where coefficients with high probability values were estimated in one particular aggregation, usually very similar coefficients, varying less than  $\pm 5$  percent, were also calculated in other aggregations. Examples are "transportation" coefficients for gasoline and diesel as well as "hard coal" and "heavy oil" emission coefficients for the power plant sector. In the case of "gas", low probability values (below the significance level) were obtained in almost all combinations even though the coefficients themselves were often very similar, i.e., within  $\pm 10$  percent of those reported in *Table 4*. Another such example is the "brown coal" coefficient for power plants.

The estimated coefficients represent emission factors for calculating baseline emissions without add-on control equipment in place; combustion modifications are assumed to have only been carried out to improve efficiency in production but not to reduce atmospheric pollution. These emission coefficients can, thus, be considered "uncontrolled", and the use of any emission reduction technology in future scenarios can be figured into the emission calculations by applying the average reduction percentage achievable in a specific sector, as follows:

$$e_{ij} = \beta_{ij} * E_{ijk} * (1-a). \quad [7]$$

The next step was to check the reasonableness of this method. The calculated  $NO_x$  emission coefficients were entered into an emission factor matrix and incorporated into RAINS; all other emission coefficients were assumed zero. The emission coefficients determined for industrial combustion were used in both RAINS' sectors, "industry" and "conversion". This was appropriate because IIASA's energy balances only include fuels actually consumed in the "conversion" sector.

By multiplication of the  $NO_x$  emission coefficients with IIASA's energy balances for 1980, total  $NO_x$  emissions per sector and per country were calculated for all Europe. The results of these calculations are shown in *Table 5* for total national  $NO_x$  emissions; they are compared to other existing emission inventories such as ECE, EMEP, and PHOXA, and the percentage difference with these inventories is indicated. In many cases (12 out of 27), this difference is smaller than 20 percent, which currently represents the desired accuracy with which modelers would like to have total national  $NO_x$  emissions reported (OECD-Air Management Policy Group, 1986). Eggleston and McInnes (1987) showed that, even with very detailed national background data, the uncertainty for British road traffic  $NO_x$  emissions, for example, is still as high as 40 percent. If compared on a total European scale (USSR excluded), the RAINS calculations overestimate the EMEP total by only four percent. If compared for all 18 European OECD countries, RAINS overpredicts total  $NO_x$  emissions by five percent, and for non-OECD countries (excluding the USSR), the model underpredicts the total by three percent. Large differences are mainly observed between RAINS calculations and ECE or EMEP estimates for non-OECD countries and those countries that show considerable differences between estimates from different inventories. Agreement between RAINS results and PHOXA estimates is within  $\pm 20$  percent for all countries, OECD and non-OECD, with the only exception of GDR.

In order to further check the reasonableness of this approach, national emission rates for each economic sector were analyzed and compared with available inventory data. Unfortunately, the availability of sector-specific emission data is limited. Therefore, such comparison was possible only in the case of emissions from transportation. This sector, however, contributes a major share to total  $NO_x$  emissions, being 53 percent on the average for the 12 OECD countries used in this analysis. *Table 6* shows the results of this comparison. In general, the over- and underestimations are of the same order of magnitude as for total  $NO_x$  emissions when RAINS calculations are compared with OECD and ECE/EMEP figures. In eight cases out of a total of 20, differences are less than  $\pm 20$  percent; in 13 cases, they are less than  $\pm 40$  percent. In some cases, a comparison between different inventories and RAINS calculations shows that differences for the transportation sector are of opposite direction than for total  $NO_x$ ; i.e., total  $NO_x$  for a specific country may have been underestimated whereas transportation emissions for the same country are overestimated, or vice versa.

Since due to the lack of data, no absolute comparison of estimated emissions is possible at a total European level, the relative shares of transportation to total  $NO_x$  emissions as calculated in RAINS are indicated for (a) all Europe, (b) OECD-Europe, and (c) non-OECD Europe. This shows that RAINS calculates on the average 54 percent of total  $NO_x$  in OECD-Europe (for all 18 European OECD countries) as originating from the transportation sector, which is in good agreement with the 53 percent on average for the 12 countries that supplied detailed data. For non-OECD countries (excluding the USSR), RAINS calculates on the average 16 percent as coming from transportation, which confirms common knowledge that the relative share by traffic to overall  $NO_x$  emissions in eastern Europe is significantly lower than in western countries. Relative contributions by traffic as reported to the ECE, range from 5 to 30 percent in eastern Europe.

Table 5. Comparison of national NO<sub>x</sub> emissions (as kt NO<sub>2</sub>) for Europe in 1980.

| Country                       | A<br>RAINS<br>/combustion/ | B<br>ECE/EMEP <sup>1)</sup><br>/combustion/ | C<br>PILLOX A <sup>2)</sup><br>/combustion/ | D<br>OECD <sup>3)</sup><br>/combustion/ | (A-B)/B*100<br>/%/ | (A-C)/C*100<br>/%/ | (A-D)/D*100<br>/%/ |
|-------------------------------|----------------------------|---------------------------------------------|---------------------------------------------|-----------------------------------------|--------------------|--------------------|--------------------|
| ALBANIA                       | 28                         | g <sup>4)</sup>                             |                                             | 211                                     |                    |                    |                    |
| AUSTRIA                       | 186                        | 216                                         |                                             | 14                                      |                    |                    | -10                |
| BELGIUM                       | 426                        | 442                                         | 508                                         | 4                                       |                    | -16                |                    |
| BULGARIA                      | 292                        | 150 <sup>5)</sup>                           |                                             | 95                                      |                    |                    | -19                |
| CSSR                          | 631                        | 1 204                                       | 779                                         | 48                                      |                    |                    |                    |
| DENMARK                       | 270                        | 251                                         | 253                                         | 8                                       |                    |                    | 7                  |
| FINLAND                       | 244                        | 280                                         |                                             | 13                                      |                    |                    |                    |
| FRANCE                        | 1 976                      | 1 867                                       |                                             | 6                                       |                    |                    |                    |
| FRG                           | 2 668                      | 3 100                                       | 3 117                                       | 13                                      |                    | -14                | -9                 |
| GDR                           | 520                        | 800                                         | 1 222                                       | 35                                      |                    | -57                |                    |
| GREECE                        | 218                        | 127 <sup>4)</sup>                           |                                             | 72                                      |                    |                    |                    |
| HUNGARY                       | 220                        | 270 <sup>5)</sup>                           | 199                                         | 19                                      |                    | 11                 |                    |
| IRELAND                       | 84                         | 67                                          | 99                                          | 25                                      |                    | -15                |                    |
| ITALY                         | 1 452                      | 1 410                                       |                                             | 3                                       |                    |                    | 1                  |
| LUXEMBURG                     | 40                         | 23                                          | 35                                          | 74                                      |                    | 14                 |                    |
| NETHERLANDS                   | 494                        | 535                                         | 465                                         | 8                                       |                    | 6                  |                    |
| NORWAY                        | 173                        | 215 <sup>6)</sup>                           |                                             | 20                                      |                    |                    | 63                 |
| POLAND                        | 1 484                      | 840 <sup>6)</sup>                           | 1 705                                       | 77                                      |                    | -13                | -10                |
| PORTUGAL                      | 149                        | 166                                         |                                             | 10                                      |                    |                    |                    |
| ROMANIA                       | 383                        | 390 <sup>4)</sup>                           |                                             | 2                                       |                    |                    |                    |
| SPAIN                         | 951                        | 780 <sup>6)</sup>                           |                                             | 22                                      |                    |                    |                    |
| SWEDEN                        | 297                        | 328                                         | 300                                         | 22                                      |                    |                    | -1                 |
| SWITZERLAND                   | 161                        | 196                                         |                                             | 18                                      |                    |                    | -17                |
| TURKEY                        | 356                        | 175 <sup>4)</sup>                           |                                             | 103                                     |                    |                    |                    |
| UK                            | 2 454                      | 1 916                                       | 2 642                                       | 28                                      | 1 867              | -7                 | 31                 |
| USSR <sup>7)</sup>            | 9 082                      | 2 790                                       |                                             | 226                                     |                    |                    |                    |
| YUGOSLAVIA                    | 339                        | 190 <sup>4)</sup>                           |                                             | 78                                      |                    |                    |                    |
| EUROPE                        | 25 598                     | 18 737                                      |                                             | 37                                      |                    |                    |                    |
| EUROPE excl. USSR             | 16 516                     | 15 947                                      |                                             | 4                                       |                    |                    |                    |
| OECD-EUROPE                   | 12 958                     | 12 284                                      |                                             | 5                                       |                    |                    |                    |
| NON OECD-EUROPE               | 12 640                     | 6 453                                       |                                             | 96                                      |                    |                    |                    |
| NON OECD-EUROPE<br>excl. USSR | 3 558                      | 3 663                                       |                                             | 3                                       |                    |                    |                    |

Notes and references:

- 1) United Nations, 1987.
- 2) Bakum and Veldt, 1986b.
- 3) Lubkert and de Tilly, 1987b.
- 4) Semb and Amble, 1981.
- 5) Year of the inventory is 1984.
- 6) Year of the inventory is 1985.
- 7) Two-thirds of total energy are assumed to be used in the European part of the USSR.

Table 6. Comparison of national  $NO_x$  emissions from the transportation sector (as kt  $NO_2$ ) for Europe in 1980.

| Country                    | A<br>RAINS | B<br>ECE/EMEP <sup>1)</sup> | C<br>PHOXA <sup>2)</sup> | D<br>OECD <sup>3)</sup> | (A-B)/B*100<br>[%] | (A-C)/C*100<br>[%] | (A-D)/D*100<br>[%] |
|----------------------------|------------|-----------------------------|--------------------------|-------------------------|--------------------|--------------------|--------------------|
| ALBANIA                    | 9          |                             |                          |                         |                    |                    |                    |
| AUSTRIA                    | 131        | 146                         |                          | 146                     | -10                |                    | -10                |
| BELGIUM                    | 210        | 242                         | 182                      |                         | -13                | 15                 |                    |
| BULGARIA                   | 89         |                             |                          |                         |                    |                    |                    |
| CSSR                       | 91         | 60                          | 132                      |                         | 52                 | -31                |                    |
| DENMARK                    | 109        | 75                          | 83                       | 81                      | 45                 | 31                 | 35                 |
| FINLAND                    | 115        | 150                         |                          | 152                     | -23                |                    | -24                |
| FRANCE                     | 1 105      | 1 021                       |                          | 1 056                   | 8                  |                    | 5                  |
| FRG                        | 1 313      | 1 748                       | 1 465                    | 1 693                   | -25                | -10                | -22                |
| GDR                        | 57         |                             | 232                      |                         |                    | -75                |                    |
| GREECE                     | 144        | 91                          |                          |                         | 58                 |                    |                    |
| HUNGARY                    | 85         | 81 <sup>4)</sup>            | 44                       |                         | 5                  | 93                 |                    |
| IRELAND                    | 54         | 21                          | 54                       |                         | 157                | 0                  |                    |
| ITALY                      | 893        | 679                         |                          | 949                     | 32                 |                    | -6                 |
| LUXEMBURG                  | 16         | 14                          | 15                       |                         | 14                 | 7                  |                    |
| NETHERLANDS                | 299        | 321                         | 246                      | 298                     | 5                  | 22                 | 0                  |
| NORWAY                     | 137        | 185 <sup>5)</sup>           |                          | 86                      | -26                |                    | 59                 |
| POLAND                     | 175        |                             | 290                      |                         |                    | -40                |                    |
| PORTUGAL                   | 107        | 106                         |                          | 106                     | 1                  |                    | 1                  |
| ROMANIA                    | 71         |                             |                          |                         |                    |                    |                    |
| SPAIN                      | 610        | 340                         |                          |                         | 79                 |                    |                    |
| SWEDEN                     | 185        | 199                         |                          | 201                     | -7                 |                    | -8                 |
| SWITZERLAND                | 108        | 138                         |                          | 136                     | -22                |                    | -21                |
| TURKEY                     | 252        |                             |                          |                         |                    |                    |                    |
| UK                         | 1 015      | 690                         | 978                      | 703                     | 47                 | 4                  | 44                 |
| USSR <sup>6)</sup>         | 4 613      | 698                         |                          |                         | 561                |                    |                    |
| YUGOSLAVIA                 | 149        |                             |                          |                         |                    |                    |                    |
| EUROPE                     | 12 142     | 47% of total $NO_x$         |                          |                         |                    |                    |                    |
| EUROPE excl. USSR          | 7 529      | 46% of total $NO_x$         |                          |                         |                    |                    |                    |
| OECD-EUROPE                | 6 952      | 54% of total $NO_x$         |                          |                         |                    |                    |                    |
| NON OECD-EUROPE            | 5 190      | 41% of total $NO_x$         |                          |                         |                    |                    |                    |
| NON OECD-EUROPE excl. USSR | 577        | 16% of total $NO_x$         |                          |                         |                    |                    |                    |

Notes and references:

- 1) United Nations, 1987.
- 2) Bakkum and Veldt, 1986b.
- 3) Lübker and de Tilly, 1987b.
- 4) Year of the inventory is 1984.
- 5) Year of the inventory is 1985.
- 6) Two thirds of total energy are assumed to be used in the European part of the USSR.



## 7. DISCUSSION

Some of the differences in total  $NO_x$  emission rates between RAINS and those reported by countries can be explained by the fact that some fuel used in individual countries is neglected because no  $NO_x$  emission coefficients for these fuels are included in RAINS. No coefficients were estimated for these fuels because their share was minor in the set of countries used in the regression analysis. Using the RAINS energy balances, relative shares of fuels in each sector have been calculated for all 27 European countries and are presented in Table 7. Adding the shares for which emission coefficients have been determined shows that only one percent of total fossil fuels remains unaccounted for in the transportation and power plant sectors, whereas 20 and 23 percent remain unaccounted for in all Europe in industrial and domestic combustion. In individual countries, these percentages may be significantly different.

Table 7. Relative fuel consumption by sector and fuel type in all Europe in 1980[%] (RAINS, 1987).

| Sector                         | Fuels: |    |    |     |    |    |    |    |
|--------------------------------|--------|----|----|-----|----|----|----|----|
|                                | HC     | BC | DC | GAS | LF | MD | HO | OS |
| Transportation                 | 1      | 0  | 0  | 0   | 51 | 48 | 0  | 0  |
| Power Plants                   | 35     | 19 | 0  | 21  | 0  | 0  | 24 | 1  |
| Domestic                       | 7      | 6  | 6  | 26  | 0  | 40 | 4  | 11 |
| Industry<br>(incl. conversion) | 14     | 2  | 15 | 25  | 4  | 14 | 25 | 1  |

LEGEND: HC: hard coal  
 BC: brown coal  
 DC: derived coal  
 LF: light fraction of oil (incl. gasoline)  
 MD: middle destillate (incl. diesel)  
 HO: heavy oil  
 OS: other solid fuels (e.g., wood)

The statistical analysis could theoretically be improved either by increasing the number of observations or by reducing the number of coefficients to be estimated. The number of observations could be increased once data from additional countries become available, [e.g., from continuing work in the international air pollution study CORINAIR (Bouscaren *et al.*, 1987)], or if time series for one or more countries were available. The latter would, however, only be of value if calculations of the time series were independent of each other to avoid co-linearity. Many techniques have been tried in this study to reduce the number of coefficients to be estimated, and it appears that any further aggregation of emission coefficients would contradict current engineering knowledge about how  $NO_x$  emission factors vary between sectors and fuels.

An attempt has been made in this study to assess the reliability of calculated emission coefficients. Reported coefficients of determination for each regression and standard errors for each estimated coefficient give an indication of the uncertainty of the coefficients and of the quality of the energy data and emission rates used in the analysis. Several additional sources of uncertainty in estimating total emission rates based on calculation by this method can, however, be identified: (1) energy data from international statistics used in RAINS have certain errors associated with them; (2) fuel use patterns may be different in countries other than those used in the regression analysis; (3) fuel use patterns may change over time; and (4) emission factors may change over time indepen-

dent of add-on emission control technology. The latter two points, therefore, introduce errors into any extrapolation of data in time, i.e., in forecasting. Point (2) introduces errors into geographical extrapolation of data because (1) some significant amount of fuel may be used in a particular sector by a specific country for which no emission coefficient was determined in the regression analysis; and (2) a weak emission coefficient such as for brown coal in "power plants" may introduce a larger mistake in eastern European countries than in OECD Europe because the amount of brown coal used in most European OECD countries in power stations is relatively small (less than 3 percent in 10 out of the 12 countries used in this study) compared to Eastern Europe (on the average for all Europe 19 percent).

Point (1) introduces errors because there are, in fact, several energy estimates available, and the sensitivity of  $NO_x$  calculations to these different energy estimates should be investigated. The uncertainty of computed country  $NO_x$  emissions can also be estimated by replacing deterministic emission factors with their frequency distributions. The frequency distributions for the emission coefficients can be derived from (1) the errors of the regression analysis used to compute these coefficients, or (2) the range of literature values.

Lastly, differences in the underlying statistics of various countries could be compared and might be explained; examples are different automobile fleet composition and average speeds, or different burner types in power plants, etc. If such distinct differences can be identified, it might be possible to group countries into classes with similar background statistics, e.g., northwestern Europe, eastern Europe, and southern Europe, and the average emission coefficient currently used in RAINS could be scaled up or down by adding a weighting factor to the equation. For countries in which no specific information is available, weighting factors could then be selected for the fuel- and sector-specific average emission factors just by knowing the group to which the particular country belongs.

## 8. SUMMARY AND CONCLUSIONS

Analysis of the principles of  $NO_x$  formation has shown that an attempt to develop an  $NO_x$  emission model for RAINS based on these principles is currently infeasible because of the large number of variables upon which nitrogen oxide formation is dependent and the lack of knowledge about these variables. Even if the necessary detail were available for one particular year, information about the variation over time of these variables is available in neither international nor in national statistics; therefore prediction of  $NO_x$  emissions is infeasible.

To date, only few  $NO_x$  emission inventory data are available for Europe that are comparable with each other. The OECD together with the European Communities are currently developing such a coherent and verified inventory for 1980 in which countries have estimated their total national  $SO_2$ ,  $NO_x$  and VOC emissions for a relatively large number of emission source categories according to a prescribed method. This has shown that the following emission source categories, identical to the ones currently used in the sulfur emissions submodel of RAINS, are also the most relevant to  $NO_x$  emission estimations: (1) transportation, (2) power plants, (3) domestic and (4) industrial combustion.

Analysis of the OECD emission data has further shown that overall  $NO_x$  emissions are most closely correlated to fossil fuel consumption rates, and aggregate emission factors available from the literature also show that  $NO_x$  emissions are fuel-specific. Energy consumption is, therefore, the only statistical parameter that is applicable to calculate emissions in all four sectors. Furthermore, energy predictions are readily available for Europe in a consistent fashion. Basing  $NO_x$  estimations on energy balances and forecasts also has the advantage that identical statistical input information is used for sulfur and nitrogen emission calculations; this is important if RAINS is to predict the overall impact of sulfur

and nitrogen deposition on the environment because, even if the absolute emission and deposition values are wrong due to wrong energy forecasts, the relative contributions by the two pollutants would remain unchanged.

In this study, the OECD data, currently available for 12 countries, were used in multiple OLS regression analysis to determine average, aggregate, sector- and fuel-specific emission coefficients for those sector/fuel combinations already used in RAINS for sulfur calculations that are most important in calculating total national  $NO_x$ . Estimation of such coefficients via regression analysis has the advantage over the use of average emission factors from the literature in that regression coefficients utilize best knowledge of 1980 emission levels.

Emission coefficients determined in this way were used by RAINS to calculate baseline (i.e., uncontrolled)  $NO_x$  emissions for all Europe in 1980. Comparison of the results with reported national totals as well as reported traffic  $NO_x$  shows generally good agreement, i.e., within 20 percent. If  $NO_x$  emissions calculated by RAINS are compared to EMEP estimates on a total European scale (USSR excluded), RAINS overestimates the EMEP total by only four percent. Relative contributions as calculated by RAINS for the transportation sector also reflect numbers reported by countries, being on the average 54 percent of total  $NO_x$  for OECD Europe and 16 percent for non-OECD Europe (excluding the USSR). For several countries and sectors, differences between calculated and reported  $NO_x$  emissions are, however, still larger than 20 percent and, therefore, some suggestions have been made as to how to improve the method. These include uncertainty analysis of emission coefficients and energy statistics and a more detailed analysis of the underlying statistics of various countries.

This paper describes how a "top-down" approach was used to extrapolate a consistent set of data from one set of countries to other countries and other years. The  $NO_x$  emission model derived in this way also allows for the testing of different future scenarios with and without emission reduction technology.

## REFERENCES

- Alcamo, J., M. Amann, J.-P. Hettelingh, M. Holmberg, L. Hordijk, J. Kämäri, L. Kauppi, P. Kauppi, G. Kornai, and A. Mäkelä (1987). Acidification in Europe: a simulation model for evaluating control strategies, *Ambio*, **16**(5), 232-244.
- Amann, M. and G. Kornai (1987). *Cost Functions for Controlling  $SO_2$  Emissions in Europe*. Working Paper WP-87-65, International Institute for Applied Systems Analysis, Laxenburg, Austria.
- Bakkum, A. and C. Veldt (1986a). *PHOXA -- Statistical Data and Emission Factors*. Discussion paper presented at the German-OECD Workshop on Comparison of Emission Inventory Data, held at Schauinsland, Black Forest, F.R. of Germany, on October 22-24, 1986.
- Bakkum, A. and C. Veldt (1986b). *PHOXA Emission Inventory - Tabulated Data and Comparison*. Presented at the German OECD Workshop on Comparison of Emission Inventory Data held at Schauinsland, Black Forest, F.R. of Germany, on October 22-24, 1986.
- Bouscaren, R., C. Veldt, and K.-H. Zierock (1986). *CORINE: Emission Inventory Feasibility Study -- Final Report*. CITEPA, Paris, France.

- Bouscaren, R., C. Veldt, K.-H. Zierock, and B. Lübkert (1987). *CORINAIR Project, Programme CORINE*. National Experts Meeting held in Brussels, Belgium, May 7-8, 1987.
- Economic Commission for Europe (ECE) (1987). *ECE Energy Data Base* (on tape). Geneva, Switzerland.
- Eggleston, H.S. and G. McInnes (1987). *Accuracy of Emission Inventories*. Presented at the EURASAP Workshop on Methodologies for Air Pollutant Emission Inventories held in Paris, June 29-July 2, 1987.
- Eliassen, A. and J. Saltbones (1983). Modeling of long range transport of sulfur over Europe: a two-year model run and some model experiments, *Atmospheric Environment*, **17**, 1457-1473.
- Energiebericht und Energiekonzept der Österreichischen Bundesregierung (1984). Bundesministerium für Handel, Gewerbe und Industrie -- Sektion V (Energie) Wien, Austria
- Handbook of Emission Factors -- Part I: Non Industrial Sources (1980). Ministry of Health and Environmental Protection, Government Publishing Office, The Hague, The Netherlands.
- Holmberg, M., J. Johnson, and L. Maxe (1987). *Assessing Aquifer Sensitivity to Acid Deposition*. Presented at the International Conference on the Vulnerability of Soil and Groundwater Pollutants, Noordwijk, The Netherlands, March 30-April 3, 1987.
- Hordijk, L. (1986). Acid rain abatement strategies in Europe. In T. Schneider (ed), *Acidification and Its Policy Implications*. Elsevier Science Publishers B.V., Amsterdam, The Netherlands, pp. 295-305.
- International Energy Agency (IEA) (1983). *Energy Statistics 1971/1981*. Organisation for Economic Co-operation and Development, Paris, France.
- International Energy Agency (IEA) (1984). *Energy Balances of OECD Countries 1970/1982*. Organisation for Economic Co-operation and Development, Paris, France.
- International Energy Agency (IEA) (1987a). *Energy Statistics 1970/1985 and Main Series from 1960, Volumes 1 and 2*. Organisation for Economic Co-operation and Development, Paris, France.
- International Energy Agency (IEA) (1987b). *Energy Balances of OECD Countries 1970, 1985 and Main Series from 1960*. Organisation for Economic Co-operation and Development, Paris, France.
- Kämäri, J. and M. Posch (1987). Regional application of a simple lake acidification model to Northern Europe. In M.B. Beck (ed), *Systems Analysis in Water Quality Management*. Pergamon Press, pp. 73-84.
- Kauppi, P., J. Kämäri, M. Posch, L. Kauppi, and E. Matzner (1986). Acidification of forest soils, *Ecological Modelling*, **33**, 231-253.
- Löblich, H.-J. (1985). *OECD -- Europa Energieverbrauch und Emissionen von SO<sub>2</sub> and NO<sub>x</sub> 1960 bis 1983*. Umweltforschungsplan des Bundesministers des Innern, Luftreinhalteung, im Auftrag des Umweltbundesamtes (unpublished material).
- Lübkert, B. (1987). *Summary of Current Knowledge and Gaps: OECD Summary -- Experience from the MAP Project*. Contribution to the Opening Session of the EURASAP Workshop on Methodologies for Air Pollutant Emission Inventories, held in Paris, June 29-July 2, 1987.
- Lübkert, B. and S. de Tilly (1987a). *An Emission Inventory for SO<sub>2</sub>, NO<sub>x</sub> and VOC's in North-Western Europe*. Presented at the 80th APCA Annual Meeting and Exhibition held in New York City, June 21-26, 1987.

- Lübkert, B. and S. de Tilly (1987b). *The OECD-MAP Emission Inventory for SO<sub>2</sub>, NO<sub>x</sub> and VOC's in Western Europe*. Presented at the EURASAP Workshop on Methodologies for Air Pollutant Emission Inventories held in Paris, June 29-July 2, 1987.
- Mäkelä, A., J. Materna, and W. Schöpp (1987). *Direct Effects of Sulfur on Forests in Europe -- A Regional Model of Risk*. Working Paper WP-87-57, International Institute for Applied Systems Analysis, Laxenburg, Austria.
- Mason, H.B., and M.A. Herther (1982). Emission inventory for stationary NO<sub>x</sub> sources (USA). In T. Schneider and L. Grant (eds), *Air Pollution by Nitrogen Oxides*. Elsevier Scientific Publ. Co., Amsterdam, The Netherlands, pp.131-141.
- Morrison, G.F. (1980). *Nitrogen Oxides from Coal Combustion - Abatement and Control*. Report No. ICHS/TR 11, IEA Coal Research, London, UK.
- OECD-Air Management Policy Group (1984). Summary Record of the Workshop on Emission Inventories in Apeldoorn, 27-28 September, 1984, ENV/AIR/M/84.10, Paris, France (restricted report).
- OECD-Air Management Policy Group (1986). Summary Record from the German-OECD Workshop on Comparison of Emission Inventory Data. ENV/AIR/86.9, Paris, France (restricted report).
- OECD (1987). Control of major air pollutants - status report, *OECD Environment Monograph*, 10, OECD, Paris, France.
- Rentz, O. (1984). Minderung von Stickoxidemissionen aus Kohlekraftwerken in Baden-Württemberg, Bericht der Kommission, Staatsministerium in Baden-Württemberg, Stuttgart, F.R. of Germany.
- Semb, A. and E. Amble (1981). *Emission of Nitrogen Oxides from Fossil Fuel Combustion in Europe*. ISBN-82-7247-277-5, Norwegian Institute for Air Research, Lillestrøm, Norway.
- Schneider, T. and L. Grant (eds) (1982). *Air Pollution by Nitrogen Oxides*. Elsevier Scientific Publ. Co., Amsterdam, The Netherlands.
- Thompson, R.E. (1979). *Nitric Oxide Controls for Coal-Fired Utility Boilers from an Applications Viewpoint*. In Proceedings of the Second NO<sub>x</sub> Control Technology Seminar, Denver, Colorado, 8-9 November 1978.
- United Nations (UN) (1987). National strategies and policies for air pollution abatement. Convention on Long-Range Transboundary Air Pollution, Report ECE/EB.AIR/14, New York, USA.
- US Department of Energy (1983). NO<sub>x</sub> Information Book. DOE/NBB-0044 UC Dist. 11, 90, Washington D.C.
- Zeldovich, B.Y. (1946). *Acta Physico Chemica USSR* 21, Nr. 4.
- Zellner, A. (1962). An efficient method of estimating seemingly unrelated regressions and tests of aggregation bias, *J. American Statistical Association*, 57, 348-368.

## ANNEX 1. NO<sub>x</sub> EMISSIONS IN OECD EUROPE

NO<sub>x</sub> emissions per RAINS source category in 1980 [in kt NO<sub>2</sub>] (OECD MAP– inventory)

| Source Sector                           | A<br>U<br>T | D<br>N<br>K | F<br>I<br>N | F<br>R<br>A | G<br>E<br>R | I<br>T<br>A | N<br>L<br>D | N<br>O<br>R | P<br>O<br>R | S<br>W<br>E | S<br>W<br>I | U<br>K<br>D | S<br>U<br>M |
|-----------------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Transportation                          | 146.0       | 81.3        | 151.6       | 1 055.7     | 1 692.8     | 948.8       | 298.1       | 87.6        | 106.3       | 201.0       | 136.0       | 703.0       | 5 608.0     |
| Powerplants                             | 20.0        | 121.1       | 106.3       | 280.0       | 803.0       | 287.0       | 79.1        | 0.0         | 19.7        | 10.0        | 0.0         | 851.0       | 2 577.3     |
| Domestic combustion                     | 10.0        | 22.4        | –           | 213.0       | 191.6       | 79.3        | 36.3        | 4.7         | –           | 48.0        | 8.6         | 65.0        | 678.8       |
| Industrial<br>combustion <sup>(1)</sup> | 30.0        | 16.0        | 11.2        | 334.4       | 266.1       | 190.4       | 42.3        | 13.4        | 13.1        | 35.8        | 49.4        | 258.0       | 1 260.2     |
| Industrial<br>processes                 | 10.0        | 1.0         | 9.2         | 61.9        | 140.2       | 50.0        | 58.0        | 10.6        | 26.6        | 33.7        | –           | –           | 401.2       |
| Other                                   | 0.0         | 1.6         | 5.2         | 5.1         | 0.0         | 0.0         | 2.9         | 0.1         | 0.0         | 0.0         | 0.0         | 47.0        | 61.9        |
| Total <sup>(2)</sup>                    | 216.0       | 243.4       | 283.6       | 1 950.1     | 3 093.6     | 1 555.5     | 516.6       | 116.4       | 165.6       | 328.5       | 194.0       | 1 924.0     | 10 587.4    |

Notes:

(1) Including all indirect process heating (e.g. boilers, furnaces, etc.).

(2) Totals may be incomplete due to unavailable data for certain source categories.