

Paris Agreement's aim of 1.5°C warming may result in many possible climates

– SUPPLEMENTARY INFORMATION

Sonia I. Seneviratne¹, Joeri Rogelj^{1,2,4}, Roland Séférian³, Richard Wartenburger¹, Myles R. Allen⁴, Michelle Cain⁴, Richard J. Millar⁴, Kristie L. Ebi⁵, Neville Ellis⁶, Ove Hoegh-Guldberg⁷, Antony J. Payne⁸, Carl-Friedrich Schleussner^{9,10,11}, Petra Tschakert⁶, and Rachel F. Warren¹²

1. Institute for Atmospheric and Climate Science, ETH Zurich, Zurich, Switzerland
2. International Institute for Applied Systems Analysis (IIASA), Laxenburg A-2361, Austria
3. Centre National de Recherches Météorologiques, Météo-France/CNRS, Toulouse, France
4. Environmental Change Institute, School of Geography and the Environment, University of Oxford, Oxford, UK
5. University of Washington, Department of Global Health, Seattle, USA
6. University of Western Australia, UWA School of Agriculture and Environment, Perth, Australia
7. Global Change Institute, University of Queensland, Brisbane, Australia
8. University of Bristol, Bristol, UK
9. Climate Analytics, Berlin, Germany
10. IRITHESys, Humboldt University, Berlin, Germany
11. Potsdam Institute for Climate Impact Research, Potsdam, Germany
12. Tyndall Centre for Climate Change, School of Environmental Sciences, University of East Anglia, Norwich, UK

Supplementary Text 1: Projections of regional changes in climate extremes

The computation of regional climate extreme indices in Table 1 and Figs. 2-3 follows the approach of Seneviratne et al (2016) and Wartenburger et al. (2017). This section provides additional background information compared to Box 1. The considered climate extremes indicators include the warming of the minimum annual night-time temperature (TNn) in the Arctic land [$^{\circ}\text{C}$], the warming of the maximum annual day-time temperature (TXx) in the contiguous United States [$^{\circ}\text{C}$], TXx warming in Central Brazil [$^{\circ}\text{C}$], (soil moisture) drying in the Mediterranean region [in units of standard deviations of pre-industrial climate variability], and increases in heavy precipitation events based on annual maximum consecutive 5-day precipitation (Rx5day) in Southern Asia [%]). The definition of the geographical domains is provided in Seneviratne et al. (2016). The estimates are derived from CMIP5 simulations for 26 climate models and all four Representative Concentration Pathways (RCP) scenarios, i.e. RCP2.6, RCP4.0, RCP6.0 and RCP8.5 (see Suppl. Table S3 for list of analysed simulations).

Supplementary Table S1: Same as Table 1, but for scenarios considered compatible with 1.5°C and 2°C warming in the 5th assessment report of IPCC (Clarke et al. 2014, Rogelj et al. 2015), including projections of changes in regional climate associated with resulting global temperature levels derived following previous studies (Seneviratne et al. 2016, Wartenburger et al. 2017)

		SCEN_1p5C Emissions pathways currently considered in line with keeping warming below 1.5°C in 2100 with 66% chance (allowing for a higher peak in temperature earlier)	SCEN_2C Emissions pathways currently considered in line with keeping warming below 2°C during the entire 21 st century with 66% chance		
		“probable” (66 th percentile) outcome ^a	“worst-case” 10% (90 th percentile) outcome ^b	“probable” (66 th percentile) outcome ^a	“worst-case” 10% (90 th percentile) outcome ^b
General characteristics of pathway	Overshoot 1.5°C in 21 st century with >50% likelihood ^c	Yes (8/8)	Yes (8/8)	Yes (60/60)	Yes (60/60)
	Overshoot 2°C in 21 st century with >50% likelihood	No (0/8)	Yes (4/8)	No (0/60)	Yes (60/60)
	Cumulative CO ₂ emissions up to peak warming (relative to 2016) ^d	510 (490, 560)	470 (410, 520)	930 (790, 1050)	900 (750, 1040)
	Cumulative CO ₂ emissions up to 2100 (relative to 2016) ^d [GtCO ₂]	-40 (-100, 10)		850 (520, 1000)	
	Global GHG emissions in 2030 ^d [GtCO ₂ y ⁻¹]	19 (17, 21)		28 (23, 32)	
	Years of global net zero CO ₂ emissions ^d	2061 (2061, 2063)		2084 (2079, 2086)	
Possible climate range at peak warming (reg+glob)	Global mean temperature anomaly at peak warming [°C]	1.65°C (1.61, 1.68°C)	2.01°C (1.95, 2.03°C)	1.90°C (1.86, 1.95°C)	2.35°C (2.29, 2.48°C)
	Warming in the Arctic ^e (TNn ^f) [°C]	4.75 °C (4.09, 5.44)	5.90 °C (4.97, 6.85)	5.63 °C (4.68, 6.59)	6.97 °C (6.13, 8.38)
	Warming in the contiguous United States ^e (TXx ^f) [°C]	2.39 °C (1.90, 2.84)	2.97 °C (2.36, 3.40)	2.77 °C (2.20, 3.30)	3.51 °C (3.05, 4.11)
	Warming in Central Brazil ^e (TXx ^f) [°C]	2.55 °C (2.12, 2.97)	3.12 °C (2.66, 3.76)	2.96 °C (2.58, 3.55)	3.66 °C (3.31, 4.21)
	Drying in the Mediterranean region ^e [std ^g] (-1: dry; -2: severely dry; -3: very severely dry)	-1.00 (-2.12, -0.39)	-1.25 (-2.21, -0.51)	-1.11 (-2.18, -0.51)	-1.36 (-2.93, -0.69)
	Increase in heavy precipitation events ^f in Southern Asia ^e [%]	9.78 % (6.52, 13.63)	11.56 % (7.04, 18.50)	10.27 % (6.50, 17.40)	16.74 % (9.60, 23.44)
Possible climate range in 2100 (reg+glob)	Global mean temperature warming in 2100 [°C]	1.41°C (1.39—1.43°C)	1.84°C (1.81—1.90°C)	1.84°C (1.76—1.89°C)	2.30°C (2.21—2.46°C)
	Warming in the Arctic ^e (TNn ^f) [°C]	4.07 °C (3.53, 4.72)	5.37 °C (4.60, 6.40)	5.37 °C (4.46, 6.38)	6.86 °C (5.83, 8.24)
	Warming in the contiguous United States ^e (TXx ^f) [°C]	2.00 °C (1.60, 2.48)	2.62 °C (2.17, 3.23)	2.62 °C (2.07, 3.24)	3.37 °C (2.88, 4.03)
	Warming in Central Brazil ^e (TXx ^f) [°C]	2.20 °C (1.95, 2.58)	2.88 °C (2.46, 3.48)	2.88 °C (2.43, 3.47)	3.57 °C (3.20, 4.18)
	Drying in the Mediterranean region ^e [std ^g]	-0.96 (-1.89, -0.28)	-1.24 (-2.27, -0.45)	-1.24 (-2.42, -0.46)	-1.38 (-2.79, -0.67)
	Increase in heavy precipitation events ^f in Southern Asia ^e [%]	7.83 % (4.20, 12.00)	10.19 % (6.34, 16.67)	10.19 % (6.46, 16.64)	15.47 % (8.72, 22.98)

^a 66th percentile estimates for global temperature (i.e. 66% likelihood of being at or below values)

^b 90th percentile estimates for global temperature (i.e. 10% likelihood of being at or above values)

^c All 1.5°C scenarios from AR5 include a substantial probability of overshooting above 1.5°C global warming before returning to 1.5°C

^d The values indicate the median and the interquartile range in parenthesis (25th percentile and 75th percentile)

^e The regional projections in these rows provide the range [median (q25, q75)] associated with the *median* global temperature outcomes of the considered mitigation scenarios at *peak warming* (see Box 1 and Suppl. Text 1 for details).

^f TNn: annual minimum night-time temperature; TXx: annual maximum day-time temperature; std: drying of soil moisture expressed in units of standard deviations of pre-industrial climate (1861-1880) variability; Rx5day: annual maximum consecutive 5-day precipitation (see Suppl. Text 1 for details)

^g Same as footnote e, but for the regional responses associated with the *median* global temperature outcomes of the considered mitigation scenarios by 2100 (see Box 1 and Suppl. Text 1 for details).

Supplementary Table S2: Same as Table 1, but with inclusion of 50% (median) values for considered emissions scenarios (see Box 1 for details).

		SCEN_1p5C Emissions pathways currently considered in line with keeping warming below 1.5°C in 2100 with 66% chance (allowing for a higher peak in temperature earlier)			SCEN_2C Emissions pathways currently considered in line with keeping warming below 2°C during the entire 21 st century with 66% chance		
		“as likely as not” (50 th percentile) outcome ^a	“probable” (66 th percentile) outcome ^b	“worst-case” 10% (90 th percentile) outcome ^c	“as likely as not” (50 th percentile) outcome ^a	“probable” (66 th percentile) outcome ^b	“worst-case” 10% (90 th percentile) outcome ^c
General characteristics of pathway	Overshoot 1.5°C in 21 st century with >50% likelihood ^d	Yes (13/13)	Yes (13/13)	Yes (13/13)	Yes (10/10)	Yes (10/10)	Yes (10/10)
	Overshoot 2°C in 21 st century with >50% likelihood	No (0/13)	No (0/13)	Yes (10/13)	No (0/10)	No (0/10)	Yes (10/10)
	Cumulative CO ₂ emissions up to peak warming (relative to 2016) ^e	710 (630,770)	720 (650, 750)	690 (650, 710)	1050 (1000, 1110)	1050(1020, 1140)	1040 (930, 1140)
	Cumulative CO ₂ emissions up to 2100 (relative to 2016) ^e [GtCO ₂]	320 (200, 340)			1030 (910, 1140)		
	Global GHG emissions in 2030 ^e [GtCO ₂ y ⁻¹]	22 (19, 31)			28 (24, 30)		
	Years of global net zero CO ₂ emissions ^e	2070 (2067, 2074)			2088 (2085, 2092)		
Possible climate range at peak warming (reg+glob)	Global mean temperature anomaly at peak warming [°C]	1.6°C (1.52-1.66)	1.75°C (1.65-1.81)	2.13°C (2.0- 2.2)	1.74°C (1.73-1.76)	1.93°C (1.9-1.94°C)	2.44°C (2.43-2.46°C)
	Warming in the Arctic ^f (TNn ^g) [°C]	4.66°C (4.05, 5.25)	4.75 °C (4.09, 5.44)	5.90 °C (4.97, 6.85)	5.01°C (4.45, 5.58)	5.63 °C (4.68, 6.59)	6.97 °C (6.13, 8.38)
	Warming in the contiguous United States ^f (TXx ^g) [°C]	2.28°C (1.91, 2.67)	2.39 °C (1.90, 2.84)	2.97 °C (2.36, 3.40)	2.53°C (2.03, 2.95)	2.77 °C (2.20, 3.30)	3.51 °C (3.05, 4.11)
	Warming in Central Brazil ^f (TXx ^g) [°C]	2.49°C (2.12, 2.86)	2.55 °C (2.12, 2.97)	3.12 °C (2.66, 3.76)	2.72°C (2.33, 3.19)	2.96 °C (2.58, 3.55)	3.66 °C (3.31, 4.21)
	Drying in the Mediterranean region ^f [std ^g] (-1: dry; -2: severely dry; -3: very severely dry)	-0.94 (-2.10, -0.38)	-1.00 (-2.12, -0.39)	-1.25 (-2.21, -0.51)	-1.26 (-2.35, -0.43)	-1.11 (-2.18, -0.51)	-1.36 (-2.93, -0.69)
	Increase in heavy precipitation events ^g in Southern Asia ^f [%]	9.22% (6.26, 13.84)	9.78 % (6.52, 13.63)	11.56 % (7.04, 18.50)	9.83% (7.03, 14.87)	10.27 % (6.50, 17.40)	16.74 % (9.60, 23.44)
Possible climate range in 2100 (reg+glob)	Global mean temperature warming in 2100 [°C]	1.29°C (1.27—1.31)	1.44°C (1.44—1.48)	1.88°C (1.85—1.93)	1.69°C (1.68—1.7)	1.89°C (1.88—1.91°C)	2.43°C (2.42—2.46°C)
	Warming in the Arctic ^h (TNn ^g) [°C]	3.78°C (3.29, 4.25)	4.07 °C (3.53, 4.72)	5.37 °C (4.60, 6.40)	4.90°C (4.30, 5.46)	5.37 °C (4.46, 6.38)	6.86 °C (5.83, 8.24)
	Warming in the contiguous United States ^h (TXx ^g) [°C]	1.82°C (1.33, 2.22)	2.00 °C (1.60, 2.48)	2.62 °C (2.17, 3.23)	2.45°C (1.98, 2.87)	2.62 °C (2.07, 3.24)	3.37 °C (2.88, 4.03)
	Warming in Central Brazil ^h (TXx ^g) [°C]	2.06°C (1.80, 2.40)	2.20 °C (1.95, 2.58)	2.88 °C (2.46, 3.48)	2.63°C (2.27, 3.00)	2.88 °C (2.43, 3.47)	3.57 °C (3.20, 4.18)
	Drying in the Mediterranean region ^h [std ^g]	-0.72 (-1.84, -0.27)	-0.96 (-1.89, -0.28)	-1.24 (-2.27, -0.45)	-1.09 (-2.23, -0.40)	-1.24 (-2.42, -0.46)	-1.38 (-2.79, -0.67)
	Increase in heavy precipitation events ^g in Southern Asia ^h [%]	6.48% (3.08, 11.15)	7.83 % (4.20, 12.00)	10.19 % (6.34, 16.67)	9.82% (6.69, 13.86)	10.19 % (6.46, 16.64)	15.47 % (8.72, 22.98)

^a 50th percentile (median) estimates for global temperature (i.e. as likely as not of being below or above values)

^b 66th percentile estimates for global temperature (i.e. 66% likelihood of being at or below values)

^c 90th percentile estimates for global temperature (i.e. 10% likelihood of being at or above values)

^d All 1.5°C scenarios from AR5 include a substantial probability of overshooting above 1.5°C global warming before returning to 1.5°C

^e The values indicate the median and the interquartile range in parenthesis (25th percentile and 75th percentile)


^f The regional projections in these rows provide the range [median (q25, q75)] associated with the *median* global temperature outcomes of the considered mitigation scenarios at *peak warming* (see Box 1 and Suppl. Text 1 for details).

^g TNn: annual minimum night-time temperature; TXx: annual maximum day-time temperature; std: drying of soil moisture expressed in units of standard deviations of pre-industrial climate (1861-1880) variability; Rx5day: annual maximum consecutive 5-day precipitation (see Suppl. Text 1 for details)


^h Same as footnote f, but for the regional responses associated with the *median* global temperature outcomes of the considered mitigation scenarios by 2100 (see Box 1 and Suppl. Text 1 for details).

Supplementary Table S3: List of 26 CMIP5 models used in the analysis of climate extremes (see Wartenburger et al. 2017 for more details).


Model	Modelling Centre	Historical	RCP2.6	RCP4.5	RCP6.0	RCP8.5
ACCESS1-0	Commonwealth Scientific and Industrial Research Organization (CSIRO) and Bureau of Meteorology (BOM), Australia	x		x		x
bcc-csm1-1	Beijing Climate Center, China Meteorological Administration	x	x	x	x	x
bcc-csm1-1-m		x	x	x	x	x
CanESM2	Canadian Centre for Climate Modelling and Analysis	x	x	x		x
CCSM4	National Center for Atmospheric Research, USA	x		x	x	x
CMCC-CM	Centro Euro-Mediterraneo per i Cambiamenti Climatici, Italy	x		x		x
CMCC-CMS		x		x		x
CNRM-CM5	Centre National de Recherches Météorologiques / Centre Européen de Recherche et Formation Avancées en Calcul Scientifique, France	x	x	x		x
CSIRO-Mk3-6-0	Commonwealth Scientific and Industrial Research Organization / Queensland Climate Change Centre of Excellence, Australia	x	x	x	x	x
FGOALS-s2	LASG, Institute of Atmospheric Physics, Chinese Academy of Sciences	x	x	x	x	x
GFDL-CM3		x	x		x	x
GFDL-ESM2G	NOAA Geophysical Fluid Dynamics Laboratory, USA	x	x	x	x	x
GFDL-ESM2M		x	x	x	x	x
HadGEM2-CC		x		x		x
HadGEM2-ES	Met Office Hadley Centre, United Kingdom	x	x		x	x
inmcm4	Institute for Numerical Mathematics, Russia	x		x		x
IPSL-CM5A-LR		x	x	x	x	x
IPSL-CM5A-MR	Institut Pierre-Simon Laplace, France	x	x		x	x
IPSL-CM5B-LR		x		x		x
MIROC-ESM	Japan Agency for Marine-Earth Science and Technology, Atmosphere and Ocean Research Institute (The University of Tokyo), and National Institute for Environmental Studies	x	x	x	x	x
MIROC-ESM-CHEM		x	x	x	x	x
MIROC5		x	x	x	x	x
MPI-ESM-LR	Max Planck Institute for Meteorology, Germany	x	x	x		x
MPI-ESM-MR		x	x	x		x
MRI-CGCM3	Meteorological Research Institute, Japan	x	x	x	x	x
NorESM1-M	Norwegian Climate Centre	x	x	x	x	x



Supplementary Figure S1: Same as Fig. 2, but additionally including median estimates (see Suppl. Table S2).



Supplementary Figure S2: Same as Fig. 3, but additionally including median estimates (see Suppl. Table S2).



Supplementary Figure S3: Spread in CMIP5 simulations for recent and present-day climate (0.5°C and 1°C global warming)

References:

Clarke, L. et al. Assessing Transformation Pathways. In: Edenhofer, O., et al. (eds.), *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press, 413-510 (2014).

Rogelj, J., et al. Energy system transformations for limiting end-of-century warming to below 1.5°C. *Nature Clim. Change*, 5(6), 519-527 (2015).

Seneviratne, S.I., Donat, M.G., Pitman, A.J., Knutti, R., & Wilby, R.L. Allowable CO₂ emissions based on regional and impact-related climate targets. *Nature*, 529, 477-483, doi:10.1038/nature16542 (2016).

Wartenburger, R. et al. Changes in regional climate extremes as a function of global mean temperature: an interactive plotting framework. *Geosci. Model Dev.*, 10, 3609–3634, <https://doi.org/10.5194/gmd-10-3609-2017> (2017).