

REPLY • OPEN ACCESS

Reply to Comment on 'Extreme weather events in early summer 2018 connected by a recurrent hemispheric wave-7 pattern'

To cite this article: Kai Kornhuber *et al* 2025 *Environ. Res. Lett.* **20** 018001

View the [article online](#) for updates and enhancements.

You may also like

- [A geospatial approach for dynamic on-road emission through open-access floating car data](#)

Pak Lun Fung, Omar Al-Jaghbeer, Jia Chen *et al.*

- [The impact of human disturbances on the regeneration layer of tropical rainforests](#)

Ricard Arasa-Gisbert, Víctor Arroyo-Rodríguez and Jorge A Meave

- [Sensitivity of the impact of aeolian sand cover on permafrost degradation in the Qinghai–Xizang Plateau](#)

Luyang Wang, Ziteng Fu, Guanli Jiang *et al.*



The Electrochemical Society
Advancing solid state & electrochemical science & technology



**249th
ECS Meeting**
May 24-28, 2026
Seattle, WA, US
*Washington State
Convention Center*

Spotlight Your Science

**Submission deadline:
December 5, 2025**

SUBMIT YOUR ABSTRACT

ENVIRONMENTAL RESEARCH
LETTERS

REPLY

OPEN ACCESS

RECEIVED
9 October 2024REVISED
13 November 2024ACCEPTED FOR PUBLICATION
21 November 2024PUBLISHED
10 December 2024

Original content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.

Reply to Comment on ‘Extreme weather events in early summer
2018 connected by a recurrent hemispheric wave-7 pattern’Kai Kornhuber^{1,2,*} , Dim Coumou³ , Stefan Petri⁴ , Scott Osprey⁵ and Stefan Rahmstorf¹¹ Energy, Climate and Environment (ECE), International Institute for Applied Systems Analysis, Laxenburg, Austria² Lamont-Doherty Earth Observatory, Columbia University, New York City, NY, United States of America³ Faculty of Science, Water and Climate Risk, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands⁴ RDI - Earth System Analysis, Potsdam Institute for Climate Impact Research, Potsdam, Germany⁵ Atmospheric, Oceanic and Planetary Physics, Oxford University, Oxford, United Kingdom

* Author to whom any correspondence should be addressed.

E-mail: kornhuber@iiasa.ac.at**Keywords:** circumglobal teleconnections, atmosphere dynamics, weather extremes**Abstract**

Circumglobal teleconnections from wave-like patterns in the mid-latitude jets can lead to synchronized weather extremes in the mid-latitudes of Northern and Southern hemispheres. The simultaneous occurrence of record breaking and persistent northern hemisphere temperature anomalies in Summer 2018 were previously discussed in the context of a persistent zonally elongated wave-7 pattern that stretched over large parts of the northern hemisphere over an extended time and led to considerable societal impacts. Various diagnostics have been put forward to quantify and detect such wave patterns, many of which incorporate low-pass time filtering to separate signal from noise. In this response we argue that advancing our understanding of the large-scale circulation's response to anthropogenic climate change and reducing associated uncertainties in future climate risk requires a diverse range of perspectives and diagnostics from both the climate and weather research communities.

**1. On the importance of selecting process
relevant time scales in climate research**

Time averaging is a common practice in climate science. Serving as a filter for noise and short-term variability, applying time averages helps to distinguish between short term fluctuations of day to day weather and long term persistent patterns associated with slowly moving components of the earth system associated with climate variability or change.

Modern reanalysis data sets, such as ERA5 offer climate data at an unprecedented hourly resolution at 36 vertical levels with an approximate horizontal resolution of 31 km at the Earth's-surface. A reasonable approach in data formatting is to trim time and spatial scales to the type of climate or weather phenomena one aims at investigating. To study long-term global climate change annual means on a global scale might be most informative [1]. To investigate certain modes of variability, such as the El Niño–Southern Oscillation (ENSO) [2] or the Madden–Julian oscillation [3] one would choose monthly to seasonal averages. For the study of individual storms

or heavy precipitation events daily or sub-daily data would be most useful [4]. When studying the large-scale atmospheric patterns that serve as steering patterns and background conditions for higher-frequency weather phenomena, weekly to monthly averages are typically most insightful. The same applies to the study of persistent heatwaves, which emerge and decline on weekly [5–7] to monthly timescales [8].

**2. On the existence of circumglobal
teleconnections in the midlatitude jets**

Globally concurrent and locally record-breaking temperatures driven by thermodynamic and dynamical responses to anthropogenic activity have revived the interest in the investigation of recurrent large-scale atmospheric patterns [9–15]. Circumglobal teleconnections (CGTs) formed and guided by the mid-latitude jet streams can be mathematically described by Rossby wave theory and have led to established theories behind CGTs over nearly a century of research [16].

A rich body of literature on recurrent stationary patterns in the midlatitude circulation exists going back decades. This body of literature describes CGTs from a phenomenological and theoretical standpoint and provides evidence for their existence using observational data and models of various complexity [17–19]. Branstator [20] showed the existence of CGTs in the Northern Hemisphere's winter (December–February), which when a disturbance is placed in the jet core, would be guided by mid-latitude waveguides in a quasi-circumglobal manner. Ding and Wang [21] showed the existence of such patterns in the Northern Hemisphere summer highlighting strong links to the Indian summer monsoon and ENSO. Assessing the leading modes of the Northern Hemisphere summer circulation, Schubert *et al* [22, 23] showed that large-scale circumglobal patterns also exist on submonthly timescales. A common feature of these studies is the employment of sub-monthly to seasonal timescales to identify, characterize and suggest theoretical models to explain the emergence of CGT, their paths and locations. In his comment to our paper Kornhuber *et al* [24] (K19 hereafter), Jacopo Riboldi (JR hereafter) states that K19 would have claimed to identify ‘a novel type of CGT mediated by Rossby waves.’ Given the numerous papers from various different groups and different generations of atmospheric scientists (also cited and discussed in K19), the above cited claim of JR cannot be considered true.

3. Quasistationary Rossby waves and synchronized weather extremes

Furthermore, quasi-stationary Rossby waves acting on timescales from weeks to months can form CGTs with strong impacts on local weather and extremes. Teng *et al* [25] identified a submonthly wave-5 CGT with strong links to U.S. heatwaves. Petoukhov *et al* [10, 26] showed that several high impact heatwaves in spring to summer were associated with high amplitude hemispheric slow-moving Rossby waves assessed on monthly and sub-monthly timescales. In addition, several singular high impact extremes were discussed in the context of quasi-stationary waves, including the 2003 European heatwave [7, 9, 15], the concurrent 2010 Russian heatwave and Pakistani floods [8, 27], the 2018 European heatwave [24, 28, 29], the 2020 record breaking Siberian heatwave [5] and the record shattering 2021 Pacific northwest heatwave [30–32]. From a statistical perspective, Screen & Simmonds [33] showed that high amplitude Rossby waves assessed on monthly hemispheric mean meridional wind can be generally considered to be significantly related to local weather extremes. Coumou *et al* [34] showed that heat extremes can become globally synchronized when hemispheric synoptic scale wave patterns on roughly monthly timescales emerge, while Wang *et al* [14] identified the impact of circumglobal patterns on local rainfall

extremes. Recently, Kornhuber *et al* [7] found that CGTs assessed on weekly timescales significantly favor the emergence of concurrent heat extremes in specific hotspot regions across the mid-latitudes.


4. Quantifying the slow-moving dynamics of synchronized weather extremes in summer 2018

In summer 2018, the Northern Hemisphere extratropics experienced several heat and precipitation extremes [29, 35, 36] some of which coincided with a strongly meandering jet [37, 38] which was categorized as an amplified wave-7 pattern [24, 28] (also see figure 1).

In K19, wave patterns were identified by applying a fast Fourier transformation (FFT) on the 15 d mean of the mid-latitude meridional wind in the upper troposphere (300 mb) following the approach described in detail in earlier publications from a similar group of authors [10, 12, 15, 26, 34]. This approach has several advantages as it characterizes the hemispheric flow through four interpretable metrics, which can be linked to linear theory without effort: wavenumber, wave-amplitude, phase position and phase speed (note that in K19 phase speeds are quantified based on daily fields before a 15 d running mean is applied, they are not based on a visual analysis as suggested by JR). To visualize the temporal evolution of this pattern and its latitudinal variation a Hovmöller diagram of the 15 d running mean meridional winds in summer 2018 was generated (figure 2(b) in K19) which illustrates the emergence of an amplified ridge-through pattern with the strongest signals over Eurasia, coinciding in time and space with the most severe heat and rainfall extremes during that summer.

5. Quantitative methods surpass visual interpretations

In his comment to K19, JR remarks that ‘figure 2(b) of K19 is actually a time-filtered version of the usually employed Hovmöller diagram, which is normally based on unfiltered meridional wind’. In contrast to JR's assertion, we want to highlight that Hovmöller diagrams are not limited to a specific time scale or investigated variable. Numerous examples and applications exist in which illustration the temporal evolution over one spatial dimension of a given variable helps in gaining new insights (see e.g. www.e-education.psu.edu/meteo820/node/549 for numerous examples of peer-reviewed research using Hovmöller diagrams on timescales from daily running means, months to years). JR continues to suggest that applying a 15 d running mean for wave amplitude detection was not clearly communicated in K19 ‘The result of this low-pass filtering operation, which is not explicitly mentioned in K19, is shown in figure 1(b)’. It is important to note that the 15 d mean



perspective is mentioned several times throughout the main manuscript and methods section. All methods including the application of a 15 d running mean follow the approaches of earlier peer-reviewed publications from the same group of authors (see above).

While a visual analysis of Hovmöller diagrams might be common practice in some communities, we want to stress that caution is advised when using these insights for physical interpretation. Quantitative methods (as applied for K19 and shown in figures 2(c)–(e)) should always be the preferred approach where possible. Wave event, phase speed and position were based on the FFT and did not rely on the visual interpretation of figure 2(b). Naturally, a 6 h Hovmöller diagram (a nitpicky reader might ask JR, why did he not go for hourly data?) allows for different insights compared to a 15 d running mean illustration. While the former allows for the investigation of small-scale dynamics and mechanisms as well as localized turbulent features, a time filtered perspective can expose persistent features and allows for a mathematical description based on linear theory.

Most importantly, we want to highlight that a natural visual interpretation in the complementary perspective of the 6 h Hovmöller diagram shown by JR (figure 1(a)) comes to the same conclusion as K19: the emergence and persistence of quasi-stationary waves in its preferred wave-7 CGT phase position, overlaid with eastward migrating high frequency patterns (which are removed by the 15 d running mean in K19). Thus, while JR claims that the approach by K19 ‘displayed an unrealistic and incomplete representation of the hemispheric Rossby wave pattern’, we assert that figure 1(a) in JR serves as a confirmation that indeed stationary eddies were present over a wide range of the Northern Hemisphere mid-latitudes for one to two weeks, which are thus not an artifact of a time filter.

More generally, while 2-dimensional representation of a complex system like the atmosphere will be incomplete, there is nothing ‘unrealistic’, or ‘misleading’ (as stated elsewhere in JR) about the Hovmöller figure shown in K19: as with any data-analysis

exercise, the figures require an interpretation which is in accordance with the data processing and under consideration of the context provided. This also relates to JRs statements on group velocity vs phase velocity, where a visual interpretation and manually drawn arrows might not be an appropriate method to gain useful insights in this case.

6. Circumglobal teleconnections amid the weather–climate schism

Various metrics exist that allow for characterization of the state of the mid latitude flow. These range from local blocking indicators [39], to longitudinally extended wave packages [40], to waviness metrics [41]. Each of these diagnostics are motivated by certain spatial and temporal scales that allow for statements on different physical processes. Naturally, their choice and interpretation should be informed by their advantages and limitations. K19 shows the existence of a preferred phase for a wave-7 pattern that is quantified on a hemispheric basis. An amplified wave-7 in its preferred phase position was detected during the synchronous weather extremes of summer 2018 and can be considered a dominant driver in their maintenance and emergence. Complementary perspectives include local blocking patterns, a stationary wave package over Eurasia, a double jet pattern [42] or a positive summer NAO [28]. In a recent study by Rousi *et al* [29] (Kornhuber and Coumou are co-authors), these perspectives were comprehensively summarized (notably also featuring a daily Hovmöller plot in figure 4, which is discussed to a large extent).

We want to conclude that the complex character of the midlatitude circulation and its response to global warming will not be captured and understood by using a single metric or time-scale alone. Multiple perspectives and approaches from dynamical meteorology, climate and weather research are needed which will help to produce a comprehensive picture through their diversity. In a recent essay Randall & Emmanuel [43] note the unfortunate separation of two distinct communities raised on ‘Weather’ and

‘Climate’, respectively, whose research centers around the atmospheric circulation but tends to use very distinct methods and scales: observations from station data, daily to sub-daily timescales, meso- to synoptic scales (weather) vs reanalysis, model data, long term means and synoptic to global scales (climate). Overcoming the challenging and pressing questions associated with the atmosphere-dynamical response to climate change will require expertise from all related fields and a scientific discussion around different hypotheses and their supporting evidence. Given the rapid rise in extremes—and our limited understanding of the underlying dynamical processes—it is essential to have this scientific debate, but this does not benefit from careless framing of specific contributions as ‘unrealistic’ or ‘misleading’.

ORCID iDs

Kai Kornhuber  <https://orcid.org/0000-0001-5466-2059>

Dim Coumou  <https://orcid.org/0000-0003-2155-8495>

Stefan Petri  <https://orcid.org/0000-0002-4379-4643>

References

- [1] Calvin K et al 2023 IPCC, 2023: *Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* ed H Lee and J Romero (Core Writing Team) (IPCC) (<https://doi.org/10.59327/IPCC/AR6-9789291691647>)
- [2] Seager R, Cane M, Henderson N, Lee D-E, Abernathey R and Zhang H 2019 Strengthening tropical Pacific zonal sea surface temperature gradient consistent with rising greenhouse gases *Nat. Clim. Change* **9** 517–22
- [3] Madden R A and Julian P R 1972 Description of global-scale circulation cells in the tropics with a 40–50 day period *J. Atmos. Sci.* **29** 1109–23
- [4] Pendergrass A G, Knutti R, Lehner F, Deser C and Sanderson B M 2017 Precipitation variability increases in a warmer climate *Sci. Rep.* **7** 17966
- [5] Gloege L, Kornhuber K, Skulovich O, Pal I, Zhou S, Ciais P and Gentile P 2022 Land-atmosphere cascade fueled the 2020 Siberian heatwave *AGU Adv.* **3** e2021AV000619
- [6] Kornhuber K, Lesk C, Schleussner C F, Jägermeyr J, Pfleiderer P and Horton R M 2023 Risks of synchronized low yields are underestimated in climate and crop model projections *Nat. Commun.* **14** 3528
- [7] Kornhuber K, Coumou D, Vogel E, Lesk C, Donges J F, Lehmann J and Horton R M 2020 Amplified Rossby waves enhance risk of concurrent heatwaves in major breadbasket regions *Nat. Clim. Change* **10** 48–53
- [8] Di Capua G, Sparrow S, Kornhuber K, Rousi E, Osprey S, Wallom D, van den Hurk B and Coumou D 2021 Drivers behind the summer 2010 wave train leading to Russian heatwave and Pakistan flooding *npj Clim. Atmos. Sci.* **4** 55
- [9] Tachibana Y, Nakamura T, Komiya H and Takahashi M 2010 Abrupt evolution of the summer Northern Hemisphere annular mode and its association with blocking *J. Geophys. Res. Atmos.* **115** 2009JD012894
- [10] Petoukhov V, Rahmstorf S, Petri S and Schellnhuber H J 2013 Quasiresonant amplification of planetary waves and recent Northern Hemisphere weather extremes *Proc. Natl Acad. Sci.* **110** 5336–41
- [11] Petoukhov V, Petri S, Kornhuber K, Thonicke K, Coumou D and Schellnhuber H J 2018 Alberta wildfire 2016: apt contribution from anomalous planetary wave dynamics *Sci. Rep.* **8** 12375
- [12] Kornhuber K, Petoukhov V, Karoly D, Petri S, Rahmstorf S and Coumou D 2017 Summertime planetary wave resonance in the Northern and Southern Hemispheres *J. Clim.* **30** 6133–50
- [13] Teng H, Leung R, Branstator G, Lu J and Ding Q 2022 Warming pattern over the Northern Hemisphere midlatitudes in boreal summer 1979–2020 *J. Clim.* **35** 3479–94
- [14] Wang S, Davies R E and Gillies R R 2013 Identification of extreme precipitation threat across midlatitude regions based on short-wave circulations *J. Geophys. Res. Atmos.* **118** 11–059
- [15] Kornhuber K, Petoukhov V, Petri S, Rahmstorf S and Coumou D 2017 Evidence for wave resonance as a key mechanism for generating high-amplitude quasi-stationary waves in boreal summer *Clim. Dyn.* **49** 1961–79
- [16] Rossby C-G et al 1939 Relation between variations in the intensity of the zonal circulation of the atmosphere and the displacements of the semi-permanent centers of action *J. Mar. Res.* **2** 38–55
- [17] Hoskins B and Woollings T 2015 Persistent extratropical regimes and climate extremes *Curr. Clim. Change Rep.* **1** 115–24
- [18] Hoskins B J and Karoly D J 1981 The steady linear response of a spherical atmosphere to thermal and orographic forcing *J. Atmos. Sci.* **38** 1179–96
- [19] Jiménez-Esteve B, Kornhuber K and Domeisen D I V 2022 Heat extremes driven by amplification of phase-locked circumglobal waves forced by topography in an idealized atmospheric model *Geophys. Res. Lett.* **49** e2021GL096337
- [20] Branstator G 2002 Circumglobal teleconnections, the jet stream waveguide, and the North Atlantic oscillation *J. Clim.* **15** 1893–910
- [21] Ding Q and Wang B 2005 Circumglobal teleconnection in the Northern Hemisphere summer* *J. Clim.* **18** 3483–505
- [22] Schubert S D, Wang H, Koster R D, Suarez M J and Groisman P Y 2014 Northern Eurasian heat waves and droughts *J. Clim.* **27** 3169–207
- [23] Schubert S, Wang H and Suarez M 2011 Warm season subseasonal variability and climate extremes in the Northern Hemisphere: the role of stationary Rossby waves *J. Clim.* **24** 4773–92
- [24] Kornhuber K, Osprey S, Coumou D, Petri S, Petoukhov V, Rahmstorf S and Gray L 2019 Extreme weather events in early summer 2018 connected by a recurrent hemispheric wave-7 pattern *Environ. Res. Lett.* **14** 054002
- [25] Teng H, Branstator G, Wang H, Meehl G A and Washington W M 2013 Probability of US heat waves affected by a subseasonal planetary wave pattern *Nat. Geosci.* **6** 1056–61
- [26] Petoukhov V, Petri S, Rahmstorf S, Coumou D, Kornhuber K and Schellnhuber H J 2016 Role of quasiresonant planetary wave dynamics in recent boreal spring-to-autumn extreme events *Proc. Natl Acad. Sci.* **113** 6862–7
- [27] Lau W K M and Kim K-M 2012 The 2010 Pakistan flood and Russian heat wave: teleconnection of hydrometeorological extremes *J. Hydrometeorol.* **13** 392–403
- [28] Drouard M, Kornhuber K and Woollings T 2019 Disentangling dynamic contributions to summer 2018 anomalous weather over Europe *Geophys. Res. Lett.* **46** 12537–46
- [29] Rousi E et al 2023 The extremely hot and dry 2018 summer in central and northern Europe from a multi-faceted weather and climate perspective *Nat. Hazards Earth Syst. Sci.* **23** 1699–718
- [30] Bartusek S, Kornhuber K and Ting M 2022 2021 North American heatwave amplified by climate change-driven nonlinear interactions *Nat. Clim. Change* **12** 1143–50

- [31] White R H *et al* 2023 The unprecedented Pacific Northwest heatwave of June 2021 *Nat. Commun.* **14** 727
- [32] Neal E, Huang C S Y and Nakamura N 2022 The 2021 Pacific Northwest heat wave and associated blocking: meteorology and the role of an upstream cyclone as a diabatic source of wave activity *Geophys. Res. Lett.* **49** e2021GL097699
- [33] Screen J A and Simmonds I 2014 Amplified mid-latitude planetary waves favour particular regional weather extremes *Nat. Clim. Change* **4** 704–9
- [34] Coumou D, Petoukhov V, Rahmstorf S, Petri S and Schellnhuber H J 2014 Quasi-resonant circulation regimes and hemispheric synchronization of extreme weather in boreal summer *Proc. Natl Acad. Sci.* **111** 12331–6
- [35] White R H, Kornhuber K, Martius O and Wirth V 2022 From atmospheric waves to heatwaves: a waveguide perspective for understanding and predicting concurrent, persistent, and extreme extratropical weather *Bull. Am. Meteorol. Soc.* **103** E923–E35
- [36] Vogel M M, Zscheischler J, Wartenburger R, Dee D and Seneviratne S I 2019 Concurrent 2018 hot extremes across Northern Hemisphere due to human-induced climate change *Earths Future* **7** 692–703
- [37] Liu X, He B, Guo L, Huang L and Chen D 2020 Similarities and differences in the mechanisms causing the European summer heatwaves in 2003, 2010, and 2018 *Earths Future* **8** e2019EF001386
- [38] Dong W, Jia X and Wu R 2024 Impact of summer Tibetan Plateau snow cover on the variability of concurrent compound heatwaves in the Northern Hemisphere *Environ. Res. Lett.* **19** 014057
- [39] Barriopedro D, García-Herrera R, Lupo A R and Hernández E 2006 A climatology of Northern Hemisphere blocking *J. Clim.* **19** 1042–63
- [40] Wirth V, Riemer M, Chang E K M and Martius O 2018 Rossby wave packets on the midlatitude waveguide—a review *Mon. Weather Rev.* **146** 1965–2001
- [41] Röthlisberger M, Pfahl S and Martius O 2016 Regional-scale jet waviness modulates the occurrence of midlatitude weather extremes *Geophys. Res. Lett.* **43** 10–989
- [42] Rousi E, Kornhuber K, Beobide-Arsuaga G, Luo F and Coumou D 2022 Accelerated western European heatwave trends linked to more-persistent double jets over Eurasia *Nat. Commun.* **13** 3851
- [43] Randall D A and Emanuel K 2024 The weather–climate schism *Bull. Am. Meteorol. Soc.* **105** E300–5