

TOPICAL REVIEW • OPEN ACCESS

The implications of overshooting 1.5 °C on Earth system tipping elements—a review

To cite this article: Paul D L Ritchie *et al* 2026 *Environ. Res. Lett.* **21** 043001

View the [article online](#) for updates and enhancements.

You may also like

- [Temporally clustered streamflow events control focused groundwater recharge in drylands](#)
Gabriel C Rau, José Bastías Espejo, R Ian Acworth *et al.*
- [Observation of CH₄, CO and N₂O in the upper troposphere and lower stratosphere over the Qinghai–Tibet Plateau](#)
You Yi, Zhaonan Cai, Yi Liu *et al.*
- [Higher efficacy of SO₂ and accumulation mode-H₂SO₄ stratospheric aerosol injection: insights from CESM2 and GEOS-Chem with advanced particle microphysics \(APM\)](#)
Fangqun Yu, Gan Luo and Arshad Nair

ENVIRONMENTAL RESEARCH
LETTERS

TOPICAL REVIEW

The implications of overshooting 1.5 °C on Earth system tipping elements—a review

Paul D L Ritchie^{1,2,27,*} , Norman J Steinert^{3,4,27} , Jesse F Abrams¹ , Hassan Alkhayuon⁵ , Constantin W Arnscheidt⁶ , Nils Bochow^{4,7,8} , Ruth R Chapman⁹ , Joseph Clarke^{1,2} , Donovan P Dennis^{4,10} , Jonathan F Donges^{4,10,11} , Bernardo M Flores^{12,13,14} , Julius Garbe⁴ , Annika Högner^{15,16} , Chris Huntingford¹⁷ , Timothy M Lenton¹ , Johannes Lohmann⁹ , Kerstin Lux-Gottschalk¹⁸ , Manjana Milkoreit¹⁹ , Tessa Möller^{15,16} , Paul Pearce-Kelly²⁰ , Laura Pereira^{10,21} , Courtney Quinn²² , Carl-Friedrich Schleussner^{15,16} , Simone M Stuenzi^{8,16} , Didier Swingedouw²³ , Larissa N Van der Laan⁹ , Kirsten Zickfeld²⁴ , and Nico Wunderling^{4,25,26,27,*} 

RECEIVED
28 August 2025

REVISED
15 January 2026

ACCEPTED FOR PUBLICATION
23 January 2026

PUBLISHED
19 February 2026

Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



- ¹ Global Systems Institute, Faculty of Environment, Science and Economy, University of Exeter, North Park Road, Exeter, United Kingdom
- ² Department of Mathematics and Statistics, Faculty of Environment, Science and Economy, University of Exeter, North Park Road, Exeter, United Kingdom
- ³ CICERO Center for International Climate Research, Oslo, Norway
- ⁴ Potsdam Institute for Climate Impact Research (PIK), Member of the Leibniz Association, Potsdam, Germany
- ⁵ School of Mathematical Sciences, University College Cork, Western Road, Cork, Ireland
- ⁶ Centre for the Study of Existential Risk, University of Cambridge, Cambridge, United Kingdom
- ⁷ Department of Mathematics and Statistics, Faculty of Science and Technology, UiT The Arctic University of Norway, Tromsø, Norway
- ⁸ Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Potsdam, Germany
- ⁹ Physics of Ice, Climate and Earth, Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
- ¹⁰ Stockholm Resilience Centre, Stockholm University, Stockholm, Sweden
- ¹¹ Max Planck Institute of Geoanthropology, Jena, Germany
- ¹² Equalsea-lab, University of Santiago de Compostela, Santiago de Compostela, Spain
- ¹³ Amazon Regional Observatory, Amazon Cooperation Treaty Organization (ORA/OTCA), Brasília, Brazil
- ¹⁴ Instituto Juruá, Manaus, Brazil
- ¹⁵ International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria
- ¹⁶ Geography Department, Humboldt-Universität zu Berlin, Berlin, Berlin, Germany
- ¹⁷ UK Centre for Ecology and Hydrology, Wallingford, Oxfordshire, United Kingdom
- ¹⁸ Centre for Analysis, Scientific Computing and Applications, Eindhoven University of Technology, Eindhoven, The Netherlands
- ¹⁹ Department of Sociology and Human Geography, University of Oslo, Oslo, Norway
- ²⁰ Zoological Society of London, Regent's Park, London, United Kingdom
- ²¹ Global Change Institute, University of the Witwatersrand, Johannesburg, South Africa
- ²² School of Natural Sciences, University of Tasmania, Hobart, Tasmania, Australia
- ²³ Environnements et Paléoenvironnements Océaniques et Continentaux (EPOC)—Université de Bordeaux, Pessac, France
- ²⁴ Department of Geography, Simon Fraser University, Vancouver, British Columbia, Canada
- ²⁵ Center for Critical Computational Studies (C3S), Goethe University Frankfurt, Frankfurt am Main, Germany
- ²⁶ Senckenberg Research Institute and Natural History Museum, Frankfurt am Main, Germany
- ²⁷ These authors have contributed equally.

* Authors to whom any correspondence should be addressed.

E-mail: paul.ritchie@exeter.ac.uk and wunderling@c3s.uni-frankfurt.de

Keywords: overshoots, tipping points, risk assessment, mitigation

Supplementary material for this article is available [online](#)

Abstract

Due to insufficient emission reductions in recent years, it is increasingly likely that global warming will exceed the 1.5 °C temperature limit in the late 2020s or 2030s. As a result, several Earth system tipping elements could, at least temporarily, have their tipping points surpassed, posing risks of large-scale and profound structural change. Tipping does not always occur immediately upon crossing such a critical threshold. If the length of time the driver is beyond the critical level is short enough, tipping could still be avoided for some slow-responding elements of the climate system.

An improved understanding is therefore needed of whether tipping remains avoidable, for which systems, and under what conditions. Here, we review how minimising the magnitude and duration of any temperature overshoot beyond 1.5 °C could decrease tipping risks. Tipping elements with fast response times, such as warm-water coral reefs, are especially vulnerable to overshoot. In contrast, those with slow response times, such as polar ice sheets, may be less sensitive to temporary overshoot. Potential interactions between tipping elements and additional human pressures, such as deforestation in the Amazon or pollution and overfishing of coral reef habitats, may further lower tipping points, narrowing the range of overshoot trajectories that can still avoid it. The vulnerability of many tipping elements, even under shorter overshoot conditions, underscores that global warming must peak below 2 °C above pre-industrial levels, return to below 1.5 °C as quickly as possible (i.e. within this century), and to around 1 °C thereafter to limit tipping point risks.

1. Introduction

Anthropogenic global warming is likely to exceed 1.5 °C relative to pre-industrial levels (Friedlingstein *et al* 2025), within the next 10 years (Rockström *et al* 2024), and current policies would take us towards 2.6 °C of global warming later this century (Climate Action Tracker 2025). However, this warming could be further amplified by possible tipping events or triggering so far unaccounted for climate-carbon cycle feedbacks (Steffen *et al* 2018, Ripple *et al* 2023a, Deutloff *et al* 2025). To eventually limit warming levels at or below 1.5 °C, it is now almost inevitable that there will be a period of temperature overshoot during which this warming target will be at least temporarily exceeded (Bustamante *et al* 2023, Bevacqua *et al* 2025, Reisinger *et al* 2025). Temperature overshoot refers to pathways of global warming (averaged over a climatological period of 30 years) that first exceed a warming target (such as 1.5 °C), followed by an eventual return to, or below, the warming target, as opposed to exceedance scenarios that permanently surpass a warming target. Therefore, temperature overshoots are often characterised by their peak warming, duration, and final stabilisation level (Schwinger *et al* 2022b, Wunderling *et al* 2023, Schleussner *et al* 2024). Two conceptual temperature overshoot pathways are illustrated in figure 1(a). Note that realistic scenarios that manage a return to 1.5 °C by the end of the century have an upper limit of peak warming of about 1.8 °C due to the feasibility and scale of carbon dioxide removal (CDR) deployment required to achieve the necessary decline (Reisinger *et al* 2025).

Many components of the climate system are judged to have critical thresholds in global warming beyond which positive feedbacks can cause irreversible and self-reinforcing change to a drastically different state, if exceeded for too long (Armstrong Mckay *et al* 2022, Lenton *et al* 2023). The critical thresholds are here referred to as *tipping points* and the components of the climate system as *tipping elements* (see

table 1 and section 4 for a description of individual tipping elements). If an element is tipped it can cause substantial negative impacts for human societies, the biosphere, and the climate system itself (Lenton *et al* 2008). During temperature overshoot, there is an increased risk of exceeding tipping points (i.e. pathways in figure 1(a) temporarily exceed an uncertainty range for a tipping point). While temperature overshoot pathways have many negative consequences, assessing whether they exceed tipping points and cause tipping is crucial, due to the potentially devastating impacts they bring.

When assessing risks of tipping from a temperature overshoot, considering the response timescale of the relevant system is important. Earth system tipping elements can be categorised by their response timescales to climate change: *slow* elements (e.g. ice sheets) have a response over multi-centennial to millennial timescales much slower than the manifestation time of current climate change itself, while *fast* tipping elements (e.g. North Atlantic Subpolar Gyre, monsoons, or warm water coral reefs) respond on similar or faster annual to few centuries timescales (Ritchie *et al* 2021, Swingedouw *et al* 2021, Bochow *et al* 2023, Lenton *et al* 2024).

For a temperature overshoot that exceeds a tipping point by a large amount and for a long time, there is a high risk of triggering tipping, even for slow elements (figure 1(b)). However, a temporary crossing of a tipping point will not necessarily result in triggering the tipping of some inherently slow systems, if the temperature overshoot is small and short (figure 1(c)) (Ritchie *et al* 2019). This is because such systems have substantial inertia and so are slow to respond to warming that has already occurred, as their response lags behind the forcing (i.e. anthropogenic greenhouse gas emissions).

Ideally, temperature exceedance of tipping points (even temporarily) should be avoided, but this may already be too late for some tipping elements (see table 1). That is, the lower (and, in the case of coral reefs, central) uncertainty ranges of some Earth

Table 1. List of tipping elements along with their global warming tipping point and tipping timescale estimates based on data from Lenton *et al* (2023), and in case of no updates, from Armstrong McKay *et al* (2022). *Note:* in instances where no minimum and/or maximum tipping timescale is given in the original source, the central estimate is also used as the minimum and/or maximum. Global tipping elements are written in bold and regional tipping elements are in regular text.

Tipping element	Global warming tipping point (°C)			Tipping timescale (years)		
	Est.	Min	Max	Est.	Min	Max
Cryosphere tipping elements						
Greenland ice sheet	1.5	0.8	3.0	10k	1k	15k
West Antarctic Ice Sheet	1.5	1.0	3.0	2k	500	13k
Marine Basins East Antarctica	3.0	2.0	6.0	2k	500	10k
Non-marine East Antarctica	7.5	6.0	10.0	10k	10k	10k
Mountain glaciers	2.0	1.5	3.0	200	50	1k
Land permafrost	1.5	1.0	2.3	200	100	300
Ecological tipping elements						
Amazon rainforest	3.5	2.0	6.0	100	50	200
Boreal forest	4.0	1.4	5.0	100	50	100
Warm-water coral reefs	1.2	1.0	1.5	10	10	10
Ocean and atmosphere tipping elements						
West African monsoon	2.8	2.0	3.5	50	10	500
North Atlantic subpolar gyre	1.8	1.1	3.8	10	5	50
Atlantic Meridional overturning circulation	4.0	1.4	8.0	50	15	300

system tipping points are already exceeded (figure 2). Further committed warming is inevitable and is almost certain to exceed 1.5 °C in the near future, therefore exceeding the lower uncertainty ranges of even more tipping points (placed at around 1.5 °C), and in the case of coral reefs exceeding the upper tipping point range, see figure 2. Therefore, limiting the peak warming and duration of temperature overshoot is crucial for avoiding many key Earth system tipping elements from tipping (Ritchie *et al* 2021). Hence, fundamental questions arise on what the limits for peak warming and overshoot duration are to avoid or limit the risk of tipping specific elements (Ritchie *et al* 2021).

The paper aims to evaluate the current status of a probabilistic risk assessment of tipping Earth system elements for temperature overshoot pathways of 1.5 °C. To address this, we consider how uncertainties in temperature overshoot and tipping element characteristics can be handled in section 2. In section 3 we assess the risk of triggering tipping during and after overshoot from a conceptual dynamical systems perspective. We explore the implications of a temperature overshoot for individual tipping elements in section 4, before we consider in section 5 how additional pressures may increase the risk of tipping. What these tipping risks imply for mitigation are identified in section 6. Finally, in section 7 we discuss what still needs to be addressed to form a quantitative probabilistic risk assessment.

2. Handling uncertainties relevant for tipping elements during and after temperature overshoots

Considering the potentially severe and irreversible consequences from triggering tipping events requires a framework for assessing and acting under uncertainty. Due to the limited predictability of the precise tipping points (Ben-Yami *et al* 2024, Boers *et al* 2025), a probabilistic risk assessment approach becomes essential. Similar methods to those used in insurance and actuarial industries can help quantify risks. Such approaches have recently been embraced in various reports for tipping points research (Laybourn *et al* 2024, Trust *et al* 2024, Saye *et al* 2025) and help quantify the likelihood of crossing tipping points under different emission scenarios (Abrams *et al* 2023, Wunderling *et al* 2023, Möller *et al* 2024, Deutloff *et al* 2025). Within this risk framework, we can systematically address different types of uncertainties in temperature overshoot and tipping element characteristics, and in cases where we cannot apply a probabilistic risk assessment yet, we detail what would be necessary to do so.

2.1. Types of uncertainties in global temperature trajectories

Understanding the characteristics of temperature overshoot pathways is crucial for assessing tipping risks and informing mitigation strategies that can minimise these dangerous consequences of climate




Figure 1. Schematic response of fast and slow systems with a common tipping point to different temperature overshoot pathways. (a) Two idealised temperature overshoot pathways that exceed a global warming target (purple dashed line), subsequently reach a level of peak warming, before returning and stabilising at the global warming target. These temperature overshoot pathways temporarily cross an uncertain tipping point (burning ember range) that is here assumed to be the same for both a fast and a slow system. One pathway has a high peak and long overshoot duration (black), the other a low peak and short overshoot duration (grey). (b) For the large and long temperature overshoot, the fast system (red) tips very quickly after the tipping point is crossed, while for the slow system (blue), tipping is delayed although it still occurs. (c) For the small and short temperature overshoot, the fast system (red) again tips, but tipping is avoided for the slow system (blue) despite also temporarily exceeding its tipping point, due to its inherent slow timescale. *Note* that the shown temperature trajectories only cover the case where the loss of a tipping element (i.e. tipping in the first place) is avoided. Instead, a recovery of an element after tipping is not shown as this would happen at even lower temperatures.

change (figure 2). Understanding the peak and duration of potential global temperature overshoots requires an examination of the relationship between greenhouse gas concentrations, emissions pathways, and the resulting warming. Deriving an emissions pathway perspective considers the full chain from greenhouse gas emissions to resulting temperature changes (Meinshausen *et al* 2024). While the concept of temperature overshoot scenarios is straightforward, projecting actual temperature trajectories from emissions involves dealing with substantial uncertainties. These uncertainties—stemming from several sources such as climate sensitivity (how much warming results from a given increase in greenhouse gases), carbon cycle feedbacks (how natural systems respond to warming), and the efficacy of mitigation measures—can be summarised as follows:

Uncertainties in climate sensitivity and resulting peak warming levels. The Transient Climate Response to cumulative carbon Emissions (TCREs) describes how much warming results from a given amount of cumulative CO₂ emissions (Allen *et al* 2009, Matthews *et al* 2009, Tachiiri *et al* 2019, Jones and Friedlingstein 2020, Meehl *et al* 2020, Nijse *et al* 2020, Winkler *et al* 2024). However, TCRE estimates vary considerably across models, with the IPCC AR6 likely range (17th–83rd percentile range) spanning 1.0 °C–2.3 °C per 1000 GtC (Canadell *et al* 2021). This uncertainty implies that even with scenarios that have identical cumulative CO₂ emissions, peak warming projections can differ substantially. Additionally, non-CO₂ forcings and carbon cycle feedbacks that may originate from the uncertainties in the impact of triggering tipping elements, such as permafrost thaw and forest changes, can




Figure 2. Overshooting 1.5 °C risks crossing Earth system tipping points. Illustrative temperature overshoot pathways, exceeding and then returning to below 1.5 °C (solid black line) and other stabilisation pathways (dashed black lines), dependent on uncertainties in future emissions and Earth system feedbacks. The depicted temperature overshoot pathway at least temporarily crosses several Earth system tipping points: whether tipping occurs depends on the peak warming, time spent beyond the tipping point and the element's response timescale. Icons denote tipping elements (global tipping elements are shown with a solid edge; regional tipping elements are shown with a dashed edge; local tipping elements are not shown; best-estimate tipping points based on data from Lenton *et al* (2023), and in case of no updates, from Armstrong Mckay *et al* (2022)). Uncertainties of tipping points are shown as shaded bars. *Current policies & Actions* lead to temperature increases of 2.6 °C (Climate Action Tracker 2025), which is far from the upper limit of 1.8 °C where a return to 1.5 °C by 2100 appears feasible (Reisinger *et al* 2025).

further extend this uncertainty range. While some of these feedbacks are already represented to varying degrees in the models used to derive current TCRE-estimates, many models (as of CMIP6) still lack a full representation of their dynamics and long-term impacts. Land and ocean carbon sinks have absorbed approximately 56% of anthropogenic emissions over 1960–2023, with the remaining portion retained in the atmosphere (Friedlingstein *et al* 2025). While the future of these carbon sinks remains uncertain, there is growing evidence of a long-term weakening of land sinks (e.g. for the Amazon rainforest; Gatti *et al* 2021, Ke *et al* 2024), whereas ocean carbon sinks, while also weakening long-term, are comparatively stable across Earth System Models (ESMs) (Tokarska *et al* 2019, Schwinger *et al* 2022b, Koven *et al* 2023, Sanderson *et al* 2025).

Uncertainties in temperature stabilisation and reversal. Even if emissions reach net-zero and are maintained at that level—known as the zero emissions commitment (ZEC) (Jones *et al* 2019, MacDougall *et al* 2020, Corner *et al* 2023), considerable uncertainty remains over how global temperatures would subsequently evolve. Some ESMs show continued warming after emissions cease, while others indicate a plateau or gradual cooling. These divergent responses primarily reflect differences in how models represent ocean heat and carbon uptake and the interplay of several positive and negative feedback processes within the Earth system that affect global warming. Prominent examples for positive climate feedbacks (meaning they amplify warming) are ice-albedo feedback, permafrost thaw or desertification (Lenton *et al* 2023, Ripple *et al* 2023b). Additionally, many tipping elements themselves possess self-reinforcing feedbacks through their disintegration, such as the large ice sheets or forest systems (e.g. Armstrong McKay *et al* 2022). Such feedbacks do not only affect TCRE during the warming ramp-up (see previous subsection) and subsequently whether stabilisation will occur after net-zero is reached, but they also strongly condition the reversibility of global temperatures following overshoot pathways. When emissions become net negative, the temperature reversal is determined by ZEC (here as committed warming after positive emissions) and TCRE (here reversed TCRE that quantifies temperature response during carbon removal). This response is further modified by (yet uncertain) corrections due to the asymmetry in temperature response to net positive and net negative emissions (Zickfeld *et al* 2016, 2021, Koven *et al* 2023, Sanderson *et al* 2025). In principle, reducing atmospheric greenhouse gas concentrations should cool the climate, yet these feedback-driven processes and hysteresis effects can delay or even prevent a return to previous climate states, particularly where tipping points are crossed. Together with current inadequate progress

in emissions reductions, which increases the risk of overshoot (Climate Action Tracker 2025, Forster *et al* 2025, World Meteorological Organization 2025), these uncertainties raise fundamental challenges for the feasibility of returning to internationally agreed temperature targets once they have been exceeded.

2.2. Implications for projected temperature overshoots and tipping

The characteristics of temperature overshoots are strongly influenced by the timing of emission reductions and carbon removals. Delayed action increases both the peak warming and the duration of the temperature overshoot, exacerbating tipping risks.

Pathways that temporarily exceed 1.5 °C with a 50% likelihood of returning to this warming level by 2100 exhibit a temperature overshoot of 0.1 °C–0.3 °C in the best estimate (Kikstra *et al* 2022). The IPCC categorises temperature overshoot scenarios as either ‘limited overshoot’ (C1), which refers to exceeding the specified limit by up to about 0.1 °C, or ‘high overshoot’ (C2), which refers to exceeding it by more than 0.1 °C and up to 0.3 °C (IPCC AR6 WGIII-SPM 2022, Schleussner *et al* 2024). Limited temperature overshoot pathways typically display exceedance of median temperature projections for up to a few decades before returning to below 1.5 °C by or before the year 2100 (IPCC SR1.5-SPM 2018). However, the uncertainty ranges around these projections are substantial, with much higher warming peaks possible due to stronger than expected climate sensitivity and Earth system feedbacks (Reisinger and Geden 2023, Schleussner *et al* 2024, Kaufhold *et al* 2025, Myhre *et al* 2025).

There has not been a thorough assessment of overshooting 2.0 °C as there has been for 1.5 °C. However, the scenario *SSP5-3.4-over* (although the underlying socio-economics in this scenario are now not likely given ongoing mitigation efforts) offers some insight into temperature pathways and the response of the carbon cycle in a pathway overshooting 2.0 °C (Melnikova *et al* 2021). Under this scenario, models show peak warming between 2.4 °C and 4.1 °C occurring between the 2040s and 2070s, after which temperatures decrease (due to carbon removal) and stabilise between 2 °C–3 °C by 2300, with temperatures remaining as high as ~3 °C above pre-industrial at the end of the 21st century. The rate of temperature decrease varies largely across models from steep to almost non-existent. Furthermore, most models show a decrease in TCRE during the cooling phase, reflecting hysteresis in the coupled carbon-climate system. Land and ocean carbon sinks weaken or even (temporarily) reverse for declining CO₂ concentration levels, leading to irreversibility in the cooling phase of an overshoot trajectory. As a result, atmospheric CO₂ concentrations decline more slowly for a given rate of anthropogenic

CO₂ removal, and the relationship between cumulative removals and temperature reduction deviates from that observed during the warming phase (Melnikova *et al* 2021).

2.3. Tipping element uncertainty

Risk assessments must account for uncertainties in tipping points, system timescales, and interactions. Central estimates for tipping points and timescales vary considerably across tipping elements (Armstrong Mckay *et al* 2022, Lenton *et al* 2023). The tipping point uncertainty range for each element also differs from very small values for the warm-water coral reefs (tipping point between 1.0 °C–1.5 °C, uncertainty: 0.5 °C) to large uncertainty ranges in the case of the Atlantic meridional overturning circulation (AMOC) (tipping point between 1.4 °C–8.0 °C, uncertainty: 6.6 °C), see figure 2. Likewise, the tipping timescale uncertainties can vary by up to two orders of magnitude for single elements. Both uncertainties will have profound impacts on a tipping element's risk of tipping during and after an overshoot (Lux-Gottschalk and Ritchie 2025). Further, tipping elements such as the AMOC or the large ice sheets, are not isolated subsystems but form a coupled system, whereby the number of stable states and tipping points can exceed the number of tipping elements. Not all interactions are fully understood, and there is large uncertainty in their strengths but also more fundamentally in their direction of interaction (Falkena and von der Heydt 2024, Wunderling *et al* 2024), and the impact of sub-system timescales on the response to forcing (Bastiaansen *et al* 2023). Interactions between elements can change effective tipping points (Wunderling *et al* 2023), and destabilising interactions introduce the risk of possible cascading transitions (Dekker *et al* 2018, Klose *et al* 2021, Wunderling *et al* 2024), which could be triggered even if only the lowest tipping point is crossed (Rosser *et al* 2024).

2.4. Structural uncertainties and alternative tipping mechanisms beyond saddle-node bifurcations

Studies of tipping during and after overshoot have mostly focused on the crossing of a smooth saddle-node bifurcation starting from an equilibrium state of the system (Ritchie *et al* 2019, 2021, Wunderling *et al* 2023), but the real Earth system is often more complicated. The initial states of relevant systems may not be simple equilibrium states (Lohmann and Ditlevsen 2021, Lohmann *et al* 2024b), opening up the possibility of other forms of tipping (Thompson *et al* 1994, Alkhayuon *et al* 2019, 2021, Ashwin and Newman 2021, Budd and Kuske 2024). For example, if the tipping element has periodic behaviour (e.g. monsoon systems such as in West Africa), the given phase of the period at the time an overshoot starts may determine if the system tips or not (Alkhayuon and Ashwin

2018). Therefore, the possibility of structural uncertainty further complicates the estimation of a peak warming and overshoot duration that avoids tipping.

Tipping can occur via other mechanisms such as rate-induced tipping, which is caused by an external forcing changing too quickly (Ritchie *et al* 2023). As one example, the AMOC, a tipping element with a tipping point between 1.4 °C and 8.0 °C (Armstrong Mckay *et al* 2022), has been shown to exhibit rate-induced tipping before a tipping point has been crossed both conceptually (Alkhayuon *et al* 2019, Chapman *et al* 2024) and in complex ocean models (Lohmann and Ditlevsen 2021, Lohmann *et al* 2024a). Rate-induced tipping complicates the commitment from overshoot scenarios, and in some cases may make them more dangerous. For example, some fast overshoots can be more dangerous than slower ones (O'Keeffe and Wiczorek 2020), as shown to be the case for a box-model of the AMOC (Ritchie *et al* 2023). Thus, the risk of tipping does not necessarily continually increase with the duration of the overshoot (Ritchie *et al* 2023). Similar behaviour might occur if a tipping element possesses multiple stable states (e.g. ice sheets, ocean circulation currents, vegetation patterns) that do not correspond to a full collapse of the system (Robinson *et al* 2012, Garbe *et al* 2020, Bastiaansen *et al* 2022, Bochow *et al* 2023, Lohmann *et al* 2024a), which could be reached for different overshoot profiles.

Structural uncertainty propagates further once interactions between tipping elements are considered (Wunderling *et al* 2024). Returning to the AMOC example, conceptual modelling already indicates that a tipping in the cryosphere can lead to rate-induced tipping in the AMOC (Lohmann *et al* 2021, Klose *et al* 2024b, Sinet *et al* 2024). There is thus a necessity to consider rate-induced tipping and other alternative scenarios in tipping cascade models to understand the full impact of various overshoot scenarios.

2.5. Modelling approaches to temperature overshoots and implications for tipping points

It is important to utilise the full modelling hierarchy when analysing the impact of temperature overshoots for tipping points. Conceptual models have the advantage of minimal computational cost, even for many stochastic realisations and long runs (Dijkstra 2024). Therefore, it is possible to qualitatively investigate tipping mechanisms for overshoot scenarios using conceptual models (e.g. Chapman *et al* 2024, Rosser *et al* 2024, Lux-Gottschalk and Ritchie 2025). Conceptual climate models generally describe subsystems of the Earth's climate and are limited in the scale and processes they model (Dijkstra 2013). Hence, conceptual models cannot provide a quantitative assessment for tipping risks during and after temperature overshoot. However, they can support the understanding of tipping behaviour in more complex models (Dijkstra 2024). Increasing scale leads to

intermediate complexity models of the ocean/atmosphere, while increasing both scale and processes results in ESMs of Intermediate Complexity (EMICs), and eventually complex ESMs.

Complex ESMs typically approximate solutions to the fundamental laws that govern the behaviour of the atmosphere, ocean, land, sea ice, land ice, and other components on a three-dimensional finite grid. While this allows for the resolution of a high number of processes, this makes it more difficult to diagnose the mechanisms causing the behaviour (e.g. tipping) observed in these complex models. Previous generations of ESMs were often limited to a horizontal resolution of 50–100 km (Eyring *et al* 2016); however, the next generation is expected to allow for coupled sub-20 km simulations (Roberts *et al* 2025). Even at this higher resolution, many important processes are still not explicitly resolved but are instead parameterised. Additionally, computational costs remain high, permitting a limited number of simulations on a 100–200 year timescale, making it challenging to investigate overshoot scenarios under consideration of all relevant uncertainties (see section above) especially in the context of crossing tipping points of slow tipping elements, which require long simulation runs. Furthermore, there have been discussions regarding the tuning process and the general ability of ESMs to represent tipping elements (Valdes 2011, Hourdin *et al* 2017). As models increase in complexity, they become more costly to run and challenging to explain output behaviour (Flato *et al* 2013). Conversely, as the number of processes and feedbacks increases, models produce a more realistic representation of overshoots.

However, there is another tool that is computationally efficient and can reproduce the behaviour of more complex climate models, providing a rapid translation of emissions into probabilistic estimates of changes to the physical climate system (Nicholls *et al* 2020), or explore the uncertainty associated with selected climate variables (Beusch *et al* 2021): These are so-called emulators. Emulators are simplified surrogates of complex models (typically ESMs) that reproduce their behaviour for a selection of variables to enable large, efficient ensemble runs within the limits of the training data. Physical emulation can be performed by simple parameterisations or statistical methods, where the emulator behaviour is tuned to reproduce the response of a given (more complex) model ensemble. Consequently, emulators can only resolve and show the impacts of non-linear and self-amplifying feedbacks if they have been accurately tuned against modelling approaches that include these processes. Because they can rapidly emulate complex model dynamics, emulators provide an excellent tool for developing quantitative frameworks for tipping-overshoot risk assessments, making them highly useful for probabilistic and ensemble-based analyses. To assess the probability of tipping for emission pathways, conceptual tipping models (e.g.

Wunderling *et al* 2021a) can be used in conjunction with climate emulators (Abrams *et al* 2023, Deutloff *et al* 2025, Möller *et al* 2024). Such approaches however, currently do not resolve direct temperature feedbacks from destabilising tipping elements (Möller *et al* 2024). Existing climate emulators are not fully tuned for overshoot scenarios due to limited training data, leading to large uncertainties about the response of the climate system. However, newer emulators have shown improved performance for such scenarios (Mathison *et al* 2025, Sandstad *et al* 2025).

These difficulties are increased by the challenges of detecting tipping points. Many systems experience *critical slowing down* before reaching a tipping point, indications of which have been modelled in a range of tipping elements including the AMOC (Boers 2021, Ditlevsen and Ditlevsen 2023) and the Amazon rainforest (Boulton *et al* 2022, Bochow and Boers 2023). There is no comparable empirical technique to indicate the exceedance of any tipping point prior to the transition to a new state. Nonetheless, modelling suggests that some tipping points may have already been crossed, however, the impacts of their crossing are not yet realised (i.e. the element is not yet tipped). Even further, it may be difficult or impossible to recognise before substantial impacts begin to unfold. According to current best estimates in the literature (Armstrong McKay *et al* 2022, Lenton *et al* 2023), it is possible that the tipping points of the Greenland ice sheet (GrIS), West Antarctic Ice Sheet (WAIS), North Atlantic Subpolar Gyre convection, warm water coral reefs, and boreal permafrost are below the current warming level (Lenton *et al* 2023, Forster *et al* 2025, Stokes *et al* 2025). However, only the warm water coral reefs are assessed to have a central tipping point estimate below 1.5 °C, where 1.5 °C corresponds to the coral reefs upper tipping point (Lenton *et al* 2023).

2.6. Approaches to constraining uncertainties

In the previous subsections we have covered many sources of uncertainties including: (i) future global temperature trajectories due to uncertainties in the relationship between greenhouse gas concentrations, emission pathways and the resulting warming; (ii) tipping element characteristics such as uncertainties in tipping points, timescales and interactions; (iii) structural and tipping mechanisms, that may cause tipping from rapid rates of increasing global warming as opposed to just the peak warming level and duration of an overshoot.

Reducing uncertainty in future forcing scenarios and tipping element characteristics is essential for estimating the response of individual tipping elements (Ritchie *et al* 2019). Given the role of uncertainty in temporarily crossing tipping points, tipping risks of selected emission pathways need to be evaluated probabilistically (Möller *et al* 2024). Using a simple conceptual climate model, Lux-Gottschalk

and Ritchie (2025) identify confidence levels for the location of the boundary that separates tipping from not tipping based on the peak overshoot and overshoot duration. They present a proof of concept, using Bayesian inference on how data could be used to better constrain uncertain system characteristics and hence also the tipping boundary. A similar approach could be followed for uncertainties in global temperature trajectories. From investigations of the impact of uncertainties in conceptual climate models (Rosser *et al* 2024), conclusions can be drawn on which model parameters have the biggest impact on tipping dynamics and thus need to be better constrained in more complex climate models.

Data-driven methods to better constrain the current proximity of tipping points have also been attempted. One avenue is to exploit critical slowing down as an early-warning signal of an approaching tipping point (Boers 2021, Boers and Rypdal 2021, Ben-Yami *et al* 2023, Ditlevsen and Ditlevsen 2023, Boers *et al* 2025). Another avenue that has recently gained attention involves using deep learning and AI-methods to predict both the occurrence and timing of tipping events (Huang *et al* 2024, Zhuge *et al* 2025). However, in the context of overshoots, extrapolating early warning signals to predict a tipping event can provide false conclusions due to assumptions on the future forcing trajectory (Ben-Yami *et al* 2024, Ashwin *et al* 2025). Additionally, early warning signals can fail when a system is not in equilibrium, such as for slow tipping elements with long internal timescales (e.g. ice sheets). Instead, new methods need to be developed to determine when an element has crossed its tipping point but not yet tipped (Gottwald and Gugole 2020). This would enable first order approximations for the chances of a return to the previous state (Ritchie *et al* 2019). Another method is to infer the strength of the underlying physical mechanisms from observations, thereby adjusting biases in ESMs, leading to better estimates of the tipping point, a so-called *observation-based constraint* (e.g. Liu *et al* 2017, Sgubin *et al* 2017, Portmann *et al* 2025).

3. Risk of triggering tipping during and after temperature overshoot


Multiple tipping elements of the Earth system could have their tipping point below 2 °C of global warming, and some below 1.5 °C (Lenton *et al* 2023, Armstrong McKay *et al* 2022; see figure 2). Consequently, even under the most optimistic emission scenarios of stabilising warming at 1.5 °C without any overshoot, it is considered *as likely as not* (33%–66% probability) that three Earth system elements will tip (see figure 3(a) based on Lenton *et al* 2023 or Armstrong McKay *et al* 2022). One of these elements will be the warm water coral reefs, which are *virtually certain* (>99% probability) to tip, given the upper range of their tipping point is 1.5 °C (Lenton

et al 2023; figure 2). The other two can be any combination of the North Atlantic Subpolar Gyre, land permafrost, and the Greenland and WAISs. However, it is considered *very unlikely* (<10% probability) for all of them to tip. However, delayed climate action and stabilising global warming at 2 °C (without an overshoot) would change the probability of five elements tipping from *very unlikely* to *as likely as not*. Mountain glaciers, boreal forests and the AMOC are the additional elements that risk being tipped under this scenario. Stabilising global warming at 3 °C would *likely* (>66% probability) cause eight elements to undergo tipping.

If the warming was subsequently reduced, such that 1.5 °C is only temporarily exceeded, the number of elements that would undergo tipping could be limited, see figure 3(b). As illustrated in figure 1, systems with a slow response timescale are more likely to avoid tipping than those with a fast response timescale for any given overshoot profile. This has been captured by some mathematical theory that describes a power law between the exceedance peak and duration, which not only depends on the tipping point but also the timescale of the system (Ritchie *et al* 2019). This theory indicates that doubling the exceedance peak requires reducing the exceedance duration of the tipping point by a factor of four to maintain the same level of tipping risk as before. Here is an example: assume that a particular tipping element would have a 1% probability of tipping if warming remains 100 years, with a 0.1 °C peak exceedance, beyond the tipping point. The theory suggests that if the peak exceedance of the same tipping point was instead 0.2 °C then the exceedance duration of the tipping point would need to be limited to 25 years to maintain the same 1% tipping risk. Utilising this theory, we here assess the tipping risk for overshoots of the 1.5 °C target (see figure 3(b)).

Limiting the temperature overshoot beyond 1.5 °C to a duration of 100 years could substantially reduce the number of elements that undergo tipping (compare figure 3(b) to 3(a)). For a peak warming of 2 °C, the number of elements that are *as likely as not* (33%–66% probability) to tip drops from five to three and tipping five becomes *very unlikely* (<10% probability) if warming is brought back to 1.5 °C (or below) within 100 years. A 100 year overshoot of 1.5 °C with a peak warming of 3 °C halves the number of elements *likely* (>66% probability) to be tipped (from eight to four) relative to staying at 3 °C. However, to limit such an overshoot scenario with a large peak to 100 years would require extraordinary carbon removal rates that are not considered to be feasible (Schleussner *et al* 2024).

Figure 4 extends this analysis to consider different overshoot durations for different levels of tipping risk (see Supplement for an animation). Using the best estimates from the latest literature (Armstrong McKay *et al* 2022, Lenton *et al* 2023), tipping elements



with fast timescales and a low warming tipping point (North Atlantic Subpolar Gyre and warm-water coral reefs) are the most susceptible to tipping even for short overshoot durations. In contrast, elements with slow timescales and a low tipping point (Greenland and WAISs) may avoid tipping if the overshoot duration is sufficiently short and warming stabilises below their tipping point (figure 4(a)). However, these ice sheets have a non-negligible probability that their tipping point is below 1.5 °C. Therefore, following a precautionary principle approach that seeks to limit the devastating long-term consequences of committing to ~10 m of sea-level rise from tipping these elements, means that limiting the time over 1.5 °C is not sufficient (figures 4(b) and (c)). Instead, warming must return to below 1.0 °C in the long term (Stokes *et al* 2025).

4. Implications of temperature overshoot for individual tipping elements

A temperature overshoot will have varying implications for the different tipping elements of the Earth system. Here we provide a brief summary for all elements considered in the previous sections, except for the terrestrial (non-marine) parts of the East Antarctic Ice Sheet, given its projected high tipping point, and the West African monsoon, which requires further research in the context of overshoot scenarios.

GrIS: The GrIS is generally assessed as a slow tipping element where a complete disappearance would occur on the time scale of centuries to millennia (Robinson *et al* 2012, Solgaard and Langen 2012, Pattyn *et al* 2018, Bochow *et al* 2023, Höning *et al* 2023, Petrini *et al* 2025). Different modelling studies show that an exceedance of the tipping point does not necessarily imply a complete melting of the GrIS but reversing the forcing can result in, at least, a partial regrowth of the ice sheet (Ridley *et al* 2010, Gregory *et al* 2020, Bochow *et al* 2023, Höning *et al* 2023, 2024). However, for some overshoot scenarios, a regrowth might not be possible as atmosphere-ice sheet interactions such as the surface mass balance-elevation feedback may impede reversibility (Pattyn *et al* 2018, Gregory *et al* 2020, Bochow *et al* 2023, Höning *et al* 2024). Additionally, there is substantial uncertainty about the tipping point global warming level (Noël *et al* 2021, Petrini *et al* 2025), which makes it difficult to assess an overshoot duration that would prevent a tipping of the GrIS. In any case, an exceedance of the tipping point implies substantial melt and therefore sea level rise, depending on the overshoot peak temperature and duration (e.g. Rückamp *et al* 2018, Bauer *et al* 2023).

WAIS: The response of the Antarctic Ice Sheet—the largest source of potential long-term sea-level rise—to global warming overshoot scenarios remains




Figure 4. Tipping risk for overshoots characterised by peak warming and time over 1.5 °C. (a) Best estimate for the cumulative number of tipped elements for overshoots characterised by their peak warming and time over 1.5 °C. (b) and (c) are the same as (a) but now an element is only included if its tipping risk exceeds 10% and 1% probability, respectively. The same global and regional tipping elements are considered as in figure 2. Each element is coloured and marked as given in the labels to the right side of the panels. Colours are selected such that red colours represent fast, and blue colours represent slow tipping timescales. Adapted from Ritchie *et al* (2025b) CC BY 4.0.

poorly constrained, with only a few modelling studies available (Mengel *et al* 2018, Bauer *et al* 2023). As a ‘slow’ tipping element, the full consequences

of past and ongoing warming may take centuries or longer to unfold (Clark *et al* 2016, Klose *et al* 2024a). Nevertheless, Antarctica is losing mass and

contributing to sea-level rise already today, with losses projected to accelerate even if global temperatures were stabilised at current levels (Reese *et al* 2023). The WAIS, in particular, is highly vulnerable due to its marine-based configuration (meaning that the ice sheet rests on bedrock submerged below the sea level), making it susceptible to runaway retreat through marine ice sheet instability (Garbe *et al* 2020). This causes the WAIS to have a particularly low tipping point, estimated at about 1 °C–3 °C of warming above pre-industrial levels (Pattyn *et al* 2018, Garbe *et al* 2020, Armstrong McKay *et al* 2022). In fact, models suggest that even under present observed warming levels, a partial WAIS collapse may already be unavoidable in the long term (Reese *et al* 2023, Chandler *et al* 2025, van den Akker *et al* 2025), though its onset could range from centuries to millennia. Ice-shelf disintegration driven by oceanic and atmospheric warming could further accelerate ice loss, compounding global sea-level rise. Some models suggest this could trigger an even more dangerous instability cascade within the ice sheet itself (DeConto *et al* 2021), though the precise mechanisms remain debated. Consequently, overshooting global warming targets, even temporarily, severely increases the risk of transgressing tipping points in Antarctica (Stokes *et al* 2025).

Marine Basins East Antarctica: The marine basins of East Antarctica hold ice equivalent to about 19.2 metres of potential sea-level rise and have been identified as a ‘global core’ climate tipping element (Lenton *et al* 2023). However, critical warming thresholds vary strongly between individual basins: The lowest tipping point in East Antarctica is found for the Wilkes Subglacial Basin and is estimated at around 2 °C–3 °C above pre-industrial levels based on modelling and paleo reconstructions (Garbe *et al* 2020, Armstrong McKay *et al* 2022, Lenton *et al* 2023). Paleoclimate data reveals that past warm periods with global temperatures as low as 1 °C–2 °C above pre-industrial levels led to significant ice retreat in this basin (Wilson *et al* 2018, Blackburn *et al* 2020). Other East Antarctic marine ice basins, such as the Aurora and Recovery Subglacial Basins, likely have higher thresholds at around 5 °C–8 °C and 6 °C–8 °C of warming, respectively (Garbe *et al* 2020, Armstrong McKay *et al* 2022, Lenton *et al* 2023). If tipping points for these basins are crossed, self-reinforcing feedbacks such as marine ice sheet instability, reduced ice shelf buttressing, and basal thawing, which together can trigger irreversible ice retreat and long-term sea-level rise (Garbe *et al* 2020, Abram *et al* 2025). In the scenario of a temperature overshoot exceeding these tipping points even temporarily, the marine basins could undergo self-sustaining retreat with substantial ice loss, committing these basins to long-term mass loss and sea-level rise, despite subsequent climate cooling efforts. Such retreat would unfold over time scales

ranging from centuries to millennia, with irreversible impacts due to hysteresis, which would require cooling far below pre-industrial levels to be reversed (Mengel and Levermann 2014, Garbe *et al* 2020).

Mountain glaciers: Mountain glaciers have been identified as a potential tipping element (e.g. Armstrong McKay *et al* 2022), though the response of mountain glaciers to global warming and potential overshoot are not uniform globally and depend on region and the individual glaciers’ sensitivity to temperature change. Like the polar ice sheets, mountain glaciers do not respond immediately to changes in climate, but rather the committed volume loss due to warming manifests over decades to centuries following the change in temperature forcing (Zekollari *et al* 2025). Few studies have investigated the response of mountain glaciers to overshoot and thus there remains considerable uncertainty in the timing and rate of mountain glacier mass loss and potential recovery. Despite their typically-linear response to warming (Rounce *et al* 2023), the few available studies demonstrate that slowly-responding glaciers will still undergo additional long-lasting glacier mass loss with overshoot than if temperatures were stabilised to the same level without a period of overshoot (Schuster *et al* 2025). Fast-responding glaciers, such as those in steeper regions (Zekollari *et al* 2020), may exhibit partial regrowth following overshoot.

AMOC: A collapse of the AMOC has been shown by models across the hierarchy, from conceptual models (Stommel 1961) to state-of-the-art ESMs (Romanou *et al* 2023, Van Westen *et al* 2024) and usually takes more than a century to occur. The main positive feedback leading to this tipping behaviour is related to the salt-advection feedback (Stommel 1961, Van Westen *et al* 2024). The AMOC’s response to overshoot scenarios can be thoroughly understood in conceptual models (e.g. Alkhayon *et al* 2019, Chapman *et al* 2024) and explored in ESMs using extreme freshwater or temperature forcing profiles (e.g. Hawkins *et al* 2011, Jackson *et al* 2023). However, the tipping point for the AMOC has the largest uncertainty range of all tipping elements (Armstrong McKay *et al* 2022, Lenton *et al* 2023) and therefore determining overshoot profiles that avoid tipping the AMOC is challenging. This is due to the large uncertainty in AMOC projections, with reduction going from 3%–72% in 2100 mainly depending on the CMIP6 model considered. As such, coordinated experiments are useful to assess the AMOC dynamics (and notably important is the strength of the salt advection feedback) in various models, in response to the same surface perturbations (contrary to emission scenarios where surface forcing of the AMOC can be very different among models). In this respect, the TIPMIP initiative (Winkelmann *et al* 2025) is welcome to better understand differences in models. Improvements

in the understanding of key oceanic processes that influence the AMOC like eddies (Hirschi *et al* 2020) and salt fluxes (Faure Ragani and Dijkstra 2025), combined with the observational-constraints framework (Portmann *et al* 2025), allows to reduce uncertainty concerning its tipping point and characteristics to better assess tipping risk related to overshooting scenarios.

North Atlantic Subpolar Gyre convection (SPG):

The SPG is a subset of the larger-scale AMOC, focusing on the subpolar gyre region, south of Iceland. This specific circulation has been suggested to have a bistability regime based on conceptual (Born and Stocker 2014) and EMIC (Levermann and Born 2007) model results. Moreover, in CMIP models multiple abrupt transitions in the SPG have been observed (Sgubin *et al* 2017, Swingedouw *et al* 2021). The mechanism underlying the tipping point in sea surface temperature, salinity, and deep convection may be linked to a critical surface salinity threshold. As surface salinity decreases, stratification strengthens, and below a certain threshold it becomes too strong to permit deep convection, even when surface temperatures decline and surface density increases. In this regime, the phase-change threshold limits further cooling: once temperatures reach the freezing point, additional heat loss leads instead to sea-ice formation rather than continued cooling. This inhibits further density increase and ultimately suppresses convection, which can be seen as a tipping point for the SPG (Swingedouw *et al* 2020, Wood *et al* 2024). The time scale of a deep convection collapse in the SPG is about one decade, making this tipping element faster than an AMOC collapse. As such, it might be very sensitive even to short-term overshoot. The response of the SPG strongly depends on the model considered, so the implications of an overshoot on the risk of collapse have not been properly assessed up to now. However, some models do show a collapse of the deep convection in the SPG for low-emission scenarios like SSP1-2.6, raising concerns for neighbouring regions like the UK and western Europe (Sgubin *et al* 2019). To improve estimates of the proximity to a deep convection collapse in the SPG linked with potential global temperature overshoots, improvements in deep convection processes and the spread of freshwater from the borders of the SPG (related to Greenland melting for instance) to the centre might be crucial to improve (e.g. Swingedouw *et al* 2022).

Amazon rainforest: The potential of climate change to cause an Amazon rainforest tipping from a closed-canopy, humid tropical forest system to an open-canopy degraded ecosystem remains an iconic risk associated with rising atmospheric greenhouse gases (Flores *et al* 2024, Brando *et al* 2025, Melnikova *et al* 2025). This leads to regional climate change and water stress to the forest system, first identified by

Friend *et al* (1997) and Cox *et al* (2000) but also in subsequent literature (Cox *et al* 2004, Malhi *et al* 2009, Nobre *et al* 2016, Parry *et al* 2022, Flores *et al* 2024). In addition to climate change alone, direct deforestation and forest degradation (e.g. from fires) also pressure the Amazon rainforest. When combined, these pressures interact, reducing the overall resilience and lowering the tipping point, potentially anticipating tipping from one adverse influence alone (Boers *et al* 2017, Lovejoy and Nobre 2019, Bochow and Boers 2023, Flores *et al* 2024). This implies that all efforts are needed for the prevention of Amazon rainforest tipping, potentially including whether deliberate reforestation could adjust the Amazon rainforest's tipping point, or create inertial effects, such that the forest would be more resilient to any imposed decadal or century-timescale temperature overshoot. Following a conceptual analysis Ritchie *et al* (2021) estimate that the Amazon rainforest may be resilient if exceedance of its warming tipping point was limited to only a few decades. However, as of yet, a dedicated study on temporary exceedance of the Amazon rainforest tipping point is still lacking. Frequent extreme droughts, such as in the years of 2005, 2010, 2015/2016 and 2023/2024 (Lewis *et al* 2011, Jiménez-Muñoz *et al* 2016, Espinoza *et al* 2024, Marengo *et al* 2024), may provide empirical evidence of how forests may respond in the future. Trees can withstand drier conditions for a certain period of time, but there are limits to adaptation (Wunderling *et al* 2022). Better ecological understanding will reveal if a temperature overshoot pathway associated with a temporary dry period in the Amazon rainforest can be withstood or not. Initial studies suggest that once a temperature overshoot period ceases, regional rainfall patterns are expected to experience lasting changes through atmosphere-vegetation feedbacks, as reduced forest cover weakens evapotranspiration and moisture recycling, leading to either higher or lower rainfall levels depending on the region (Kim *et al* 2022, Oh *et al* 2022, Schleussner *et al* 2024).

Boreal forests: Boreal forests, one of the largest terrestrial biomes on Earth, store about a third of the global terrestrial carbon above and below ground (Bradshaw and Warkentin 2015) and destabilisation could result in additional global warming due to the release of greenhouse gases (Armstrong McKay *et al* 2022). The biome does not exhibit a gradual decline in tree cover towards the ecological limits, but demonstrates high susceptibility to abrupt shifts between stable but competing forest, grassland-steppe, and shrubland states with differing effects on surface albedo and disturbance regimes (Scheffer *et al* 2012, Gauthier *et al* 2015). Two primary tipping processes influence boreal forest dynamics and the associated carbon sequestration rates. Firstly, southern dieback is driven by warming-induced drought stress,

coupled with increased fire frequency, lower pest resilience, and limited seed availability, which constrains regrowth, leading to rapid shifts in forest composition and density, and facilitating the expansion of steppe or temperate forests at regional to sub-continental spatial scales (Rao *et al* 2023). On the other hand, northern forest expansion into tundra areas is expected due to longer and warmer growing seasons and would reduce winter surface albedo, potentially reinforcing warming effects. Recent evidence suggests that this shift is much slower than the southern steppe transition but that improved growing conditions have increased the productivity in the northern boreal forests (Berner and Goetz 2022, Rotbarth *et al* 2023). Therefore, for a temperature overshoot, the risk of boreal forest dieback in the south is greater than boreal forest expansion in the north (assuming the same tipping point). While boreal forest loss is generally reversible on centennial timescales, nonlinear biome shift at the southern margin, to woodland or treeless states, alter the ecosystem state at regional to sub-continental spatial scales. A temperature overshoot could in turn lead to the loss of ecosystem-protected and -modified permafrost, which exhibits a strong coupling between forest cover and active layer dynamics, particularly in terms of water availability (Stuenzi *et al* 2022). The associated permafrost thaw and ground ice melt is not reversible within centennial timescales.

Warm-water coral reefs: Warm-water coral reefs, which play a critical role in supporting marine biodiversity and human livelihoods, are among the most vulnerable ecosystems to anthropogenic driven climate change, with a tipping point for collapse closely tied to rising global temperatures and CO₂ concentrations. Collapse is defined as having insufficient living cover (typically <10%) and diversity of hard corals to support the wide diversity of species and ecological interactions typical of a coral reef (Lenton *et al* 2023). Stressors such as rising sea surface temperatures (i.e. more severe and frequent marine heatwaves), ocean acidification, overfishing and pollution are the primary drivers of potentially irreversible transformations (Pearce-Kelly *et al* 2025). The likelihood of reaching these tipping points is high, as many reefs already experience conditions beyond their thermal tolerance (Muñiz-Castillo *et al* 2019, IPCC AR6 WGII-SPM 2022, Cooley *et al* 2023, Lenton *et al* 2023, Goreau and Hayes 2024, Klein *et al* 2024, Reimer *et al* 2024, Byrne *et al* 2025). A global mean surface temperature increase of 1.2 °C above pre-industrial levels (range 1 °C–1.5 °C) and atmospheric CO₂ concentrations exceeding 350 ppm are recognised as key tipping points for mass bleaching events and coral reef collapse that would lead to biodiversity and ecosystem services loss (Veron *et al* 2009, Hansen *et al* 2013, IPCC AR6 WGII-SPM 2022, Lenton *et al* 2023, Pearce-Kelly *et al* 2025).

Previous generally acknowledged estimates for temperature driven coral reef loss is 70%–90% loss at 1.5 °C warming and 99% loss at 2 °C (Frieler *et al* 2013, Schleussner *et al* 2016, Hoegh-Guldberg *et al* 2018, Díaz *et al* 2019, Souter *et al* 2021, Armstrong McKay *et al* 2022, Reimer *et al* 2024). Finer-scale modelling projects a 100% loss at 2.0 °C (Dixon *et al* 2022, Kalmus *et al* 2022). Moreover, co-occurring stressors (Setter *et al* 2022), such as ocean acidification, pollution, storms and disease, together with cascading Earth system and ecological impacts, may further lower these tipping points, accelerating ecosystem collapse (Pearce-Kelly *et al* 2025). Corals have very fast response timescales (mortality may play out over weeks to months for thermal stress, or years for chronic threats such as disease and pollution, but prolonged failure to recover over a decade or more is necessary to qualify as collapsed), suggesting anything other than the briefest overshoot period is likely to cause irreversible damage. Recovery from such tipping events is (partially) possible (Donner and Carilli 2019, Devlin 2022, Anderson 2025) but challenging due to slowed coral recruitment and calcification rates following repeated disturbances. A reliance on climate resilient corals and the existence of connected coral refuges offer the greatest hope for these critical ecosystems not to die-out completely (Camp *et al* 2018, Bartels *et al* 2023, McClanahan *et al* 2024, Dubé *et al* 2025, Greiner *et al* 2025). Ongoing mass coral mortality events increasingly compromise recovery potential of corals. This creates hysteresis, where reefs fail to recover even if environmental conditions improve. Hence, when emission-driven temperature overshoot is considered, lower target temperatures with a temporary overshoot can mean little difference in the coral survival compared to stabilisation scenarios at higher temperatures (Tachiiri *et al* 2019).

Land permafrost: Overshooting temperature targets presents a substantial risk to permafrost stability and its climate impacts due to its potential to release vast amounts of greenhouse gases from the large quantities of carbon in the form of frozen soil organic matter (Schuur *et al* 2022) in an almost irreversible manner. While permafrost thaw is characterised by multiple regional-scale tipping processes, including abrupt thermokarst lake formation and slope slumping, these processes do not aggregate to a single globally coherent tipping point (Nitzbon *et al* 2024). Therefore, permafrost is likely not a tipping element on a global scale. This means that, in contrast to other tipping elements, permafrost thaw commences with every increment of global warming and therefore there is no maximal warming limit up to which climate change impacts are minimal to permafrost (Nitzbon *et al* 2024). The severity of permafrost degradation under overshoot depends on both the peak and duration of the overshoot, i.e. cumulative

heat exposure (Gasser *et al* 2018, Schleussner *et al* 2024). Even if an overshoot is temporary, heat can penetrate deep into the soil, causing continued permafrost degradation. While some may refreeze if temperatures decline after the warming peak, much of the carbon loss is likely irreversible due to the slow formation timescales of permafrost. Even if temperature levels eventually decline, sustained microbial decomposition of previously frozen organic matter causes continued carbon emissions over centennial to millennial timescales and reinforces warming through a self-perpetuating, although relatively weak, positive feedback loop (de Vrese and Brovkin 2021, Schwinger *et al* 2022b, Ji *et al* 2025). Furthermore, permafrost thaw does not occur uniformly, as some areas experience localised abrupt thaw, leading to rapid carbon loss and landscape destabilisation, i.e., degradation of ice-rich permafrost, and subsequent rapid ground subsidence and the formation of thermokarst landscapes (Turetsky *et al* 2020, Park *et al* 2025). Temperature overshoot enhances the risk of causing localised abrupt thaw. Overshooting temperature targets also increases the risk of tipping cascades, where permafrost degradation—in addition to its carbon-climate feedback—interacts with other Earth system components, such as boreal forest dieback (Alfaro-Sánchez *et al* 2024), including wildfire occurrence (Kim *et al* 2024) or a modified ocean circulation (Schwinger *et al* 2022a, Park *et al* 2025), potentially accelerating global climate change.

5. Additional pressures on tipping elements

In addition to direct global warming impacts, further pressures such as direct anthropogenic interference, tipping element interactions and Earth system feedbacks may threaten the stability of tipping elements, and the feasibility of realising/maintaining stabilisation target levels. This means that global warming tipping points may effectively be lower than their reported value in the recent literature (e.g. Armstrong McKay *et al* 2022, Lenton *et al* 2023) where they are studied in isolation, consequently reducing the allowable peak and duration for a temperature overshoot that does not cause tipping.

Additional anthropogenic pressures: Ecological tipping elements suffer from additional anthropogenic interference such as deforestation in the Amazon forest, or overfishing, ocean acidification and pollution in warm-water coral reefs, further reducing their resilience. Comparable pressures are evident in the cryosphere: soot and black carbon deposition on snow and ice accelerate melting, while infrastructure development in permafrost regions destabilises ground systems and releases greenhouse gases; and in the atmosphere: emissions of aerosols and short-lived climate forcers (like methane and tropospheric

ozone) alter radiative forcing patterns and regional monsoon dynamics. Combined with global warming, these additional pressures may cause these tipping elements to have their tipping point at effectively lower temperatures than would be the case for global warming pressure alone (e.g. Hughes *et al* 2018, Lovejoy and Nobre 2018, Setter *et al* 2022, Lenton *et al* 2023, Pearce-Kelly *et al* 2025). Or, in turn, opportunities for improved management and local agency offer to limit these pressures and provide increased resilience within these ecological tipping elements.

Interactions between tipping elements: Most direct interactions between tipping elements are assessed as destabilising (Lenton *et al* 2023, Wunderling *et al* 2024). Therefore, interactions between tipping elements exert additional pressure on the resilience of tipping elements and become relevant at 1.5 °C of global warming or higher. This means that increasing temperatures can trigger a first element into tipping that in turn effectively lowers the warming tipping point for subsequent tipping elements via direct physical interactions (Wunderling *et al* 2021a). The AMOC sits centrally in a network of tipping elements and is therefore often viewed as a mediator of a possible cascade of tipping events (Wunderling *et al* 2021a). For instance, a destabilisation of the GrIS could cause a collapse of the AMOC, that could then impact the Amazon rainforest or monsoon systems. In such an accelerating cascade of tipping elements (Ritchie *et al* 2025a), an overshoot that may only temporarily cross the GrISs tipping point and not cause tipping (due to its slow timescale), may still be sufficient to cause tipping of the AMOC as a result of its faster timescale (Klose *et al* 2024b). Conversely, due to the relatively fast timescales of the AMOC, a slowdown could stabilise the slower responding GrIS and lead to a mutual stabilisation of both tipping elements (Pöppelmeier and Stocker 2025). Alternatively, melting from the Antarctic ice sheets could have a stabilising effect on the AMOC (Sinet *et al* 2023, 2025, Knight and Condron 2025).

Earth system feedbacks: Further, there may be Earth system feedbacks that alter the risk for triggering tipping elements. Many tipping elements would cause feedbacks that increase global warming levels if they tip (Armstrong McKay *et al* 2022, Lenton *et al* 2023, Ripple *et al* 2023b, Deutloff *et al* 2025). For instance, the large ice sheets would add up to 0.2 °C of additional global warming if they disintegrate due to the ice-albedo feedback and likewise 0.2 °C for a dieback of the Amazon rainforest because of decreased carbon uptake. Further positive climate feedbacks include cloud feedbacks (Bjorndal *et al* 2020, Ceppi and Nowack 2021) or the permafrost carbon feedback (MacDougall *et al* 2020, Steinert and Sanderson 2025). However, there are also few but

potentially strong negative feedbacks on global temperatures (e.g. 0.5 °C cooling due to AMOC tipping) (Armstrong McKay *et al* 2022). Therefore, the overall effect of temperature feedbacks of disintegrated tipping elements on tipping risks remains uncertain and requires improved constraints of their feedback mechanisms, direction and magnitude (Bdolach *et al* 2025). Furthermore, changes in the land and ocean carbon sinks under ongoing global warming may affect global warming pathways themselves. So far, around 60% of all emissions have been taken up by land and ocean carbon sinks and only the remaining 40% contribute to atmospheric CO₂-increase and global warming (Friedlingstein *et al* 2025). There is now growing evidence of a long-term weakening of land carbon sinks (e.g. in the Amazon rainforest or permafrost (Gatti *et al* 2021, Ke *et al* 2024)). On the other hand, while also weakening on the long term, ocean carbon sinks seem to be comparatively stable across ESMs (Tokarska *et al* 2019, Schwinger *et al* 2022b, Koven *et al* 2023, Sanderson *et al* 2025).

6. Implications of climate tipping risks for mitigation ambition during and after temperature overshoot

Considering these findings, the prospects of temperature overshoot must inform climate mitigation ambition in the near and longer term. Given the potential of initiating tipping processes during overshoot, and the severe risks tipping events present to societies, the ambition reflected in greenhouse gas emission reductions should aim to minimise tipping risks by limiting the peak and duration of temperature overshoot, and bringing temperatures back down to levels well below 1.5 °C.

A race to net-zero CO₂: Limiting overshoot requires mitigation strategies that minimise the global peak temperature, keeping it as close to 1.5 °C as possible, and returning to below 1.5 °C as soon as possible (Palter *et al* 2018, Wunderling *et al* 2023, Schleussner *et al* 2024). Peak temperature and return time are mutually dependent—the higher the peak, the longer the overshoot duration likely becomes. Minimising the global peak temperature (McKenna *et al* 2021) requires immediate peaking of global greenhouse gas emissions and stringent emission reductions thereafter towards global net zero CO₂ emissions by 2050 (IPCC AR6 WGIII-SPM 2022).

Limits and feasibility of minimising overshoot: Substantial net negative emissions are required to achieve a rapid reversal in warming. This is needed to limit the higher risks of crossing tipping points associated with large and long overshoot pathways, given that already under present global warming levels, we enter the uncertainty ranges for triggering several tipping elements such as GIS, WAIS or the subpolar gyre

(Armstrong McKay *et al* 2022, Lenton *et al* 2025). Achieving net negative emissions requires anthropogenic CDR to exceed remaining emissions that cannot be eliminated. However, these must be viewed in light of the very real risks that these interventions pose for marine and terrestrial systems, as well as human wellbeing (Heck *et al* 2016, 2018, Pereira *et al* 2024). CDR refers to a set of methods aimed at actively extracting CO₂ from the atmosphere and durably storing it in terrestrial, geological, or ocean reservoirs. There is an immediate need to invest in and bring to scale sustainable methods of CDR as a complement to—not a substitute for—mitigation to enable future temperature reduction. Achieving net zero greenhouse gas emissions as of Article 4.1 of the Paris Agreement requires CDR deployment to achieve net-negative CO₂ emissions to balance out residual non-CO₂ emissions and may therefore lead to a long-term temperature decline (Fuglestad *et al* 2018, Schleussner *et al* 2022). Achieving a faster and sustained temperature decline may require going even beyond net-zero greenhouse gases.

Current assessments indicate that CDR rates in overshoot 1.5 °C scenarios range from 7 to 10 GtCO₂/yr (Smith *et al* 2024). With a current rate of CDR of 2 GtCO₂/yr from conventional methods such as Afforestation, Forestry and Other Land Use, massive upscaling will be required to achieve the removal rate required to return warming to below 1.5 °C by 2100 (Smith *et al* 2024). These removals would need to be in tandem with reductions in CO₂ emissions to near zero. Even higher rates of CDR would be needed to reduce temperatures below the most sensitive tipping points (at around 1 °C warming). While several technological CDR approaches, such as bioenergy with carbon capture and storage, direct air carbon capture and storage, enhanced weathering, or ocean-based methods, are under active development, most remain at early deployment stages and face considerable challenges in terms of cost, energy demand, infrastructure, and scalability (Smith *et al* 2024) and risks (Heck *et al* 2016). Land-based methods are limited by land availability, ecosystem trade-offs, and saturation effects. Additionally, there are capacity limits for some forms of CDR, which constrain the space for feasible CDR deployment (Gidden *et al* 2025). Consequently, achieving sustained removal rates of several gigatonnes per year would require rapid technological innovation, global policy coordination, and major financial and institutional support (Smith *et al* 2024). Even if CO₂ removal rates required to restore temperature to 1.5 °C after various levels of overshoot prove to be technically feasible, large upscaling of CDR is subject to various sustainability and feasibility constraints, including political preferences and social acceptability (Schleussner *et al* 2024, Smith *et al* 2024).

The reversibility of global temperatures remains uncertain, and achieving net-negative CO₂ emissions

might be required should Earth system feedbacks exceed current expectations, with the implication of potentially greater deployment of CDR (Schleussner *et al* 2024, Kaufhold *et al* 2025). Furthermore, the potential reversibility of global mean temperature does not mean that the impacts of climate change can be reversed. While some impacts may be reversible with lowering global mean temperature, others such as sea level rise, loss of ecosystem functionality, increased risks of species extinction, and glacier and permafrost loss are not reversible on timescales of decades to millennia (Tokarska and Zickfeld 2015, Ehlert and Zickfeld 2018, Drouet *et al* 2021, Bauer *et al* 2023). Further, even if temperature can be reversed on a global scale, there will likely be substantial impacts on regional-to-local scales as consequence of the overshoot (Pfleiderer *et al* 2024, Schleussner *et al* 2024, Steinert *et al* 2025) or of the interventions taken to drawdown the CO₂ that has already been or will continue to be emitted should drastic decarbonisation efforts not materialise (Pereira *et al* 2024). These factors together highlight the need for climate governance to evolve to collectively manage global emissions and temperature pathways rather than focus only on achieving the long-term temperature goal. This pathway focus has direct implementation potential in the ambition cycle of the Paris Agreement, e.g., with explicit considerations in Nationally Determined Contributions, the Global Stocktake, and the Enhanced Transparency Framework.

Long-term temperature stabilisation: The Paris Agreement temperature goal sets upper limits for global mean temperature increase above pre-industrial levels in light of the ultimate Objective of the Climate Convention to avoid ‘dangerous anthropogenic interference’ (United Nations Framework Convention on Climate Change) (UNFCCC 1992). The Paris Agreement climate objectives are designed to achieve long-term temperature decline and do not specify a ‘safe’ level of global temperature stabilisation (Schleussner *et al* 2022). Emerging science on tipping risks indicates that temperature levels well below 1.5 °C, potentially around 1 °C above pre-industrial levels, would be required to limit tipping risks in the long-run and be a more ‘just’ level, limiting harm to people and ecosystems (Breyer *et al* 2023, Rockström *et al* 2023, Gupta *et al* 2024, Möller *et al* 2024). Considering this emerging evidence, the current temperature goal may require amendments e.g., in a future Periodic Review process under the UNFCCC.

7. Conclusion

Recent assessments indicate that current global carbon emissions remain at high levels, making at least

a temporary overshoot of the 1.5 °C target increasingly likely. Combined with evidence that some tipping points may already have been crossed, this trajectory heightens the risk that additional Earth system tipping elements could be pushed beyond their tipping points.

Five, and possibly up to eight, tipping elements of the Earth system may reach their tipping point below 2 °C of global warming, with some tipping points potentially exceeded below 1.5 °C (see figure 2). The risk of tipping for individual elements is found to increase for larger and longer overshoots. Minimising the peak of an overshoot is crucial, but arguably minimising the duration of an overshoot is of even greater importance. However, these characteristics are interrelated—invariably a higher peak temperature will result in a longer overshoot duration. Therefore, limiting peak global warming to as close to 1.5 °C as possible will contribute to ensuring that the overshoot duration is limited too. This will help to avoid as many tipping points as possible from being exceeded, particularly those associated with fast elements (e.g. North Atlantic Subpolar Gyre), where it will be difficult to prevent tipping once the tipping point is exceeded, even if overshoot duration is comparatively short. Keeping the overshoot duration as short as possible and then pursuing efforts to reduce warming to levels of 1 °C and below, will aid in preventing slow elements (e.g. polar ice sheets) from tipping in the long term despite some maybe already having reached their tipping points (in terms of level of warming).

The current theoretical understanding on whether an overshoot profile, characterised by overshoot peak and duration, will trigger tipping for a given element has been derived from a simple dynamical context. Therefore, more research is required to determine how this scales with complexity, for example where the tipping mechanism cannot be represented by passing through a saddle-node bifurcation or if the system is chaotic. Moreover, deriving dynamically simple emulators based on saddle-node bifurcation tipping that resemble more complex dynamics could address this complexity gap. Additionally, global mean temperature is often depicted as the main driver of the tipping elements, arguably for policy relevance. However, many of the elements respond to other quantities such as regional temperatures, including polar amplification that cannot be captured fully by global temperature. Further research on how this affects risk assessments is still required.

Further research is needed to better understand the reversibility of elements if they were to tip. This includes determining how much further global warming must be reduced, compared to the level that triggered tipping in the first instance, to restore an element to its original state. The magnitude and timescale of such hysteresis effects, and how they differ across Earth system tipping elements, are still poorly

constrained. Further, rate-induced tipping requires considerably more research: while rapidly reducing temperatures after an overshoot could shorten the period of high tipping risk, fast changes in forcing may destabilise systems that are sensitive to the rate of change rather than just absolute level. Exploring these interactions systematically across different temperature pathways is crucial for anticipating unintended consequences of both overshoot and rapid-reversal scenarios. More generally, reducing key uncertainties on tipping points, the strength and interactions of Earth system feedbacks, and the value of climate sensitivity remains fundamental for narrowing the range of possible futures and for designing strategies that minimise tipping risks. However, uncertainty can never be fully eliminated and so this calls for a comprehensive risk assessment approach to quantify tipping likelihoods for various temperature overshoot pathways and for changes in climate governance to limit the risks related to Earth tipping processes. Therefore, we envision that the following steps towards a novel risk assessment for tipping elements subjected to overshoot pathways can (and should) be taken:

- 1. Derive simplified process-based dynamics from complex ESMs.** Recent advances in the process-based representation of tipping elements within complex ESMs now enable the extraction of their dynamics in a simplified form. Large-scale modelling efforts such as TIPMIP (Winkelmann *et al* 2025) and related projects, as well as complex system-specific modelling efforts (for example, simulations with ocean models such as Veros (Lohmann *et al* 2024a) or ice sheet models such as PISM (Bochow *et al* 2023)) already provide a solid foundation for this development.
- 2. Integrate process-based tipping dynamics into efficient climate models for development of a probabilistic tipping framework.** These simplified representations can be incorporated into reduced-complexity and computationally efficient modelling tools such as FaIR (Möller *et al* 2024, Deutloff *et al* 2025) and PyCascades (Wunderling *et al* 2021b, Bdolach *et al* 2025). This would allow for large-scale Monte Carlo simulations across a range of overshoot scenarios (such as in Wunderling *et al* 2023, Möller *et al* 2024, Ritchie *et al* 2025b). The resulting distribution of tipping likelihoods would constitute the probabilistic risk assessment for tipping elements. However, unlike earlier approaches, which have their limitations, this more comprehensive framework can provide process-based knowledge from ESMs on characteristics such as the tipping point and the tipping timescale.
- 3. Extend the framework to include spatially explicit tipping dynamics.** In subsequent stages, the simplified tipping dynamics should be enriched by incorporating spatially explicit dynamics. Spatially resolved dynamics are particularly relevant for tipping elements exhibiting basin or region-specific thresholds, such as the Antarctic ice sheet (Garbe *et al* 2020) and the Amazon rainforest (Boulton *et al* 2022), where local feedbacks can govern the onset and propagation of tipping. In this way, the phenomenon of fragmented tipping, where only a part of the spatial domain has transitioned to an alternative state, can be addressed as well (Bastiaansen *et al* 2022).

Achieving the rapid reversal in warming needed to prevent many tipping elements from tipping, requires substantial net negative emissions. The longer CO₂ emission reductions are delayed, the larger net negative emissions must be. Additionally, the larger the overshoot, the larger the need for negative emissions and the implications of those for our planet's critical biophysical systems. Potential pathways that overshoot 1.5 °C and return to this level by 2100 with 50% probability require up to five times the current rates of CDR, which are mainly achieved through conventional methods such as Afforestation, Forestry and Other Land Use (Smith *et al* 2024). These methods, together with alternatives that would be required if upscaling is not possible, carry their own risks and impacts for people and ecosystems (Heck *et al* 2018, Pereira *et al* 2024). Additional feedbacks have the capacity to impede the reversal of warming, such as the potential of permafrost thawing and Amazon rainforest dieback to amplify global warming (e.g. Deutloff *et al* 2025) or the weakening of land carbon sinks under climate change. Therefore, urgent climate action to minimise tipping risks is required, also considering additional anthropogenic pressures, interactions between tipping elements as well as uncertainties in tipping points and system timescales. This means that peak overshoot temperature (below 2.0 °C), overshoot duration (below 1.5 °C by the end of this century), and long-term warming stabilisation (at or below 1.0 °C) must be limited to sufficiently low levels to prevent Earth system elements from tipping in the short and long term.

Data availability statement

The data and the code that support the findings of this study are openly available at the following URL/DOI: URL/DOI: https://github.com/PaulRitchie89/Overshoots_review (Ritchie 2025).

Tipping risk overshoot animation available at <http://doi.org/10.1088/1748-9326/ae3cad/data1>.

Acknowledgments

This review article has been carried out within the framework of the Global Tipping Points Report 2025. P D L R acknowledges support from the European Union's Horizon Europe research and innovation programme under Grant Agreement No. 101137601 (ClimTip): Funded by the European Union. This is ClimTip contribution #137. P D L R, C H, T M L and D S also acknowledge support by the UK Advanced Research and Invention Agency (ARIA) via the project 'AdvanTip' and D S via the Project 'SORTED'. N W is grateful for funding support from the Center for Critical Computational Studies and the Klaus Tschira Foundation (under the Grant Agreement ID 25545). N J S has been supported by the Research Council of Norway through the project NAVIGATE (Grant 352142). D S acknowledges support from the European Union's Horizon Europe research and innovation programme under Grant Agreement No. 101137673 (TipESM) funded by the European Union. J F A acknowledges support from the PREDICT project which has received funding from the European Space Agency (ESA) under ESA Contract No. 4000146344/24/I-LR. J F A also acknowledges support from the Earth Commission, which is hosted by Future Earth. C Q acknowledges support from the Australian Research Council under Discovery Early Career Researcher Award Project Number DE250101025. K L-G acknowledges the Irène Curie fellowship of the Eindhoven University of Technology. T M acknowledges financial support by the Luxembourgish Fond National de la Recherche (FNR) (under the AFR PhD Grant 18824314). N B acknowledges support from the Research Council of Norway (Project Number 314570).

Conflict of interest

The authors declare no competing financial interests.

Author contributions

Paul D L Ritchie  0000-0002-7649-2991
 Conceptualization (equal), Formal analysis (lead),
 Project administration (equal), Supervision (equal),
 Visualization (lead), Writing – original draft (equal),
 Writing – review & editing (lead)

Norman J Steinert  0000-0002-2154-5857
 Conceptualization (equal), Formal
 analysis (supporting), Project
 administration (equal), Supervision (equal),
 Visualization (supporting), Writing – original
 draft (equal), Writing – review & editing (lead)

Jesse F Abrams  0000-0003-0411-8519
 Writing – original draft (equal), Writing – review &
 editing (supporting)

Hassan Alkhayouon  0000-0001-8117-4907
 Writing – original draft (equal), Writing – review &
 editing (supporting)

Constantin W Arnscheidt  0000-0002-0791-1814
 Writing – original draft (equal), Writing – review &
 editing (supporting)

Nils Bochow  0000-0003-4578-5780
 Writing – original draft (equal), Writing – review &
 editing (supporting)

Ruth R Chapman  0000-0002-7927-325X
 Writing – original draft (equal), Writing – review &
 editing (supporting)

Joseph Clarke  0000-0002-6250-1543
 Writing – original draft (equal), Writing – review &
 editing (supporting)

Donovan P Dennis  0000-0002-5423-7703
 Writing – original draft (equal), Writing – review &
 editing (supporting)

Jonathan F Donges  0000-0001-5233-7703
 Writing – original draft (equal), Writing – review &
 editing (supporting)

Bernardo M Flores  0000-0003-4555-5598
 Writing – original draft (equal), Writing – review &
 editing (supporting)

Julius Garbe  0000-0003-3140-3307
 Writing – original draft (equal), Writing – review &
 editing (supporting)

Annika Högner  0000-0002-4178-9664
 Writing – original draft (equal), Writing – review &
 editing (supporting)

Chris Huntingford  0000-0002-5941-7770
 Writing – original draft (equal), Writing – review &
 editing (supporting)

Timothy M Lenton  0000-0002-6725-7498
 Conceptualization (supporting), Writing – original
 draft (equal), Writing – review &
 editing (supporting)

Johannes Lohmann  0000-0002-6323-6243
 Writing – original draft (equal), Writing – review &
 editing (supporting)

Kerstin Lux-Gottschalk  0000-0001-8412-2707
 Writing – original draft (equal), Writing – review &
 editing (supporting)

Manjana Milkoreit  0000-0002-7862-3561
 Writing – original draft (equal), Writing – review &
 editing (supporting)

Tessa Möller  0000-0001-7757-1985
 Writing – original draft (equal), Writing – review &
 editing (supporting)

- Paul Pearce-Kelly  [0000-0003-2356-7061](#)
Writing – original draft (equal), Writing – review & editing (supporting)
- Laura Pereira  [0000-0002-4996-7234](#)
Writing – original draft (equal), Writing – review & editing (supporting)
- Courtney Quinn  [0000-0001-5298-5233](#)
Writing – original draft (equal), Writing – review & editing (supporting)
- Carl-Friedrich Schleussner  [0000-0001-8471-848X](#)
Writing – original draft (equal), Writing – review & editing (supporting)
- Simone M Stuenzi  [0000-0002-6071-289X](#)
Writing – original draft (equal), Writing – review & editing (supporting)
- Didier Swingedouw  [0000-0002-0583-0850](#)
Writing – original draft (equal), Writing – review & editing (supporting)
- Larissa N Van der Laan  [0000-0002-8892-9011](#)
Writing – original draft (equal), Writing – review & editing (supporting)
- Kirsten Zickfeld  [0000-0001-8866-6541](#)
Writing – original draft (equal), Writing – review & editing (supporting)
- Nico Wunderling  [0000-0002-3566-323X](#)
Conceptualization (equal), Formal analysis (lead), Project administration (equal), Supervision (equal), Visualization (lead), Writing – original draft (equal), Writing – review & editing (lead)

References

- Abram N J *et al* 2025 Emerging evidence of abrupt changes in the Antarctic environment *Nature* **644** 621–33
- Abrams J F *et al* 2023 Committed global warming risks triggering multiple climate tipping points *Earth's Future* **11** e2022EF003250
- Alfaro-Sánchez R, Richardson A D, Smith S L, Johnstone J F, Turetsky M R, Cumming S G, Le Moine J M and Baltzer J L 2024 Permafrost instability negates the positive impact of warming temperatures on boreal radial growth *Proc. Natl Acad. Sci.* **121** e2411721121
- Alkhayuon H M and Ashwin P 2018 Rate-induced tipping from periodic attractors: partial tipping and connecting orbits *Chaos Interdiscip. J. Nonlinear Sci.* **28** 033608
- Alkhayuon H, Ashwin P, Jackson L C, Quinn C and Wood R A 2019 Basin bifurcations, oscillatory instability and rate-induced thresholds for Atlantic meridional overturning circulation in a global oceanic box model *Proc. R. Soc. A* **475** 20190051
- Alkhayuon H, Tyson R C and Wiczorek S 2021 Phase tipping: how cyclic ecosystems respond to contemporary climate *Proc. R. Soc. A* **477** 20210059
- Allen M R, Frame D J, Huntingford C, Jones C D, Lowe J A, Meinshausen M and Meinshausen N 2009 Warming caused by cumulative carbon emissions towards the trillionth tonne *Nature* **458** 1163–6
- Anderson G D 2025 Recovery of coral cover at Lizard island Australia 6 years post-disturbance *Front. Mar. Sci.* **11** 1509455
- Armstrong Mckay D I, Staal A, Abrams J F, Winkelmann R, Sakschewski B, Loriani S, Fetzer I, Cornell S E, Rockström J and Lenton T M 2022 Exceeding 1.5 C global warming could trigger multiple climate tipping points *Science* **377** eabn7950
- Ashwin P, Bastiaansen R, von der Heydt A S and Ritchie P 2025 Early warning skill, extrapolation and tipping for accelerating cascades *Proc. A* **481** 20250405
- Ashwin P and Newman J 2021 Physical invariant measures and tipping probabilities for chaotic attractors of asymptotically autonomous systems *Eur. Phys. J. Special Top.* **230** 3235–48
- Bartels N, Dilernia N J, Howlett L and Camp E F 2023 Stress event for “super corals” in Great Barrier Reef mangrove lagoon *Mar. Biodiver.* **53** 64
- Bastiaansen R, Ashwin P and von der Heydt A S 2023 Climate response and sensitivity: time scales and late tipping points *Proc. R. Soc. A* **479** 20220483
- Bastiaansen R, Dijkstra H A and von der Heydt A S 2022 Fragmented tipping in a spatially heterogeneous world *Environ. Res. Lett.* **17** 045006
- Bauer N *et al* 2023 Exploring risks and benefits of overshooting a 1.5 C carbon budget over space and time *Environ. Res. Lett.* **18** 054015
- Bdolach T, Kurths J and Yanchuk S 2025 Tipping in an adaptive climate network model *Chaos Interdiscip. J. Nonlinear Sci.* **35** 053157
- Ben-Yami M, Morr A, Bathiany S and Boers N 2024 Uncertainties too large to predict tipping times of major Earth system components from historical data *Sci. Adv.* **10** eadl4841
- Ben-Yami M, Skiba V, Bathiany S and Boers N 2023 Uncertainties in critical slowing down indicators of observation-based fingerprints of the Atlantic overturning circulation *Nat. Commun.* **14** 8344
- Berner L T and Goetz S J 2022 Satellite observations document trends consistent with a boreal forest biome shift *Glob. Change Biol.* **28** 3275–92
- Beusch L, Nicholls Z, Gudmundsson L, Hauser M, Meinshausen M and Seneviratne S I 2021 From emission scenarios to spatially resolved projections with a chain of computationally efficient emulators: MAGICC (v7.5.1)–MESMER (v0.8.1) coupling *Geosci. Model. Dev. Discuss.* **2021** 1–26
- Bevacqua E, Schleussner C F and Zscheischler J 2025 A year above 1.5° C signals that Earth is most probably within the 20-year period that will reach the Paris Agreement limit *Nat. Clim. Change* **15** 1–4
- Bjordal J, Storelvmo T, Alterskjær K and Carlsen T 2020 Equilibrium climate sensitivity above 5 °C plausible due to state-dependent cloud feedback *Nat. Geosci.* **13** 718–21
- Blackburn T, Edwards G H, Tulaczyk S, Scudder M, Piccione G, Hallet B, McLean N, Zachos J C, Cheney B and Babbe J T 2020 Ice retreat in Wilkes Basin of East Antarctica during a warm interglacial *Nature* **583** 554–9
- Bochow N and Boers N 2023 The South American monsoon approaches a critical transition in response to deforestation *Sci. Adv.* **9** eadd9973
- Bochow N, Poltronieri A, Robinson A, Montoya M, Rypdal M and Boers N 2023 Overshooting the critical threshold for the Greenland ice sheet *Nature* **622** 528–36
- Boers N 2021 Observation-based early-warning signals for a collapse of the Atlantic Meridional overturning circulation *Nat. Clim. Change* **11** 680–8
- Boers N, Liu T, Bathiany S, Ben-Yami M, Blaschke L L, Bochow N and Smith T 2025 Destabilization of Earth system tipping elements *Nat. Geosci.* **18** 1–12
- Boers N, Marwan N, Barbosa H M J and Kurths J 2017 A deforestation-induced tipping point for the South American monsoon system *Sci. Rep.* **7** 41489
- Boers N and Rypdal M 2021 Critical slowing down suggests that the western Greenland Ice Sheet is close to a tipping point *Proc. Natl Acad. Sci.* **118** e2024192118
- Born A and Stocker T F 2014 Two stable equilibria of the Atlantic subpolar gyre *J. Phys. Oceanogr.* **44** 246–64

- Boulton C A, Lenton T M and Boers N 2022 Pronounced loss of Amazon rainforest resilience since the early 2000s *Nat. Clim. Change* **12** 271–8
- Bradshaw C J and Warkentin I G 2015 Global estimates of boreal forest carbon stocks and flux *Glob. Planet Change* **128** 24–30
- Brando P M et al 2025 Tipping points of Amazonian forests: beyond Myths and toward solutions *Annu. Rev. Environ. Resour.* **50** 97–131
- Breyer C, Keiner D, Abbott B W, Bamber J L, Creutzig F, Gerhards C, Mühlbauer A, Nemet G F and Terli Ö 2023 Proposing a 1.0 C climate target for a safer future *PLoS Clim.* **2** e0000234
- Budd C and Kuske R 2024 Dynamic tipping and cyclic folds, in a one-dimensional non-smooth dynamical system linked to climate models *Physica D* **457** 133949
- Bustamante M et al 2023 Ten new insights in climate science 2023 *Glob. Sustain.* **7** e19
- Byrne M et al 2025 Catastrophic bleaching in protected reefs of the Southern Great Barrier Reef *Limnol. Oceanogr. Lett.* **10** 340–8
- Camp E E, Schoepf V, Mumby P J, Hardtke L A, Rodolfo-Metalpa R, Smith D J and Suggett D J 2018 The future of coral reefs subject to rapid climate change: lessons from natural extreme environments *Front. Mar. Sci.* **5** 4
- IPCC AR6 WGI Canadell J G et al 2021 Global carbon and other biogeochemical cycles and feedbacks *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* ed V Masson-Delmotte et al (Cambridge University Press) ch 5, pp 673–816
- Ceppi P and Nowack P 2021 Observational evidence that cloud feedback amplifies global warming *PNAS* **118** e2026290118
- Chandler D M, Langebroek P M, Reese R, Albrecht T, Garbe J and Winkelmann R 2025 Antarctic Ice Sheet tipping in the last 800,000 years warns of future ice loss *Commun. Earth Environ.* **6** 420
- Chapman R, Sinet S and Ritchie P D 2024 Tipping mechanisms in a conceptual model of the Atlantic meridional overturning circulation *Weather* **79** 316–23
- Clark P U et al 2016 Consequences of twenty-first-century policy for multi-millennial climate and sea-level change *Nat. Clim. Change* **6** 360–9
- Climate Action Tracker 2025 Warming projections global update. Little change in warming outlook for four years; new 2035 climate targets make no difference (available at: <https://climateactiontracker.org/publications/warming-projections-global-update-2025/>) (Accessed 20 November 2025)
- Cooley S, Schoeman D, Bopp L, Boyd P, Donner S, Kiessling W and Simmons C T 2023 *Oceans and Coastal Ecosystems and Their Services. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* ed H-O Pörtner et al (Cambridge University Press) pp 379–550
- Corner S et al 2023 The zero emissions commitment and climate stabilization *Front. Sci.* **1** 1170744
- Cox P M, Betts R A, Collins M, Harris P P, Huntingford C and Jones C D 2004 Amazonian forest dieback under climate-carbon cycle projections for the 21st century *Theor. Appl. Climatol.* **78** 137–56
- Cox P M, Betts R A, Jones C D, Spall S A and Totterdell I J 2000 Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model *Nature* **408** 184–7
- de Vrese P and Brovkin V 2021 Timescales of the permafrost carbon cycle and legacy effects of temperature overshoot scenarios *Nat. Commun.* **12** 2688
- DeConato R M et al 2021 The Paris Climate Agreement and future sea-level rise from Antarctica *Nature* **593** 83–89
- Dekker M M, Von Der Heydt A S and Dijkstra H A 2018 Cascading transitions in the climate system *Earth Syst. Dyn.* **9** 1243–60
- Deutloff J, Held H and Lenton T M 2025 High probability of triggering climate tipping points under current policies modestly amplified by Amazon dieback and permafrost thaw *Earth Syst. Dyn.* **16** 565–83
- Devlin M J 2022 Coral reefs: the good and not so good news with future bright and dark spots for coral reefs through climate change *Glob. Change Biol.* **28** 4506
- Díaz S, Settele J, Brondizio E S, Ngo H T, Guèze M, Agard J and Zayas C 2019 Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental science-policy platform on biodiversity and ecosystem services *Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*
- Dijkstra H A 2013 *Nonlinear Climate Dynamics* (Cambridge University Press)
- Dijkstra H A 2024 The role of conceptual models in climate research *Physica D* **457** 133984
- Ditlevsen P and Ditlevsen S 2023 Warning of a forthcoming collapse of the Atlantic meridional overturning circulation *Nat. Commun.* **14** 1–12
- Dixon A M, Forster P M, Heron S F, Stoner A M K, Beger M and Storto A 2022 Future loss of local-scale thermal refugia in coral reef ecosystems *PLoS Clim.* **1** e0000004
- Donner S D and Carilli J 2019 Resilience of Central Pacific reefs subject to frequent heat stress and human disturbance *Sci. Rep.* **9** 3484
- Drouet L et al 2021 Net zero-emission pathways reduce the physical and economic risks of climate change *Nat. Clim. Change* **11** 1070–6
- Dubé C E, Pogoreutz C, Reverter M, Boitel C, Mills S C, Aqua A and Brahmi C 2025 Bleaching of reef organisms: current knowledge, challenges, and future directions into the Anthropocene *Future Coral Reefs* **21** 41–72
- Ehlert D and Zickfeld K 2018 Irreversible ocean thermal expansion under carbon dioxide removal *Earth Syst. Dyn.* **9** 197–210
- Espinoza J-C, Jimenez J C, Marengo J A, Schongart J, Ronchail J, Lavado-Casimiro W and Ribeiro J V M 2024 The new record of drought and warmth in the Amazon in 2023 related to regional and global climatic features *Sci. Rep.* **14** 8107
- Eyring V, Bony S, Meehl G A, Senior C A, Stevens B, Stouffer R J and Taylor K E 2016 Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization *Geosci. Model. Dev.* **9** 1937–58
- Falkena S K and von der Heydt A S 2024 Subpolar gyre variability in CMIP6 models: is there a mechanism for bistability? (arXiv:2408.16541)
- Faure Ragani A and Dijkstra H A 2025 Physical characterization of the boundary separating safe and unsafe AMOC overshoot behavior *Earth Syst. Dyn.* **16** 1287–301
- IPCC AR5 WGI Flato G et al 2013 Evaluation of climate models *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker et al (Cambridge University Press) ch 9, pp 741–866
- Flores B M et al 2024 Critical transitions in the Amazon forest system *Nature* **626** 555–64
- Forster P M et al 2025 Indicators of Global climate change 2024: annual update of key indicators of the state of the climate system and human influence *Earth Syst. Sci. Data Discuss.* **17** 1–72
- Friedlingstein P, O'sullivan M, Jones M W, Andrew R M, Hauck J, Landschützer P and Zeng J 2025 Global carbon budget 2024 *Earth Syst. Sci. Data* **17** 965–1039
- Frieler K, Meinshausen M, Golly A, Mengel M, Lebek K, Donner S D and Hoegh-Guldberg O 2013 Limiting global warming to 2 C is unlikely to save most coral reefs *Nat. Clim. Change* **3** 165–70
- Friend A D, Stevens A K, Knox R G and Cannell M G R 1997 A process-based, terrestrial biosphere model of ecosystem dynamics (Hybrid v3. 0) *Ecol. Model.* **95** 249–87
- Fuglestedt J, Rogelj J, Millar R J, Allen M, Boucher O, Cain M, Forster P M, Kriegler E and Shindell D 2018 Implications of

- possible interpretations of ‘greenhouse gas balance’ in the Paris Agreement *Phil. Trans. R. Soc. A* **376** 20160445
- Garbe J, Albrecht T, Levermann A, Donges J F and Winkelmann R 2020 The hysteresis of the Antarctic ice sheet *Nature* **585** 538–44
- Gasser T, Kechiar M, Ciaia P, Burke E J, Kleinen T, Zhu D, Huang Y, Ekici A and Obersteiner M 2018 Path-dependent reductions in CO₂ emission budgets caused by permafrost carbon release *Nat. Geosci.* **11** 830–5
- Gatti L V et al 2021 Amazonia as a carbon source linked to deforestation and climate change *Nature* **595** 388–93
- Gauthier S, Bernier P, Kuuluvainen T, Shvidenko A Z and Schepaschenko D G 2015 Boreal forest health and global change *Science* **349** 819–22
- Gidden M J et al 2025 A prudent planetary limit for geologic carbon storage *Nature* **645** 124–32
- Goreau T J and Hayes R L 2024 2023 Record marine heat waves: coral reef bleaching HotSpot maps reveal global sea surface temperature extremes, coral mortality, and ocean circulation changes *Oxford Open Clim. Change* **4** kgae005
- Gottwald G A and Gugole F 2020 Detecting regime transitions in time series using dynamic mode decomposition *J. Stat. Mech.* **179** 1028–45
- Gregory J M, George S E and Smith R S 2020 Large and irreversible future decline of the Greenland ice sheet *The Cryosphere* **14** 4299–322
- Greiner A, Darling E S, Wenger A, Krkošek M and Fortin M J 2025 Identifying priority stepping stone reefs to maintain global networks of connected coral reefs *bioRxiv* 2025–08
- Gupta J et al 2024 A just world on a safe planet: a lancet planetary health–earth commission report on earth-system boundaries, translations, and transformations *Lancet Planet Health* **8** e813–e873
- Hansen J et al 2013 Assessing “dangerous climate change”: required reduction of carbon emissions to protect young people, future generations and nature *PLoS One* **8** e81648
- Hawkins E, Smith R S, Allison L C, Gregory J M, Woollings T J, Pohlmann H and De Cuevas B 2011 Bistability of the Atlantic overturning circulation in a global climate model and links to ocean freshwater transport *Geophys. Res. Lett.* **38** L10605
- Heck V, Donges J F and Lucht W 2016 Collateral transgression of planetary boundaries due to climate engineering by terrestrial carbon dioxide removal *Earth Syst. Dyn.* **7** 783–96
- Heck V, Gerten D, Lucht W and Popp A 2018 Biomass-based negative emissions difficult to reconcile with planetary boundaries *Nat. Clim. Change* **8** 151–5
- Hirschi J J-M et al 2020 The Atlantic meridional overturning circulation in high-resolution models *J. Geophys. Res. Oceans* **125** e2019JC015522
- Hoegh-Guldberg O, Kennedy E V, Beyer H L, McClennen C and Possingham H P 2018 Securing a long-term future for coral reefs *Trends. Ecol. Evol.* **33** 936–44
- Höning D, Willeit M, Calov R, Klemann V, Bagge M and Ganopolski A 2023 Multistability and transient response of the Greenland ice sheet to anthropogenic CO₂ emissions *Geophys. Res. Lett.* **50** e2022GL101827
- Höning D, Willeit M and Ganopolski A 2024 Reversibility of Greenland ice sheet mass loss under artificial carbon dioxide removal scenarios *Environ. Res. Lett.* **19** 024038
- Hourdin F et al 2017 The art and science of climate model tuning *Bull. Am. Meteorol. Soc.* **98** 589–602
- Huang Y, Bathiany S, Ashwin P and Boers N 2024 Deep learning for predicting rate-induced tipping *Nat. Mach. Intell.* **6** 1556–65
- Hughes T P et al 2018 Global warming transforms coral reef assemblages *Nature* **556** 492–6
- IPCC AR6 WGI-SPM 2021 Summary for Policymakers *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* ed V Masson-Delmotte et al (Cambridge University Press) pp 3–32
- IPCC AR6 WGII-SPM 2022 Summary for Policymakers *Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* ed H-O Pörtner et al (Cambridge University Press) pp 3–33
- IPCC AR6 WGIII-SPM 2022 Summary for Policymakers. In: *Climate Change 2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* ed P R Shukla et al (Cambridge University Press) (<https://doi.org/10.1017/9781009157926.001>)
- IPCC SR1.5-SPM 2018 Summary for Policymakers *Global Warming of 1.5 °C. An IPCC Special Report on the Impacts of Global Warming of 1.5 °C above Pre-industrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty* ed V Masson-Delmotte et al (Cambridge University Press) pp 3–24
- Jackson L C, Hewitt H T, Bruciaferri D, Calvert D, Graham T, Guiavarc’h C, Menary M B, New A L, Roberts M and Storkey D 2023 Challenges simulating the AMOC in climate models *Phil. Trans. R. Soc. A* **381** 20220187
- Ji D, Cui M, Chen Y and Dai Y 2025 Permafrost response in northern high-latitude regions to 1.5 C warming and overshoot scenarios achieved via solar radiation modification *J. Geophys. Res. Atmos.* **130** e2024JD041772
- Jiménez-Muñoz J C, Mattar C, Barichivich J, Santamaria-Artigas A, Takahashi K, Malhi Y, Sobrino J A and Schrier G V D 2016 Record-breaking warming and extreme drought in the Amazon rainforest during the course of El Niño 2015–2016 *Sci. Rep.* **6** 33130
- Jones C D et al 2019 The zero emissions commitment model intercomparison project (ZECMIP) contribution to C4MIP: quantifying committed climate changes following zero carbon emissions *Geosci. Model. Dev.* **12** 4375–85
- Jones C D and Friedlingstein P 2020 Quantifying process-level uncertainty contributions to TCRE and carbon budgets for meeting Paris Agreement climate targets *Environ. Res. Lett.* **15** 074019
- Kalmus P, Ekanayaka A, Kang E, Baird M and Gierach M 2022 Past the precipice? Projected coral habitability under global heating *Earth’s Future* **10** e2021EF002608
- Kaufhold C, Willeit M, Talento S, Ganopolski A and Rockström J 2025 Interplay between climate and carbon cycle feedbacks could substantially enhance future warming *Environ. Res. Lett.* **20** 044027
- Ke P et al 2024 Low latency carbon budget analysis reveals a large decline of the land carbon sink in 2023 *Natl Sci. Rev.* **11** nwae367
- Kikstra J S et al 2022 The IPCC sixth assessment report WGIII climate assessment of mitigation pathways: from emissions to global temperatures *Geosci. Model. Dev.* **15** 9075–109
- Kim I-W, Timmermann A, Kim J-E, Rodgers K B, Lee S-S, Lee H and Wieder W R 2024 Abrupt increase in Arctic-Subarctic wildfires caused by future permafrost thaw *Nat. Commun.* **15** 7868
- Kim S-K, Shin J, An S-I, Kim H-J, Im N, Xie S-P, Kug J-S and Yeh S-W 2022 Widespread irreversible changes in surface temperature and precipitation in response to CO₂ forcing *Nat. Clim. Change* **12** 834–40
- Klein S G, Roch C and Duarte C M 2024 Systematic review of the uncertainty of coral reef futures under climate change *Nat. Commun.* **15** 2224
- Klose A K, Coulon V, Pattyn F and Winkelmann R 2024a The long-term sea-level commitment from Antarctica *The Cryosphere* **18** 4463–92
- Klose A K, Donges J F, Feudel U and Winkelmann R 2024b Rate-induced tipping cascades arising from interactions between the Greenland ice sheet and the Atlantic meridional overturning circulation *Earth Syst. Dyn.* **15** 635–52

- Klose A K, Wunderling N, Winkelmann R and Donges J F 2021 What do we mean, 'tipping cascade'? *Environ. Res. Lett.* **16** 125011
- Knight J and Condron A 2025 Freshwater from Antarctica mitigates the risk of an AMOC slowdown (available at: www.researchsquare.com/article/rs-5369100/v1)
- Koven C D, Sanderson B M and Swann A L 2023 Much of zero emissions commitment occurs before reaching net zero emissions *Environ. Res. Lett.* **18** 014017
- Laybourn L, Abrams J F, Benton D, Brown K, Evans J, Elliott J, Swingedouw D, Lenton T M and Dyke J G 2024 The security blind spot: cascading climate impacts and tipping points threaten national security, IPPR (available at: www.ippr.org/articles/security-blind-spot)
- Lenton T M *et al* 2024 Remotely sensing potential climate change tipping points across scales *Nat. Commun.* **15** 343
- Lenton T M, Held H, Kriegler E, Hall J W, Lucht W, Rahmstorf S and Schellnhuber H J 2008 Tipping elements in the Earth's climate system *Proc. Natl Acad. Sci.* **105** 1786–93
- Lenton T M, Mckay D I A, Loriani S, Abrams J F, Lade S J, Donges J F and Rockström J 2023 The global tipping points report 2023 (University of Exeter)
- Lenton T M, Milkoreit M, Willcock S, Abrams J F, Mc Kay D L A, Buxton J E and Wieners C 2025 Global tipping points report 2025
- Levermann A and Born A 2007 Bistability of the Atlantic subpolar gyre in a coarse-resolution climate model *Geophys. Res. Lett.* **34** L24605
- Lewis S L, Brando P M, Phillips O L, Van Der Heijden G M F and Nepstad D 2011 The 2010 Amazon drought *Science* **331** 554
- Liu W, Xie S-P, Liu Z and Zhu J 2017 Overlooked possibility of a collapsed Atlantic Meridional Overturning Circulation in warming climate *Sci. Adv.* **3** e1601666
- Lohmann J, Castellana D, Ditlevsen P D and Dijkstra H A 2021 Abrupt climate change as rate-dependent cascading tipping point *Earth Syst. Dyn.* **12** 819–35
- Lohmann J, Dijkstra H A, Jochum M, Lucarini V and Ditlevsen P D 2024a Multistability and intermediate tipping of the Atlantic Ocean circulation *Sci. Adv.* **10** eadi4253
- Lohmann J and Ditlevsen P D 2021 Risk of tipping the overturning circulation due to increasing rates of ice melt *PNAS* **118** e2017989118
- Lohmann J, Wuyts B, Ditlevsen P D and Ashwin P 2024b On the predictability of possible storylines for forced complex systems *J. Phys. Complex* **5** 035015
- Lovejoy T E and Nobre C 2018 Amazon tipping point *Sci. Adv.* **4** eaat2340
- Lovejoy T E and Nobre C 2019 Amazon tipping point: last chance for action *Sci. Adv.* **5** eaba2949
- Lux-Gottschalk K and Ritchie P D L 2025 Uncertainty quantification for overshoots of tipping thresholds *Earth Syst. Dyn.* **16** 1153–68
- MacDougall A H *et al* 2020 Is there warming in the pipeline? A multi-model analysis of the Zero Emissions Commitment from CO₂ *Biogeosciences* **17** 2987–3016
- Malhi Y, Aragão L E O C, Galbraith D, Huntingford C, Fisher R, Zelazowski P, Sitch S, McSweeney C and Meir P 2009 Exploring the likelihood and mechanism of a climate-change-induced dieback of the Amazon rainforest *Proc. Natl Acad. Sci.* **106** 20610–5
- Marengo J A, Cunha A P, Espinoza J-C, Fu R, Schöngart J, Jimenez J C, Costa M C, Ribeiro J M, Wongchuig S and Zhao S 2024 The drought of Amazonia in 2023–2024 *Am. J. Clim. Change* **13** 567–97
- Mathison C *et al* 2025 A rapid-application emissions-to-impacts tool for scenario assessment: probabilistic regional impacts from model patterns and emissions (PRIME) *Geosci. Model. Dev.* **18** 1785–808
- Matthews H D, Gillett N P, Stott P A and Zickfeld K 2009 The proportionality of global warming to cumulative carbon emissions *Nature* **459** 829–32
- McClanahan T R *et al* 2024 Diversification of refugia types needed to secure the future of coral reefs subject to climate change *Conserv. Biol.* **38** e14108
- McKenna C M, Maycock A C, Forster P M, Smith C J and Tokarska K B 2021 Stringent mitigation substantially reduces risk of unprecedented near-term warming rates *Nat. Clim. Change* **11** 126–31
- Meehl G A, Senior C A, Eyring V, Flato G, Lamarque J-F, Stouffer R J, Taylor K E and Schlund M 2020 Context for interpreting equilibrium climate sensitivity and transient climate response from the CMIP6 Earth system models *Sci. Adv.* **6** eaba1981
- Meinshausen M *et al* 2024 A perspective on the next generation of Earth system model scenarios: towards representative emission pathways (REPs) *Geosci. Model. Dev.* **17** 4533–59
- Melnikova I, Boucher O, Cadule P, Ciais P, Gasser T, Quilcaille Y, Shiogama H, Tachiiri K, Yokohata T and Tanaka K 2021 Carbon cycle response to temperature overshoot beyond 2 C: an analysis of CMIP6 models *Earth's Future* **9** e2020EF001967
- Melnikova I, Hajima T, Shiogama H, Hayashi M, Ito A, Nishina K, Tachiiri K and Yokohata T 2025 Amazon dieback beyond the 21st century under high-emission scenarios by Earth System models *Commun. Earth Environ.* **6** 670
- Mengel M and Levermann A 2014 Ice plug prevents irreversible discharge from East Antarctica *Nat. Clim. Change* **4** 451–5
- Mengel M, Nauels A, Rogelj J and Schlessner C-F 2018 Committed sea-level rise under the Paris Agreement and the legacy of delayed mitigation action *Nat. Commun.* **9** 601
- Möller T, Högner A E, Schlessner C-F, Bien S, Kitzmann N H, Lamboll R D, Rogelj J, Donges J F, Rockström J and Wunderling N 2024 Achieving net zero greenhouse gas emissions critical to limit climate tipping risks *Nat. Commun.* **15** 6192
- Muñiz-Castillo A I, Rivera-Sosa A, Chollett I, Eakin C M, Andrade-Gómez L, McField M and Arias-González J E 2019 Three decades of heat stress exposure in Caribbean coral reefs: a new regional delineation to enhance conservation *Sci. Rep.* **9** 11013
- Myhre G, Hodnebrog Ø, Loeb N and Forster P M 2025 Observed trend in Earth energy imbalance may provide a constraint for low climate sensitivity models *Science* **388** 1210–3
- Nicholls Z R J *et al* 2020 Reduced complexity model intercomparison project phase 1: introduction and evaluation of global-mean temperature response *Geosci. Model. Dev.* **13** 5175–90
- Nijse F J M M, Cox P M and Williamson M S 2020 Emergent constraints on transient climate response (TCR) and equilibrium climate sensitivity (ECS) from historical warming in CMIP5 and CMIP6 models *Earth Syst. Dyn.* **11** 737–50
- Nitzbon J *et al* 2024 No respite from permafrost-thaw impacts in the absence of a global tipping point *Nat. Clim. Change* **14** 573–85
- Nobre C A, Sampaio G, Borma L S, Castilla-Rubio J C, Silva J S and Cardoso M 2016 Land-use and climate change risks in the Amazon and the need of a novel sustainable development paradigm *Proc. Natl Acad. Sci.* **113** 10759–68
- Noël B, Van Kampenhou L, Lenaerts J T M, Van de Berg W J and Van Den Broeke M R 2021 A 21st century warming threshold for sustained Greenland ice sheet mass loss *Geophys. Res. Lett.* **48** e2020GL090471
- O'Keeffe P E and Wicczorek S 2020 Tipping phenomena and points of no return in ecosystems: beyond classical bifurcations *SIAM J. Appl. Dyn. Syst.* **19** 2371–402
- Oh J-H, An S-I, Shin J and Kug J-S 2022 Centennial memory of the Arctic Ocean for future Arctic climate recovery in response to a carbon dioxide removal *Earth's Future* **10** e2022EF002804

- Palter J B, Frölicher T L, Paynter D and John J G 2018 Climate, ocean circulation, and sea level changes under stabilization and overshoot pathways to 1.5 K warming *Earth Syst. Dyn.* **9** 817–28
- Park S-W, Mun J-H, Lee H, Steinert N J, An S-I, Shin J and Kug J-S 2025 Continued permafrost ecosystem carbon loss under net-zero and negative emissions *Sci. Adv.* **11** eadn8819
- Parry I M, Ritchie P D L and Cox P M 2022 Evidence of localised Amazon rainforest dieback in CMIP6 models *Earth Syst. Dyn.* **13** 1667–75
- Pattyn F *et al* 2018 The Greenland and Antarctic ice sheets under 1.5 C global warming *Nat. Clim. Change* **8** 1053–61
- Pearce-Kelly P *et al* 2025 Considerations for determining warm-water coral reef tipping points *Earth Syst. Dyn.* **16** 275–92
- Pereira L M *et al* 2024 Equity and justice should underpin the discourse on tipping points *Earth Syst. Dyn.* **15** 341–66
- Petrini M, Scherrenberg M D W, Muntjewerf L, Vizcaino M, Sellevold R, Leguy G R, Lipscomb H and Goelzer H 2025 A topographically controlled tipping point for complete Greenland ice sheet melt *Cryosphere* **19** 63–81
- Pfleiderer P, Schleussner C F and Sillmann J 2024 Limited reversal of regional climate signals in overshoot scenarios *Environ. Res.* **3** 015005
- Pöppelmeier F and Stocker T F 2025 Mutual stabilization of AMOC and GRIIS due to different transient response to warming *Environ. Res. Lett.* **20** 094015
- Portmann V, Chavent M and Swingedouw D 2025 ClimLoco1.0: Climate variable confidence interval of multivariate linear observational constraint *EGUsphere* **18** 1–32
- Rao M P *et al* 2023 Approaching a thermal tipping point in the Eurasian boreal forest at its southern margin *Commun. Earth Environ.* **4** 247
- Reese R *et al* 2023 The stability of present-day Antarctic grounding lines—Part 2: onset of irreversible retreat of Amundsen Sea glaciers under current climate on centennial timescales cannot be excluded *The Cryosphere* **17** 3761–83
- Reimer J D *et al* 2024 The fourth global coral bleaching event: where do we go from here? *Coral. Reefs* **43** 1121–5
- Reisinger A *et al* 2025 Overshoot: a conceptual review of exceeding and returning to global warming of 1.5° C *Annu. Rev. Environ. Resour.* **50** 185–217
- Reisinger A and Geden O 2023 Temporary overshoot: origins, prospects, and a long path ahead *One Earth* **6** 1631–7
- Ridley J, Gregory J M, Huybrechts P and Lowe J 2010 Thresholds for irreversible decline of the Greenland ice sheet *Clim. Dyn.* **35** 1049–57
- Ripple W J *et al* 2023a The 2023 state of the climate report: entering uncharted territory *BioScience* **73** 841–50
- Ripple W J, Wolf C, Lenton T M, Gregg J W, Natali S M, Duffy P B, Rockström H J and Schellnhuber H J 2023b Many risky feedback loops amplify the need for climate action *One Earth* **6** 86–91
- Ritchie P D L, Alkhayouon H, Cox P M and Wiczorek S 2023 Rate-induced tipping in natural and human systems *Earth Syst. Dyn.* **14** 669–83
- Ritchie P D L, Clarke J J, Cox P M and Huntingford C 2021 Overshooting tipping point thresholds in a changing climate *Nature* **592** 517–23
- Ritchie P D L, Bastiaansen R, von der Heydt A S and Ashwin P 2025a The role of coupling and timescales for interacting tipping elements (arXiv:2509.03996v1)
- Ritchie P D, Huntingford C and Cox P M 2025b ESD Ideas. Climate tipping is not instantaneous—the duration of an overshoot matters *Egusphere* (<https://doi.org/10.5194/esd-16-1523-2025>)
- Ritchie P 2025 Python scripts to generate figures for paper titled ‘The implications of overshooting 1.5C on Earth system tipping elements - a review’ *GitHub* (available at: https://github.com/PaulRitchie89/Overshoots_review)
- Ritchie P, Karabacak Ö and Sieber J 2019 Inverse-square law between time and amplitude for crossing tipping thresholds *Proc. R. Soc. A* **475** 20180504
- Roberts M J *et al* 2025 High-resolution model intercomparison project phase 2 (HighResMIP2) towards CMIP7 *Geosci. Model. Dev.* **18** 1307–32
- Robinson A, Calov R and Ganopolski A 2012 Multistability and critical thresholds of the Greenland ice sheet *Nat. Clim. Change* **2** 429–32
- Rockström J, Gupta J, Qin D, Lade S J, Abrams J F, Andersen L S and Zhang X 2023 Safe and just Earth system boundaries *Nature* **619** 102–11
- Rockström J, Warszawski L and Schuster A 2024 A stable planet is a prerequisite for development *Horizons* **27** 36–49 (available at: www.jstor.org/stable/48790036)
- Romanou A, Rind D, Jonas J, Miller R, Kelley M, Russell G, Orbe C, Nazarenko L, Lato R and Schmidt G A 2023 Stochastic bifurcation of the North Atlantic circulation under a midrange future climate scenario with the NASA-GISS ModelE *J. Clim.* **36** 6141–61
- Rosser J P, Winkelmann R and Wunderling N 2024 Polar ice sheets are decisive contributors to uncertainty in climate tipping projections *Commun. Earth Environ.* **5** 702
- Rotbarth R, Van Nes E H, Scheffer M, Jepsen J U, Vindstad O P L, Xu C and Holmgren M 2023 Northern expansion is not compensating for southern declines in North American boreal forests *Nat. Commun.* **14** 3373
- Rounce D R *et al* 2023 Global glacier change in the 21st century: every increase in temperature matters *Science* **379** 78–83
- Rückamp M, Falk U, Frieler K, Lange S and Humbert A 2018 The effect of overshooting 1.5° C global warming on the mass loss of the Greenland ice sheet *Earth Syst. Dyn.* **9** 1169–89
- Sanderson B M *et al* 2025 flat10MIP: an emissions-driven experiment to diagnose the climate response to positive, zero, and negative CO₂ emissions *Geosci. Model Dev.* **18** 5699–724
- Sandstad M, Steinert N J, Baur S and Sanderson B M 2025 METEORv1.0.1: a novel framework for emulating multi-timescale regional climate responses *Geosci. Model Dev.* **18** 8269–312
- Saye L, Bettis O, Bedenham G, Hampshire O, Lenton T M and Abrams J F 2025 Planetary solvency—finding our balance with nature (available at: <https://actuaries.org.uk/news-and-media-releases/news-articles/2025/jan/16-jan-25-planetary-solvency-finding-our-balance-with-nature/>)
- Scheffer M, Hirota M, Holmgren M, Van Nes E H and Chapin F S III 2012 Thresholds for boreal biome transitions *Proc. Natl Acad. Sci.* **109** 21384–9
- Schleussner C-F *et al* 2016 Differential climate impacts for policy-relevant limits to global warming: the case of 1.5 C and 2 C *Earth Syst. Dyn.* **7** 327–51
- Schleussner C-F *et al* 2024 Overconfidence in climate overshoot *Nature* **634** 366–73
- Schleussner C-F, Ganti G, Rogelj J and Gidden M J 2022 An emission pathway classification reflecting the Paris Agreement climate objectives *Commun. Earth Environ.* **3** 135
- Schuster L, Maussion F, Rounce D R, Ulltee L, Schmitt P, Lacroix F and Schleussner C F 2025 Irreversible glacier change and trough water for centuries after overshooting 1.5° C *Nat. Clim. Change* **15** 634–41
- Schuur E A G *et al* 2022 Permafrost and climate change: carbon cycle feedbacks from the warming Arctic *Annu. Rev. Environ. Resour.* **47** 343–71
- Schwinger J, Asaadi A, Goris N and Lee H 2022a Possibility for strong northern hemisphere high-latitude cooling under negative emissions *Nat. Commun.* **13** 1095
- Schwinger J, Asaadi A, Steinert N J and Lee H 2022b Emit now, mitigate later? Earth system reversibility under overshoots of different magnitudes and durations *Earth Syst. Dyn.* **13** 1641–65
- Setter R O, Franklin E C, Mora C and Mumby P J 2022 Co-occurring anthropogenic stressors reduce the timeframe of environmental viability for the world’s coral reefs *PLoS Biol.* **20** e3001821

- Sgubin G, Swingedouw D, Drijfhout S, Mary Y and Bennabi A 2017 Abrupt cooling over the North Atlantic in modern climate models *Nat. Commun.* **8** 14375
- Sgubin G, Swingedouw D, García de Cortázar-atauri I, Ollat N and Van Leeuwen C 2019 The impact of possible decadal-scale cold waves on viticulture over Europe in a context of global warming *Agronomy* **9** 397
- Sinet S, Ashwin P, von der Heydt A S and Dijkstra H A 2024 AMOC stability amid tipping ice sheets: the crucial role of rate and noise *Earth Syst. Dyn.* **15** 859–73
- Sinet S, Von Der Heydt A S and Dijkstra H A 2023 AMOC stabilization under the interaction with tipping polar ice sheets *Geophys. Res. Lett.* **50** e2022GL100305
- Sinet S, von der Heydt A S and Dijkstra H A 2025 West Antarctic meltwater can prevent an AMOC Collapse (arXiv:2502.17104)
- Smith S, Geden O, Gidden M, Lamb W, Nemet G, Minx J and Vaughan N 2024 *The State of Carbon Dioxide Removal* (available at: <https://doi.org/10.17605/OSF.IO/F85QJ>)
- Solgaard A M and Langen P L 2012 Multistability of the Greenland ice sheet and the effects of an adaptive mass balance formulation *Clim. Dyn.* **39** 1599–612
- Souter D, Planes S, Wicquart J, Logan M, Obura D and Staub F 2021 Status of coral reefs of the world: 2020: executive summary (available at: <https://bvearmb.do/handle/123456789/3190>)
- Steffen W et al 2018 Trajectories of the earth system in the Anthropocene *Proc. Natl Acad. Sci.* **115** 8252–9
- Steinert N J and Sanderson B M 2025 Normalizing the permafrost carbon feedback contribution to TCRE and ZEC *EGU sphere*, 2025 in press pp 1–14
- Steinert N J, Schwinger J, Chadwick R, Kug J S and Lee H 2025 Irreversible land water availability changes from a potential ITCZ shift during temperature overshoot *Earth's Future* **13** e2024EF005787
- Stokes C R, Bamber J L, Dutton A and DeConto R M 2025 Warming of + 1.5° C is too high for polar ice sheets *Commun. Earth Environ.* **6** 1–12
- Stommel H 1961 Thermohaline convection with two stable regimes of flow *Tellus* **13** 224–30
- Stuenzi S M, Kruse S, Boike J, Herzschuh U, Oehme A, Pstryakova L A, Westermann S and Langer M 2022 Thermohydrological impact of forest disturbances on ecosystem-protected permafrost *J. Geophys. Res.* **127** e2021JG006630
- Swingedouw D, Bily A, Esquerdo C, Borchert L F, Sgubin G, Mignot J and Menary M 2021 On the risk of abrupt changes in the North Atlantic subpolar gyre in CMIP6 models *Ann. New York Acad. Sci.* **1504** 187–201
- Swingedouw D, Houssais M N, Herbaut C, Blaizot A C, Devilliers M and Deshayes J 2022 AMOC recent and future trends: a crucial role for oceanic resolution and Greenland melting? *Front. Clim.* **4** 838310
- Swingedouw D, Ifejika Speranza C, Bartsch A, Durand G, Jamet C, Beaugrand G and Conversi A 2020 Early warning from space for a few key tipping points in physical, biological, and social-ecological systems *Surv. Geophys.* **41** 1237–84
- Tachiiri K, Hajima T and Kawamiya M 2019 Increase of the transient climate response to cumulative carbon emissions with decreasing CO₂ concentration scenarios *Environ. Res. Lett.* **14** 124067
- Thompson J M T, Stewart H B and Ueda Y 1994 Safe, explosive, and dangerous bifurcations in dissipative dynamical systems *Phys. Rev. E* **49** 1019
- Tokarska K B and Zickfeld K 2015 The effectiveness of net negative carbon dioxide emissions in reversing anthropogenic climate change *Environ. Res. Lett.* **10** 094013
- Tokarska K B, Zickfeld K and Rogelj J 2019 Path independence of carbon budgets when meeting a stringent global mean temperature target after an overshoot *Earth's Future* **7** 1283–95
- Trust S B, Saye O, Bedenham L, Lenton G, Abrams T M and Kemp J F 2024 Climate scorpion—the sting is in the tail: introducing planetary solvency
- Turetsky M R, Abbott B W, Jones M C, Anthony K W, Olefeldt D, Schuur E A and McGuire A D 2020 Carbon release through abrupt permafrost thaw *Nat. Geosci.* **13** 138–43
- UNFCCC (United Nations Framework Convention on Climate Change) 1992 'Articles 1–26' (available at: <http://unfccc.int/resource/docs/convkp/conveng.pdf>)
- Valdes P 2011 Built for stability *Nat. Geosci.* **4** 414–6
- van den Akker T, Lipscomb W H, Leguy G R, Bernales J, Berends C J, van de Berg W J and van de Wal R S 2025 Present-day mass loss rates are a precursor for West Antarctic Ice Sheet collapse *Cryosphere* **19** 283–301
- Van Westen R M, Kliphuis M and Dijkstra H A 2024 Physics-based early warning signal shows that AMOC is on tipping course *Sci. Adv.* **10** eadk1189
- Veron J E, Hoegh-Guldberg O, Lenton T M, Lough J M, Obura D O, Pearce-Kelly P A U L, Sheppard C R C, Spalding M, Stafford-Smith M G and Rogers A D 2009 The coral reef crisis: the critical importance of < 350 = “ppm = “ > *Mar. Poll. Bull.* **58** 1428–36
- Wilson D J, Bertram R A, Needham E F, van de Flierdt T, Welsh K J, McKay R M, Mazumder A, Riesselman C R, Jimenez-Espejo F J and Escutia C 2018 Ice loss from the East Antarctic Ice Sheet during late Pleistocene interglacials *Nature* **561** 383–6
- Winkelmann R, Dennis D, Donges J F, Loriani S, Klose A K, Abrams J F and Yang S 2025 The tipping points modelling intercomparison project (TIPMIP): assessing tipping point risks in the Earth system *Earth Syst. Dyn. Discuss.* (<https://doi.org/10.5194/egusphere-2025-1899>)
- Winkler A J, Myneni R, Reimers C, Reichstein M, Brovkin V and Xu Y 2024 Carbon system state determines warming potential of emissions *PLoS One* **19** e0306128
- Wood R A, Baker J A, Beaugrand G, Boutin J, Conversi A, Donner R V and Swingedouw D 2024 Opportunities for earth observation to inform risk management for ocean tipping points *Surv. Geophys.* **46** 443–502
- World Meteorological Organization 2025 State of the Global Climate 2024, WMO-No. 1368
- Wunderling N et al 2024 Climate tipping point interactions and cascades: a review *Earth Syst. Dyn.* **15** 41–74
- Wunderling N, Donges J F, Kurths J and Winkelmann R 2021a Interacting tipping elements increase risk of climate domino effects under global warming *Earth Syst. Dyn.* **12** 601–19
- Wunderling N, Krönke J, Wohlfarth V, Kohler J, Heitzig J, Staal A, Willner S, Winkelmann R and Donges J F 2021b Modelling nonlinear dynamics of interacting tipping elements on complex networks: the PyCascades package *Eur. Phys. J. Special Top.* **230** 3163–76
- Wunderling N, Staal A, Sakschewski B, Hirota M, Tuinenburg O A, Donges J F, Barbosa H M J and Winkelmann R 2022 Recurrent droughts increase risk of cascading tipping events by outpacing adaptive capacities in the Amazon rainforest *Proc. Natl Acad. Sci.* **119** e212077119
- Wunderling N, Winkelmann R, Rockström J, Loriani S, Armstrong McKay D I, Ritchie P D L, Sakschewski B and Donges J F 2023 Global warming overshoots increase risks of climate tipping cascades in a network model *Nat. Clim. Change* **13** 75–82

- Zekollari H *et al* 2025 Glacier preservation doubled by limiting warming to 1.5° C versus 2.7° C *Science* **388** 979–83
- Zekollari H, Huss M and Farinotti D 2020 On the imbalance and response time of glaciers in the European Alps *Geophys. Res. Lett.* **47** e2019GL085578
- Zhuge C, Li J and Chen W 2025 Deep learning for predicting the occurrence of tipping points *R. Soc. Open Sci.* **12** 242240
- Zickfeld K, Azevedo D, Mathesius S and Matthews H D 2021 Asymmetry in the climate–carbon cycle response to positive and negative CO₂ emissions *Nat. Clim. Change* **11** 613–7
- Zickfeld K, MacDougall A H and Matthews H D 2016 On the proportionality between global temperature change and cumulative CO₂ emissions during periods of net negative CO₂ emissions *Environ. Res. Lett.* **11** 055006