



International Institute for
Applied Systems Analysis
www.iiasa.ac.at

On the Local and Global Convergence of a Reduced Quasi- Newton Method

Gilbert, J.C.

IIASA Working Paper

WP-87-113

October 1987



Gilbert JC (1987). On the Local and Global Convergence of a Reduced Quasi-Newton Method. IIASA Working Paper. IIASA, Laxenburg, Austria: WP-87-113 Copyright © 1987 by the author(s). <http://pure.iiasa.ac.at/id/eprint/2939/>

Working Papers on work of the International Institute for Applied Systems Analysis receive only limited review. Views or opinions expressed herein do not necessarily represent those of the Institute, its National Member Organizations, or other organizations supporting the work. All rights reserved. Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage. All copies must bear this notice and the full citation on the first page. For other purposes, to republish, to post on servers or to redistribute to lists, permission must be sought by contacting repository@iiasa.ac.at

WORKING PAPER

ON THE LOCAL AND GLOBAL CONVERGENCE
OF A REDUCED QUASI-NEWTON METHOD

Jean Charles GILBERT

October 1987
WP-87-113

**ON THE LOCAL AND GLOBAL CONVERGENCE
OF A REDUCED QUASI-NEWTON METHOD**

Jean Charles GILBERT

October 1987
WP-87-113

Working Papers are interim reports on work of the International Institute for Applied Systems Analysis and have received only limited review. Views or opinions expressed herein do not necessarily represent those of the Institute or of its National Member Organizations.

**INTERNATIONAL INSTITUTE FOR APPLIED SYSTEMS ANALYSIS
A-2361 Laxenburg, Austria**

FOREWORD

This paper deals with the approximation of solutions of general equality constrained optimization problems and proposed an algorithm of the reduced type that locally may be an alternative to the Newton method. The algorithm has indeed the same rate of convergence as Newton method while only operators of reduced order are needed.

Those types of methods may be particularly useful when the number of constraints is large while the number of free parameters remains reasonable.

A technique to globalize the local method is also proposed and studied.

Alexander B. Kurzhanski
Chairman
System and Decision Sciences Program

Abstract: In optimization in \mathbb{R}^n with m nonlinear equality constraints, we study the local convergence of reduced quasi-Newton methods, in which the updated matrix is of order $n-m$. In particular, we give necessary and sufficient conditions for q-superlinear convergence (in one step). We introduce a device to globalize the local algorithm which consists in determining a step on an arc in order to decrease an exact penalty function. We give conditions so that asymptotically the step will be equal to one.

Abbreviated title: A reduced quasi-Newton method.

Key words: Constrained Optimization, Successive Quadratic Programming, Reduced Quasi-Newton Method, Superlinear Convergence, Exact Penalty Function, Arc Search, Step-size Selection Procedure, Global Convergence.

Subject classification AMS(MOS): 49D05, 49D30, 65K05.

1. Introduction

Let ω be an open convex set in \mathbb{R}^n , f be a real-valued function on ω and c map ω to \mathbb{R}^m , where $m < n$. We shall suppose that f and c are functions of class C_b^ν with $\nu \geq 3$, i.e. f and c are supposed to be three times continuously differentiable with bounded derivatives on ω . We shall endow \mathbb{R}^n with its canonical basis. We are interested in algorithms for solving the following minimization problem with equality constraints:

$$\min \{ f(x) : x \in \mathbb{R}^n, c(x) = 0 \}. \quad (1.1)$$

In addition to the smoothness of f and c , we shall assume that c is a *submersion* on ω , that is to say that the $m \times n$ Jacobian matrix

$$A_x := A(x) \quad (1.2)$$

of partial derivatives of c at x is supposed to be surjective for all x in ω . If ω is "large" this is a very strong hypothesis, but it is usual to suppose that the gradients of the constraints are linearly independent at a solution of (1.1) and, therefore, this hypothesis is satisfied in a neighborhood of a solution. Then, if x_* is a local minimizer for problem (1.1), there exists a unique Lagrange multiplier λ_* so that the first order optimality conditions are satisfied:

$$\begin{cases} c(x_*) = 0, \\ \nabla f(x_*) + A_*^T \lambda_* = 0, \end{cases} \quad (1.3)$$

where $\nabla f(x_*)$ is the vector of partial derivatives of f at x_* and $A_* := A(x_*)$. The quantity in the left hand side of the second equation is the first derivative with respect to x at (x_*, λ_*) of the Lagrangian $l(x, \lambda) := f(x) + (\lambda, c(x))$. The second order sufficient condition will also be assumed: the $n \times n$ Hessian matrix L_* of second derivatives with respect to x of l at (x_*, λ_*) is supposed to be positive definite in the null space $N(A_*)$ of A_* . For further references, we gather those hypotheses under the name of

assumption A:

- f, c are in $C_b^\nu(\omega)$ with $\nu \geq 3$,
- c is a submersion,
- (x_*, λ_*) satisfies (1.3),
- $h^T L_* h > 0$ for all h in \mathbb{R}^n with $h \neq 0$ and $A_* h = 0$.

Quasi-Newton methods, also called variable metric or secant methods, are methods for solving a system of nonlinear equations on \mathbb{R}^N , say $F(x_*) = 0$, that generate a sequence of points (x_k) and a sequence of nonsingular matrices (J_k) of order N

from the data of a point x_1 , and a nonsingular matrix J_1 by the formula:

$$x_{k+1} = x_k - J_k^{-1} F(x_k),$$

where J_k is updated at each iteration according to the following scheme:

$$J_{k+1} = U(J_k, \gamma_k, \sigma_k),$$

$$\gamma_k := F(x_{k+1}) - F(x_k),$$

$$\sigma_k := x_{k+1} - x_k.$$

The rule U is designed in order that J_{k+1} will satisfy the secant equation $J_{k+1}\sigma_k = \gamma_k$ and then will improve the approximation by J_k of the Jacobian matrix $\nabla F(x_*)$ at the solution x_* . These methods are particularly attractive because second order derivatives need not be calculated and because a q-superlinear rate of convergence for (x_k) can be obtained (see the review paper by Dennis and Moré (1977)), i.e.

$$\frac{\|x_{k+1} - x_*\|}{\|x_k - x_*\|} \rightarrow 0 \text{ as } k \rightarrow \infty. \quad (1.4)$$

Such a method could be used for solving system (1.3), but matrices of order $n+m$ should be updated.

The aim of this paper is to introduce and study such quasi-Newton methods but with updated matrices of order $n-m$.

The *Successive Quadratic Programming* (SQP) method proposed by Wilson (1963) and Han (1976) improves the method described above with regard to the order of the updated matrices. In this method x_{k+1} is obtained from x_k by solving a quadratic program with linear constraints:

$$\begin{cases} \min [f'(x_k) \cdot (x - x_k) + \frac{1}{2} (x - x_k)^T L_k (x - x_k)], \\ x \in \mathbb{R}^n \text{ and } c(x_k) + c'(x_k) \cdot (x - x_k) = 0, \end{cases} \quad (1.5)$$

where we have used a dot to separate the linear operators $f'(x_k)$ and $c'(x_k)$ from their argument $(x - x_k)$. The matrix L_k of order n is updated in order to approximate L_* , the Hessian of the Lagrangian. Under assumption A, the solution of (1.5) can be written in the form (see Gabay (1982,b)):

$$x_{k+1} = x_k - A_k^- c(x_k) - Z_k^- (Z_k^{-T} L_k Z_k^-)^{-1} [g(x_k) - Z_k^{-T} L_k A_k^- c(x_k)], \quad (1.6)$$

where A_k^- is any right inverse of $A_k := A(x_k)$, Z_k^- is any $n \times (n-m)$ matrix whose columns form a basis of the tangent space $N(A_k)$ to the manifold $M_k := c^{-1}(c(x_k))$ at x_k and $g(x_k)$ is the *reduced gradient* defined by

$$g(x_k) := Z_k^{-T} \nabla f(x_k) \in \mathbb{R}^{n-m}. \quad (1.7)$$

The first part $(-A_k^- c(x_k))$ of the displacement in (1.6) is a *restoration step*, i.e. a Newton-like step for solving $c(x_*) = 0$. It belongs to $R(A_k^-)$ which is a complementary space to $N(A_k)$ in \mathbb{R}^n . The second part of the displacement in (1.6) is a *minimization step* belonging to the tangent space $R(Z_k^-) = N(A_k)$ to M_k at x_k .

Let us consider for a while the case where the constraints are linear:

$$c(x) := Ax + b = 0, \quad (1.8)$$

where A is an $m \times n$ surjective matrix and b is a vector in \mathbb{R}^m . As in the nonlinear case, let us introduce Z^- , an $n \times (n-m)$ matrix whose columns generate $N(A) : AZ^- = 0$. Suppose that the first iterate belongs to the plane of the constraints: $c(x_1) = 0$. Then, any point x satisfying the constraints (1.8) can be expressed by using a reduced variable u in \mathbb{R}^{n-m} ($x = x_1 + Z^-u$) and the problem of minimizing f subject to the constraints (1.8) is equivalent to the one of minimizing $\varphi(u) := f(x_1 + Z^-u)$ on $\{u \in \mathbb{R}^{n-m} : x_1 + Z^-u \in \omega\}$:

$$\min \{\varphi(u) : u \in \mathbb{R}^{n-m}, x_1 + Z^-u \in \omega\}. \quad (1.9)$$

By considering the optimality equation $\nabla\varphi(u_*) = Z^{-T}\nabla f(x_1 + Z^-u_*) = 0$, a quasi-Newton method for solving problem (1.9) generates a sequence (u_k) in \mathbb{R}^{n-m} and a sequence (G_k) of nonsingular matrices of order $n-m$ such that

$$u_{k+1} = u_k - G_k^{-1} \nabla\varphi(u_k).$$

By setting $x_k := x_1 + Z^-u_k$, we obtain

$$x_{k+1} = x_k - Z^- G_k^{-1} g(x_k), \quad (1.10)$$

where $g(x_k) := Z^{-T}\nabla f(x_k)$ is the reduced gradient of f at x_k and G_k is updated as follows:

$$\begin{aligned} G_{k+1} &= U(G_k, \gamma_k, \sigma_k), \\ \gamma_k &:= g(x_{k+1}) - g(x_k), \\ Z^- \sigma_k &:= x_{k+1} - x_k, \end{aligned} \quad (1.11)$$

in order to approximate $\nabla^2\varphi(u_*)$, the Hessian of φ at u_* , which is also the *reduced Hessian of f at x_** :

$$Z^{-T} \nabla^2 f(x_*) Z^-.$$

The conditions so that the rate of convergence of the sequence (u_k) will be q -superlinear can be satisfied and because of the injectivity of Z^- , the same conditions will assure the q -superlinear convergence of the sequence (x_k) .

The algorithm (1.10)-(1.11) is a *reduced quasi-Newton method* because the order of the updated matrices is $n-m$ rather than n in the SQP. Our aim is to study such methods in the case of nonlinear constraints. They are particularly well adapted to problems with n large and $n-m$ small. That situation appears for example in the parametric identification of nonlinear sources in elliptic partial differential equations. If a finite element method is used to discretize the state equations (the constraints), m is large, say of the order of 1000, whereas the number $n-m$ of identifiable parameters is usually small: 2 or 3 in the example of Blum, Gilbert and Thooris (1985). In that case, a reduced quasi-Newton method is usable while the SQP method is not because of the order of the matrices that should be updated. Another advantage in developing reduced quasi-Newton methods comes from the fact that, under assumption A, the projected Hessian of the Lagrangian is positive definite at the optimum. Therefore, positive definite quasi-Newton approximations of that operator can be generated, in particular by the BFGS formula which is at present widely believed to be the best update formula. We see that reduced quasi-Newton methods appears rather natural. So, it is important to generalize the algorithm (1.10)-(1.11) in case the constraints are nonlinear.

This can be done by using the implicit function theorem in order to obtain a reduced objective function:

$$\varphi(u) = f(\xi(u)),$$

where $\xi : u \in V \subset \mathbb{R}^{n-m} \rightarrow \xi(u) \in \omega \subset \mathbb{R}^n$ is a parametric representation of the regular manifold $c^{-1}(0)$ around $x_* := \xi(u_*)$. We have $c(\xi(u)) = 0$ for all u in the neighborhood V of u_* . This is the basic idea of methods like the Generalized Reduced Gradient (GRG) method (Abadie and Carpentier (1969)). In fact, the parametric representation $\xi(u)$ is usually not known and this leads to several difficulties. Because the method asks the generated sequence (x_k) to be feasible ($c(x_k) = 0$ for all k), and because this cannot be achieved exactly in practice, some criterion has to be introduced to decide when to stop the restoration steps, i.e. how well the equality $x_{k+1} = \xi(u_{k+1})$ has to be realized (Mukai and Polak (1978)). Another difficulty appears when x_k is far from x_* and a step-size has to be introduced in the u -space in order to globalize the method. Indeed, every time a step-size is tried, an infinite number of restoration steps have to be done: see Gabay (1975), Gabay and Luenberger (1976), Mukai and Polak (1978) and Gabay (1982,a).

On the other hand, some non-feasible reduced quasi-Newton methods have been developed recently. Gabay (1982,b) has studied the following algorithm:

$$x_{k+1} = x_k + r_k^1 + t_k^1, \quad (1.12)$$

$$r_k^1 := -A_k^- c(x_k), \quad (1.13)$$

$$t_k^1 := -Z_k^- G_k^{-1} g(x_k), \quad (1.14)$$

where A_k^- is any right inverse of A_k , Z_k^- is any $n \times (n-m)$ matrix whose columns form a basis of the tangent space $N(A_k)$, G_k is a nonsingular matrix of order $(n-m)$ and $g(x_k)$ is the reduced gradient of f at x_k . The *tangent step* t_k^1 in (1.12), tangent to the manifold M_k , has the same structure as the displacement in (1.10) except for the basis Z_k^- , which changes here at each iteration. The *restoration step* r_k^1 in (1.12) is introduced to improve the feasibility of the sequence. The displacement in (1.12) can also be deduced from the displacement (1.6) of the SQP method by dropping the last part of the minimization step and by considering G_k as an approximation of the projected Hessian $Z_k^{-T} L(x_k, \lambda_k) Z_k^-$.

For their part, Coleman and Conn (1982,a) have studied the following algorithm:

$$x_{k+1} = x_k + r_k^2 + t_k^2, \quad (1.15)$$

$$r_k^2 := - A_k^- c(x_k + t_k^2), \quad (1.16)$$

$$t_k^2 := - Z_k^- G_k^{-1} g(x_k), \quad (1.17)$$

where Z_k^- is an $n \times (n-m)$ matrix whose columns form an orthogonal basis of $N(A_k)$ and A_k^- is the Penrose pseudo-inverse of A_k : $A_k^- := A_k^T (A_k A_k^T)^{-1}$. A relevant difference with the algorithm of Gabay lies in the restoration step in which the constraints are evaluated at $x_k + t_k^2$, after the tangent step, rather than at x_k in the algorithm (1.12)–(1.14).

The study of both algorithms (1.12)–(1.14) and (1.15)–(1.17) shows that when the matrices G_k are suitably chosen and the initial point x_1 is close to x_* , the sequence (x_k) generated by any of those algorithms converges to x_* q-superlinearly in two steps, that is to say:

$$\frac{\|x_{k+1} - x_*\|}{\|x_{k-1} - x_*\|} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

This rate of convergence is not so good as the rate (1.4) obtained with reduced quasi-Newton methods when the constraints are linear or with the SQP method. On the other hand, counter-examples have been given by Byrd (1985) and Yuan (1985) for which both methods of Gabay and Coleman and Conn do not converge better than with a two step q-superlinear rate of convergence. Therefore, the question of the rate of convergence of reduced quasi-Newton methods seemed closed. However, Byrd (1984) and Gilbert (1986,a) have shown (independently) that the sequence $(x_k + t_k^2)$ of Coleman and Conn's algorithm converges q-superlinearly in one step. This fact makes this method competitive. Indeed, the quasi-Newton version of Coleman and Conn's algorithm (see Coleman and Conn (1984)) needs two linearizations of the constraints per iteration. Consequently, as mentioned by Byrd (1984), it was thought that a super-linear step was made in this algorithm for each four constraint linearizations. The result of convergence of Byrd (1984) and Gilbert (1986,a) shows however that one

superlinear step is made for each two constraint linearizations. Furthermore, it can be shown (see Gilbert (1986,a)) that the use of an update criterion allows linearization of the constraints only once per iteration. Therefore, the method makes a superlinear step for each constraint linearization.

In this paper, we first show how the algorithm (1.15)–(1.17) can be obtained from a very general principle. If unconstrained optimization problems are related to the solution of nonlinear equations (via the optimality condition $f'(x_*) = 0$), we see from (1.3) that constrained optimization problems are closely related to the solution of two coupled nonlinear equations:

$$c(x_*) = 0 \text{ in } \mathbb{R}^m, \quad (1.18)$$

$$g(x_*) = 0 \text{ in } \mathbb{R}^{n-m}. \quad (1.19)$$

The equation (1.19) expresses the vanishment of the reduced gradient defined in (1.7) and is obtained by projecting the second equation of (1.3) on the tangent space $N(A_*)$. A "decoupling" method for solving (1.18)–(1.19) is introduced in *Section 2*. At x_k , the first part of the step of the method consists in doing a Newton-like displacement for solving (1.18). This leads to a point y_k . Then, x_{k+1} is obtained by doing a Newton-like displacement for solving (1.19) from the point y_k :

$$y_k = x_k - A_k^- c(x_k), \quad (1.20)$$

$$x_{k+1} = y_k - B_k^- g(y_k). \quad (1.21)$$

A_k^- is a right inverse of $\nabla c(x_*)$ and B_k^- is a right inverse of $\nabla g(x_*)$. We shall show that only conditions on B_k^- have to be imposed in order to ensure the local q-quadratic convergence (in one step) of the process. In *Section 4*, we apply this algorithm to constrained optimization, when g in (1.19) has the special structure (1.7), and it takes the form of a reduced method. Its extension to reduced quasi-Newton method is then easily done. This presentation gives, in our opinion, some insight into the method (1.15)–(1.17) and generalizes in some way the choice of the operators A_k^- in (1.16) and Z_k^- in (1.17) made by Coleman, Conn and Byrd. On the one hand, A_k^- may be any right inverse of A_k (with a smoothness hypothesis of $A_k^- = A(x_k)^-$ according to x_k) and not necessary the Penrose pseudo-inverse and on the other hand, the columns of Z_k^- may form any basis of $N(A_k)$ and not necessarily an orthogonal basis. This remark may be crucial in some applications like the one mentioned above where the "partitioned framework" (see *Section 9*) occurs naturally. In *Section 5*, we give a necessary and sufficient condition of q-superlinear convergence of the method that is weaker than the sufficient condition given by Byrd (1984).

The globalization of the local method could then be done as in the paper of Coleman and Conn (1982,b). In *Section 6*, however, we examine another globalizing technique essentially based on the ideas of Han (1977) for the SQP method (see also Danilin

and Pschenichny (1965)). We introduce the following exact penalty function:

$$\Theta_p(x) = f(x) + p \|c(x)\|_1, \quad (1.22)$$

where p is a large enough penalty parameter and $\|\bullet\|_1$ is the l_1 norm on \mathbb{R}^m . We look for x_* by minimizing Θ_p on ω . The idea is then to obtain a descent direction for Θ_p at the current iterate from the displacement calculated by the local algorithm (1.20)–(1.21). Contrary to what happens with the SQP method, our total displacement is not a descent direction for Θ_p any more. So, we shall introduce a descent arc, being inspired in that way by the work of Gabay (1982,b) for his algorithm and Mayne and Polak (1982) for the SQP method, although in those algorithms, the arc was introduced for other reasons. A search on the arc is done in order to decrease the penalty function Θ_p with the help of an Armijo-like criterion. This gives a theorem assuring the global convergence of the method. Furthermore, under natural conditions, the "Maratos effect" is avoided: the step-size is equal to one after a finite number of iterations. Therefore there is a smooth transition from the global to the local method that does not prevent the q-superlinear convergence from occurring.

If (v_k) is a sequence in a normed space $(E, \|\bullet\|_E)$ and (α_k) is a sequence of positive numbers, we shall say that (v_k) is a big O of (α_k) (we shall note $v_k = O(\alpha_k)$) if the sequence $(\|v_k\|_E/\alpha_k)$ is bounded and we shall say that (v_k) is a small o of (α_k) (we shall note $v_k = o(\alpha_k)$) if the sequence $(\|v_k\|_E/\alpha_k)$ converges to zero. We shall say that two positive real sequences (α_k) and (β_k) are *equivalent* (we shall note $\alpha_k \sim \beta_k$) if $\alpha_k = O(\beta_k)$ and $\beta_k = O(\alpha_k)$. We shall note v^i , the i -th component of a vector v in E . If A is a linear operator from $(E, \|\bullet\|_E)$ to $(F, \|\bullet\|_F)$, we shall note $\|A\| = \sup \{\|Av\|_F : \|v\|_E \leq 1\}$. If A and B are two square matrices of the same order, we shall note $A \leq B$ when $(B-A)$ is positive semi definite.

This paper constitutes a revised version of a part of the report number RR-482 of INRIA in which some techniques for updating the reduced matrix have also been investigated (see also Gilbert (1987)). A variant of the method is given in Gilbert (1986,b).

2. A decoupling method for solving two nonlinear coupled equations

Let us consider the following coupled system of nonlinear equations:

$$\begin{cases} F(x) = 0 \\ G(x) = 0 \end{cases} \quad (2.1)$$

where F and G are supposed smooth and map \mathbb{R}^n to \mathbb{R}^m and \mathbb{R}^{n-m} ($m < n$) respectively. Let x_* be a solution of (2.1) and let us denote by A_* the $m \times n$ Jacobian matrix of F at x_* and B_* the $(n-m) \times n$ Jacobian matrix of G at x_* . We shall say that x_* is a *regular solution* of (2.1) if the Jacobian matrix of the system (2.1),

$$J_* := \begin{bmatrix} A_* \\ B_* \end{bmatrix}, \quad (2.2)$$

is nonsingular. This will be the case if and only if $N(A_*) \cap N(B_*) = \{0\}$. We would like to define a Newton-like method for solving (2.1) without having to inverse the Jacobian $J(x_k)$ of (2.1) at x_k . For that, let us suppose that x_* is a regular solution. Then A_* and B_* are surjective and we can introduce a right inverse A_*^- of A_* and a right inverse B_*^- of B_* :

$$A_* A_*^- = I_m, \quad B_* B_*^- = I_{n-m}. \quad (2.3)$$

Two algorithms using A_*^- and B_*^- can be considered and we introduce them with the help of the fixed point maps ξ_1 and ξ_2 . The first one is

$$x_{k+1} = \xi_1(x_k), \quad (2.4)$$

$$\xi_1(x) := x - A_*^- F(x) - B_*^- G(x) \quad (2.5)$$

and the second one is

$$x_{k+1} = \xi_2(x_k) := (\psi \circ \varphi)(x_k), \quad (2.6)$$

$$\varphi(x) := x - A_*^- F(x), \quad (2.7)$$

$$\psi(y) := y - B_*^- G(y). \quad (2.8)$$

These algorithms are somewhat "ideal". Indeed, the matrices A_* and B_* are not known and so neither are A_*^- and B_*^- . But they are simpler to study than their implementable versions that will be introduced later. The relations (2.3) do not determine the right inverses A_*^- and B_*^- completely. Therefore, we can try to choose them so that the sequences generated by the algorithms (2.4)–(2.5) and (2.6)–(2.8) will have a good local behavior. The next two propositions show that this is possible: we can get conditions on A_*^- and B_*^- in order to have $\xi_1'(x_*) = 0$ and $\xi_2'(x_*) = 0$, which will ensure the q-quadratic rate of convergence for both algorithms. We shall say that an n row matrix is a *basis* of a given subspace of \mathbb{R}^n if its columns form a basis of that subspace.

Proposition 2.1. Suppose that F and G are differentiable at x_* , a regular solution of (2.1). Then, the following statements are equivalent:

- (i) $\xi_1'(x_*) = 0$,
- (ii) $R(A_*^-) = N(B_*)$ and $R(B_*^-) = N(A_*)$,
- (iii) for any right inverse A_*^- of A_* and any basis Z_*^- of $N(A_*)$, we have

$$A_*^- = [I - Z_*^- (B_* Z_*^-)^{-1} B_*] A_*^- \quad (2.9)$$

$$B_*^- = Z_*^- (B_* Z_*^-)^{-1}. \quad (2.10)$$

Proof. First, we prove (i) \iff (ii). Statement (i) is equivalent to

$$I = A_*^- A_* + B_*^- B_* . \quad (2.11)$$

The right hand side of (2.11) is equal to $[A_*^- B_*^-]J_*$. Then, (2.11) means that $[A_*^- B_*^-]$ is the inverse of J_* and therefore is equivalent to $I = J_*[A_*^- B_*^-]$, i.e. $A_* B_*^- = 0$ and $B_* A_*^- = 0$, which is equivalent to statement (ii) because the matrices A_* , B_* , A_*^- and B_*^- have full rank. Next, we prove (ii) \implies (iii). Let Z_*^- be any basis of $N(A_*)$: $R(Z_*^-) = N(A_*)$. Because J_* is nonsingular, $B_* Z_*^-$ is nonsingular. Indeed, if u in \mathbb{R}^{n-m} satisfies $B_* Z_*^- u = 0$, we have $J_* Z_*^- u = 0$ (because $A_* Z_*^- = 0$), then $Z_*^- u = 0$ because J_* is nonsingular and $u = 0$ because Z_*^- is injective. Then, by multiplying (2.11) to the right by Z_*^- , we get $Z_*^- = B_*^- (B_* Z_*^-)$ and therefore (2.10). (2.9) is obtained by multiplying (2.11) to the right by any right inverse A_*^- of A_* and by using (2.10). It remains to prove (iii) \implies (ii). If we take $A_*^- = A_*^-$ in (2.9), we obtain $B_* A_*^- = 0$ and so $R(A_*^-) = N(B_*)$ because these spaces have the same dimension m . Because $A_* Z_*^- = 0$, (2.10) gives $A_* B_*^- = 0$ and so $R(Z_*^-) = N(A_*)$ because these spaces have the same dimension $n-m$. \square

Proposition 2.2. Suppose that F and G are differentiable at x_* , a regular solution of (2.1). Then, the following statements are equivalent:

- (i) $\xi_2'(x_*) = 0$,
- (ii) $R(B_*^-) = N(A_*)$,
- (iii) for any basis Z_*^- of $N(A_*)$, we have $B_*^- = Z_*^- (B_* Z_*^-)^{-1}$.

Proof. The equivalence (i) \iff (ii) comes from the fact that $\xi_2'(x_*) = (I - B_*^- B_*) (I - A_*^- A_*)$ and that the spaces $N(A_*) = R(I - A_*^- A_*)$ and $R(B_*^-) = N(I - B_*^- B_*)$ have the same dimension $n-m$. To prove that (i) implies (iii), let Z_*^- be any basis of $N(A_*)$. By multiplying to the right both sides of

$$(I - B_*^- B_*)(I - A_*^- A_*) = 0$$

by Z_*^- , we get $Z_*^- = B_*^- (B_* Z_*^-)$ and therefore (iii) because $B_* Z_*^-$ is nonsingular (see the proof of proposition 2.1). From (iii), we get $A_* B_*^- = 0$ by multiplying to the left $B_*^- = Z_*^- (B_* Z_*^-)^{-1}$ by A_* and $A_* B_*^- = 0$ is equivalent to (ii) because $R(B_*^-)$ and $N(A_*)$ have the same dimension $n-m$. \square

In the statement (iii) of proposition 2.1, we could equivalently have given to B_* the role of A_* . We also see that the right inverses A_*^- and B_*^- are completely determined by the condition (i) of proposition 2.1 and do not depend on the choice of A_*^- and Z_*^- in (iii). Similarly, the right inverse B_*^- is completely determined by condition (i) of proposition 2.2 and does not depend on the choice of Z_*^- in (iii).

From proposition 2.1, we see that $\xi'_1(x_*) = 0$ if and only if $[A_*^- \ B_*^-]$ is the inverse of J_* . This means that the algorithm (2.4)–(2.5) is in fact the “ideal” (with J_*^{-1} rather than $J(x_k)^{-1}$) *Newton method* for solving (2.1) (see the displacement in (2.5)), the method we wanted to avoid. On the other hand, proposition 2.2 shows that the algorithm (2.6)–(2.8) needs fewer conditions to have a good local behavior than the algorithm (2.4)–(2.5). The fact that no conditions are required on the right inverse A_*^- means that any solver of the first equation in (2.1) can be used in (2.7), independently of the second equation of (2.1), whereas this is not true for the solver B_*^- of the second equation of (2.1) that has to be adapted to the first equation.

The results of propositions 2.1 and 2.2 have a geometrical interpretation. In the ideal Newton method, (x_k) will converge q-quadratically if the displacements $(-A_*^- F(x_k))$ and $(-B_*^- G(x_k))$ belong to the tangent space at x_* to the manifolds defined respectively by the pre-image of 0 by F and G . In the method (2.6)–(2.8) only the second step $(-B_*^- G(y_k))$ has to belong to the tangent space $N(A_*)$; the first step is arbitrary (apart from the fact that A_*^- has to be a right inverse of A_*).

3. A change of coordinates

Before applying the results of the previous section to constrained optimization, let us give some examples of right inverses A_x^- of A_x and basis Z_x^- of $N(A_x)$. The formalism adopted here has been introduced by Gabay (1982,a).

Once the injective matrices A_x^- and Z_x^- have been chosen, the columns of $[A_x^- \ Z_x^-]$ form a new basis of \mathbb{R}^n . Indeed, $R(A_x)$ is a complementary space of $N(A_x) = R(Z_x^-)$. To make a change of coordinates in that new basis, it is convenient to introduce the additional $(n-m) \times n$ matrix Z_x given in the following proposition.

Proposition 3.1. Let A_x be an $m \times n$ ($m < n$) surjective matrix, A_x^- be any right inverse of A_x and Z_x^- be any basis of $N(A_x)$. Then, there exists a unique $(n-m) \times n$ matrix Z_x such that

$$Z_x A_x^- = 0, \quad (3.1)$$

$$Z_x Z_x^- = I_{n-m}. \quad (3.2)$$

Furthermore, we have

$$I_n = A_x^- A_x + Z_x^- Z_x. \quad (3.3)$$

Proof. Existence and unicity of the matrix Z_x come from the nonsingularity of $[A_x^- \ Z_x^-]$ and (3.3) comes from the fact that $[A_x^T \ Z_x^T]^T$ is the inverse of $[A_x^- \ Z_x^-]$. \square

The relation (3.1) shows that $N(Z_x) = R(A_x^-)$ (the matrices A_x^- and Z_x have full rank) and (3.2) shows that Z_x^- is a right inverse of Z_x . The equality (3.3) can be used to introduce a change of coordinates. Indeed, by applying it to a vector ξ of \mathbb{R}^n , we see that $A_x \xi$ are the coordinates of ξ in $R(A_x^-) = N(Z_x)$ and $Z_x \xi$ are the coordinates of ξ in $R(Z_x^-) = N(A_x)$.

A first choice of matrices A_x^- and Z_x^- , which is frequently made in constrained optimization, defines what could be called the *orthogonal framework*: A_x^- is the Moore-Penrose pseudo-inverse of A_x (see Ben, Israel and Greville (1974)) and Z_x^- is an orthogonal basis of $N(A_x)$ for the scalar product $(\xi, \eta) = \sum_{i=1}^n \xi^i \eta^i$. We have:

$$A_x^- = A_x^T (A_x A_x^T)^{-1}, \quad (3.4)$$

$$Z_x^{-T} Z_x^- = I_{n-m}.$$

Then $Z_x = Z_x^{-T}$ is the unique matrix satisfying (3.1) and (3.2). We see that $R(A_x^-)$ is orthogonal to $N(A_x)$.

Another choice of matrices A_x^- and Z_x^- can be made when a separation of variables occurs naturally, as in optimal control problems or in parameter identification problems. This is also the framework adopted to introduce the GRG method (Abadie and Carpentier (1969)) and could be called the *partitioned framework*: A_x is supposed to be partitioned in two submatrices

$$A_x = [C_x \ D_x], \quad (3.5)$$

where the $m \times m$ matrix C_x is nonsingular and D_x has dimension $m \times (n-m)$. The right inverse A_x^- is then taken as

$$A_x^- = \begin{bmatrix} C_x^{-1} \\ 0 \end{bmatrix} \quad (3.6)$$

and the basis of $N(A_x)$ is

$$Z_x^- = \begin{bmatrix} -C_x^{-1} D_x \\ I_{n-m} \end{bmatrix}. \quad (3.7)$$

Then $Z_x = [0 \ I_{n-m}]$ is the unique matrix satisfying (3.1) and (3.2).

In the following, we shall suppose that the choice of (A_x^-, Z_x^-) is a smooth function of x :

assumption B:

the function $x \rightarrow (A_x^-, Z_x^-, Z_x)$ is bounded on ω and is in $C_b^{\nu-1}(\omega)$ with $\nu \geq 3$.

This assumption is satisfied for A_x^- given by (3.4). With regard to Z_x^- , the question is trickier, although the assumption can be satisfied locally by projection on $N(A_x)$ of a basis Z_x^- of $N(A_*)$ (see Gill, Murray, Saunders and Wright (1985) and Byrd and Schnabel (1986)).

4. A reduced quasi-Newton method for constrained optimization

In this section, we apply the results of Section 2 to constrained optimization. The first step consists in reducing the size of the optimality system (1.3). This can be done because the second optimality condition can be expressed by $n-m$ equations rather than n , in fact, by the vanishment of the $n-m$ coordinates of the orthogonal projection of $\nabla f(x_*)$ on $N(A_*)$. If Z_x^- is any basis of $N(A_*)$, the orthogonal projector on $N(A_*)$ is $Z_x^- (Z_x^{-T} Z_x^-)^{-1} Z_x^{-T}$. Then, the second equation of (1.3) is projected on $N(A_*)$ by multiplying it by Z_x^{-T} . Using the definition (1.7) of the reduced gradient, the system (1.3) can be rewritten as follows:

$$\begin{cases} c(x_*) = 0, \\ g(x_*) = 0. \end{cases} \quad (4.1)$$

In order to apply the previous results, we need to calculate the first derivative of g at x_* . This can be done as in Stoer (1984) and Nocedal and Overton (1985):

$$\nabla g(x_*) = \nabla (Z_x^- T (\nabla f(x) + A_x^T \lambda_*))(x_*) = Z_x^{-T} L_*. \quad (4.2)$$

The Jacobian matrix of (4.1),

$$\begin{bmatrix} A_* \\ Z_x^{-T} L_* \end{bmatrix},$$

is nonsingular because of the surjectivity of A_* and the second order sufficient condition which is equivalent to the nonsingularity of

$$G_* := Z_x^{-T} L_* Z_x^-. \quad (4.3)$$

We shall denote by $H_* := G_*^{-1}$.

Now, let us apply algorithm (2.4)–(2.5) to the system (4.1). By using statement (iii) of proposition 2.1 and (4.2), we get the following q-quadratically convergent algorithm:

$$x_{k+1} = x_k - [I - Z_*^- H_* Z_*^{-T} L_*] A_*^- c(x_k) - Z_*^- H_* g(x_k), \quad (4.4)$$

where A_*^- is any right inverse of A_* (playing the role of A_*^- in (2.9)) and Z_*^- is any basis of $N(A_*)$. This is exactly the "ideal" SQP method (see (1.6)). So, we obtain a result of Goodman (1985) according to which the SQP method is a Newton method for solving (4.1). We shall not go further with that method.

If we apply algorithm (2.6)–(2.8) to the system (4.1), we get, by using statement (iii) of proposition 2.2 and (4.2):

$$\bar{y}_k = x_k - A_*^- c(x_k), \quad (4.5)$$

$$\bar{x}_{k+1} = \bar{y}_k - Z_*^- G_*^{-1} g(\bar{y}_k), \quad (4.6)$$

where A_*^- is any right inverse of A_* and Z_*^- is any basis of $N(A_*)$. The following lemma is a consequence of proposition 2.2.

Lemma 4.1. Suppose that assumptions A and B are satisfied and let x_* be a solution of (1.1). There exists a positive constant C that depends only on f and c such that if x_k, \bar{y}_k given by (4.5) and \bar{x}_{k+1} given by (4.6) are in ω , we have

$$\|\bar{x}_{k+1} - x_*\| \leq C \|\bar{x}_k - x_*\|^2. \quad (4.7)$$

From the q-quadratically convergent algorithm (4.5)–(4.6), a quasi-Newton method is easily introduced. In (4.6), G_* is replaced by an approximation G_k and Z_*^- is replaced by $Z(y_k)^-$ which intervenes in the calculation of the reduced gradient $g(y_k)$. If A_*^- in (4.5) is replaced by $A(x_k)^-$, the constraints will have to be linearized twice per iteration: at x_k to calculate $A(x_k)^-$ and at y_k to calculate the basis $Z(y_k)^-$. Since the constraints have to be linearized at y_k to calculate the reduced gradient in (4.6), we avoid two linearizations of the constraints by replacing A_*^- in (4.5) by $A(y_{k-1})^-$. So we obtain the following local algorithm:

$$y_k = x_k - A(y_{k-1})^- c(x_k), \quad (4.8)$$

$$x_{k+1} = y_k - Z(y_k)^- G_k^{-1} g(y_k). \quad (4.9)$$

We shall denote by r_k the restoration step and t_k the tangent step:

$$r_k := -A(y_{k-1})^- c(x_k), \quad (4.10)$$

$$t_k := -Z(y_k)^- G_k^{-1} g(y_k). \quad (4.11)$$

We shall also use the total displacements

$$d_k := r_k + t_k, \quad (4.12)$$

$$e_k := t_k + r_{k+1}. \quad (4.13)$$

Practically, the algorithm cannot start with (4.8) from a point x_1 without knowing a point y_0 . So, we shall suppose in the following that the algorithm starts with (4.9) from a point y_0 in ω .

This is really the same type of algorithm as Coleman and Conn's method (1.15)–(1.17) if we exchange in (4.8)–(4.9) y_k with x_k and x_{k+1} with $x_k + t_k$. However, our point of view shows that there is no reason to take a restoration step orthogonal to $N(A(y_{k-1}))$ in (4.10) or to calculate an orthogonal basis $Z(y_k)^-$ of $N(A(y_k))$ in (4.11). In particular, this validates the use of the partitioned framework (3.5)–(3.7) that occurs often in practice. We will see that contrary to the sequence (y_k) in (4.8)–(4.9), which does not usually converge q-superlinearly in one step (see the examples of Byrd (1985) and Yuan (1985)), the sequence (x_k) will converge q-superlinearly as expected from the behavior of the ideal algorithm (4.5)–(4.6).

In fact, it is not essential to reduce the size of the optimality system before applying algorithm (2.6)–(2.8). The same method (4.8)–(4.9) can be obtained when the method is applied to the optimality conditions (1.3). In this case, $B_* = [L_* A_*^T]$ and

$$B_*^- = \begin{bmatrix} Z_*^- G_*^{-1} Z_*^{-T} \\ A_*^{-T} (I - L_* Z_*^- G_*^{-1} Z_*^{-T}) \end{bmatrix},$$

where A_*^- is any right inverse of A_* and Z_*^- is any basis of $N(A_*)$. Furthermore, that way gives an iteration scheme for the Lagrange multipliers (λ_k) :

$$\lambda_{k+1} = -A(y_k)^{-T} \nabla f(y_k) + A(y_k)^{-T} L_k Z(y_k)^- G_k^{-1} g(y_k), \quad (4.14)$$

where L_k is an approximation of L_* . Therefore, if L_k in (4.14) is correctly updated, we shall have also $\lambda_{k+1} - \lambda_* = o(\|x_k - x_*\|)$; indeed, λ_k does not intervene in the iteration. The formula (4.14) simplifies the one obtained when the quasi-Newton method is applied to (1.3), which writes (see Gabay (1982,b)):

$$\lambda_{k+1} = -A(x_k)^{-T} [I - L_k Z(x_k)^- G_k^{-1} Z(x_k)^{-T}] [\nabla f(x_k) - L_k A(x_k)^- c(x_k)].$$

The algorithm (4.8)–(4.9) is a *reduced quasi-Newton method* because the only matrix to update is the approximation G_k of G_* and it is of order $n-m$. Unfortunately, this is not the case any more when the sequence (λ_k) is generated by (4.14) since L_k intervenes in the formula and L_k is of order n .

In the next two propositions, we study the local linear convergence of algorithm (4.8)–(4.9).

Lemma 4.2. Suppose that assumptions *A* and *B* are satisfied. There exist positive constants $\bar{\delta}$, C_1 , C_2 and C_3 that depend only on f , c and ω such that on the one hand, $\|G_k - G_*\| \leq \bar{\delta}$ implies that G_k is nonsingular with

$\|G_k^{-1}\| \leq C_3$ and on the other hand,

$$\|y_{k-1} - x_*\| \leq (1 + C_1) \delta, \quad (4.15)$$

$$\|x_k - x_*\| \leq \delta, \quad (4.16)$$

$$\|G_k - G_*\| \leq \delta \quad (4.17)$$

with $0 < \delta \leq \bar{\delta}$ imply that y_k and x_{k+1} are well defined by (4.8)–(4.9) and satisfy

$$\|y_k - x_*\| \leq (1 + C_1) \delta, \quad (4.18)$$

$$\|x_{k+1} - x_*\| \leq C_2 \delta \|x_k - x_*\|. \quad (4.19)$$

Proof. We shall denote by C_i ($i=1,2,\dots$) any positive constant that depends only on f, c and ω . According to assumptions A and B , there exists a positive constant C_1 such that

$$\max (\|A(y)\|, \|A(y)^-\|, \|Z(y)^-\|, \|g'(y)\|) \leq C_1, \quad (4.20)$$

for all y in ω and

$$\max (\|\bar{y}_k - x_*\|, \|y_k - x_*\|) \leq (1 + C_1) \|x_k - x_*\|, \quad (4.21)$$

if y_{k-1} and x_k are in ω and \bar{y}_k and y_k are calculated from x_k by (4.5) and (4.8) respectively. Both inequalities in (4.21) are obtained from (4.5) and (4.8) by using Taylor's expansion on $c(x_k)$. Let ϵ , η and $\bar{\delta}$ be three fixed positive constants such that

$$B(x_*, \epsilon) \subset \omega, \quad (4.22)$$

$$\|G_*^{-1}\| \leq \eta, \quad (4.23)$$

$$\bar{\delta} < \frac{1}{\eta}, \quad (4.24)$$

$$(1 + C_1) \bar{\delta} < \epsilon, \quad (4.25)$$

where $B(x_*, \epsilon)$ denotes the ball of radius ϵ centred at x_* . These constants ϵ , η and $\bar{\delta}$ depend only on f , c and ω . If G_k satisfies $\|G_k - G_*\| \leq \bar{\delta}$ then, by (4.24) and (4.23), G_k is nonsingular and satisfies (see for example Schwartz (1981), chap. II, theorem 62):

$$\|G_k^{-1}\| \leq \frac{1}{\frac{1}{\eta} - \bar{\delta}} =: C_3. \quad (4.26)$$

This proves the first part of the lemma. For the second part, let us suppose that (4.15)–(4.17) are satisfied with δ in $]0, \bar{\delta}]$. According to (4.15), (4.25) and (4.22), y_{k-1} belongs to ω and according to (4.16), (4.25) and (4.22), x_k belongs to ω . Therefore y_k is

well defined by (4.8) and we have (4.21). This inequality and (4.16) show (4.18). Now, according to (4.21), (4.16), (4.25) and (4.22), y_k and \bar{y}_k belong to ω . So, \bar{x}_{k+1} and x_{k+1} are well defined by (4.6) and (4.9) respectively. From (4.26), (4.23), (4.17) and $G_k^{-1} - G_*^{-1} = G_k^{-1} (G_* - G_k) G_*^{-1}$, we deduce:

$$\|G_k^{-1} - G_*^{-1}\| \leq \|G_k^{-1}\| \|G_*^{-1}\| \|G_k - G_*\| \leq C_4 \delta.$$

Let C_5 be the constant given by lemma 4.1. Then, Taylor expansions give easily the following inequalities:

$$\|x_{k+1} - x_*\| \leq \|x_{k+1} - \bar{x}_{k+1}\| + \|\bar{x}_{k+1} - x_*\|,$$

$$\begin{aligned} \|x_{k+1} - \bar{x}_{k+1}\| &\leq C_6 \|y_k - \bar{y}_k\| + C_7 \|y_k - x_*\|^2 \\ &\quad + C_4 C_8 \delta \|y_k - x_*\|, \end{aligned}$$

$$\|y_k - \bar{y}_k\| \leq C_9 \|y_{k-1} - x_*\| \|x_k - x_*\|.$$

By combining these inequalities with (4.7), (4.15), (4.16) and (4.21), we get (4.19) with $C_2 = C_5 + C_6 C_9(1+C_1) + C_7(1+C_1)^2 + C_4 C_8(1+C_1)$. \square

Theorem 4.3. Suppose that assumptions *A* and *B* are satisfied. There exists a positive constant C that depends only on f , c and ω such that if κ is a real number in $]0, 1[$ and if

$$\|y_0 - x_*\| \leq C \kappa, \tag{4.27}$$

$$\|G_k - G_*\| \leq C \kappa \text{ for all subscripts } k, \tag{4.28}$$

then algorithm (4.8)–(4.9) generates from y_0 a sequence (x_k) in ω that converges q-linearly to x_* and

$$\|x_{k+1} - x_*\| \leq \kappa \|x_k - x_*\| \tag{4.29}$$

for all subscripts k .

Proof. Let $\bar{\delta}$, C_1 , C_2 and C_3 be the positive constants given by lemma 4.2. Then, if G_{k-1} satisfies $\|G_{k-1} - G_*\| \leq \bar{\delta}$, we have $\|G_{k-1}^{-1}\| \leq C_3$. By expanding $g(y_{k-1})$ about x_* (with (4.2)), (4.9) shows that for $k \geq 1$, we have

$$\|x_k - x_*\| \leq (1+C_4) \|y_{k-1} - x_*\|, \tag{4.30}$$

where C_4 is a positive constant that depends only on f , c and ω . Then, the theorem can be proved with $C := \min(\bar{\delta}, 1/C_2)/(1+C_4)$. Indeed, if (4.27) and (4.28) are satisfied, we see, with the help of (4.30), that (4.15)–(4.17) are satisfied for $k = 1$ and $\delta := (1+C_4) C \kappa \leq \bar{\delta}$:

$$\|y_0 - x_*\| \leq C \kappa \leq \delta \leq (1 + C_1)\delta,$$

$$\|x_1 - x_*\| \leq (1 + C_4) \|y_0 - x_*\| \leq \delta,$$

$$\|G_1 - G_*\| \leq C\kappa \leq \delta.$$

We can then apply lemma 4.2 and because $C_2 \delta \leq \kappa$, (4.19) shows that (4.29) is satisfied for $k=1$. The fact that κ is less than 1, (4.28) and (4.18) for $k=1$ show that (4.15)–(4.17) are still satisfied for $k=2$. So we can conclude by induction. \square

The next proposition gives some useful estimates and equivalences.

Proposition 4.4. Suppose that assumptions *A* and *B* are satisfied. Let (G_k) be a sequence of nonsingular matrices of order $n-m$. Let (x_k) in ω and (y_k) in ω be the sequences generated by algorithm (4.8)–(4.9) starting from a point y_0 in ω . If (x_k) and (y_k) converge to a solution x_* of (1.1), we have

$$r_k = -A_*^- A_* (x_k - x_*) + o(\|x_k - x_*\|), \quad (4.31)$$

$$y_k - x_* = Z_*^- Z_* (x_k - x_*) + o(\|x_k - x_*\|). \quad (4.32)$$

If moreover (G_k) and (G_k^{-1}) are bounded, we have

$$\|d_k\| \sim \|x_k - x_*\|, \quad (4.33)$$

$$\|e_k\| \sim \|y_k - x_*\|. \quad (4.34)$$

Proof. From the definition (4.10) of r_k , the expansion of $c(x_k)$ about x_* and the convergence of (y_k) , we get (4.31). Then, by using the identity (3.3) and (4.31) on $y_k = x_k + r_k$, we obtain (4.32). Next, by using the boundedness of (G_k^{-1}) and (4.2), we see that $t_k = -Z_*^- G_k^{-1} Z_*^T L_*(y_k - x_*) + o(\|y_k - x_*\|)$. Finally, with (4.12), (4.31) and (4.32), this gives

$$d_k = -[A_*^- A_* + Z_*^- G_k^{-1} G_* Z_*] (x_k - x_*) + o(\|x_k - x_*\|).$$

This estimate shows that $d_k = 0(\|x_k - x_*\|)$. To prove the converse, we only have to show that the operator in square brackets is nonsingular with bounded inverse. If this were not the case, there would exist a subsequence \mathcal{K} of subscripts and a sequence $(\xi_k: k \in \mathcal{K})$ in \mathbb{R}^n such that

$$\|\xi_k\| = 1 \text{ for } k \text{ in } \mathcal{K}, \quad (4.35)$$

$$[A_*^- A_* + Z_*^- G_k^{-1} G_* Z_*] \xi_k \rightarrow 0 \text{ for } k \text{ in } \mathcal{K}. \quad (4.36)$$

By multiplying (4.36) by A_* (resp. Z_*), we should obtain $A_* \xi_k \rightarrow 0$ (resp. $G_k^{-1} G_* Z_* \xi_k \rightarrow 0$), from which we should deduce $Z_* \xi_k \rightarrow 0$ because of the

boundedness of (G_k) and the nonsingularity of G_* . Finally, with (3.3), we should have $\xi_k \rightarrow 0$, which would contradict (4.35). So, (4.33) is proved. The proof of (4.34) is similar and is based on the estimate

$$e_k = - [A_*^- A_* + Z_*^- G_k^{-1} Z_*^{-T} L_*] (y_k - x_*) + o(\|y_k - x_*\|). \quad \square$$

5. Conditions for q-superlinear convergence

Theorem 4.3 has an immediate corollary which states that if in addition to (4.28), the sequence (G_k) converges to G_* then (x_k) converges to x_* q-superlinearly (see for example the argument in the proof of corollary 3.5 in Han (1976)). However, this assumption on (G_k) is usually not satisfied when these matrices are generated by quasi-Newton formulas. Assuming that (x_k) converges to x_* , the next theorem gives necessary and sufficient conditions on (G_k) in order to have the q-superlinear convergence of (x_k) . It is the analogue of theorem 2.2 of Denis and Moré (1974) valid for quasi-Newton methods in optimization without constraints.

Theorem 5.1. Suppose that assumptions *A* and *B* are satisfied and that (y_k) and (x_k) are generated in ω from a point y_0 by algorithm (4.8)–(4.9) with a sequence (G_k) of nonsingular matrices. Suppose that (x_k) and (y_k) converge to x_* . Then, the following statements are equivalent:

- (i) (x_k) converges q-superlinearly,
- (ii) $g(y_{k+1}) = o(\|x_k - x_*\|)$,
- (iii) $(G_k - G_*) Z(y_k) t_k = o(\|x_k - x_*\|)$.

Proof. The estimate (4.32) shows that

$$y_k - x_* = O(\|x_k - x_*\|). \quad (5.1)$$

It is then easy to get (we use $A(y_k) - A_* = O(\|y_k - x_*\|)$, $t_k \rightarrow 0$, (4.32) and $A_* Z_*^- = 0$):

$$\begin{aligned} A_*(x_{k+1} - x_*) &= A_*(y_k - x_*) + A(y_k)t_k + o(\|y_k - x_*\|) \\ &= o(\|x_k - x_*\|). \end{aligned} \quad (5.2)$$

According to (3.3), it remains to estimate $Z_*(x_{k+1} - x_*)$. This will depend on the quality of the tangent step t_k .

Let us first prove the equivalence (i) \iff (ii). With (4.2) and (4.32), we have

$$g(y_{k+1}) = Z_*^{-T} L_* (y_{k+1} - x_*) + o(\|y_{k+1} - x_*\|)$$

$$= G_* Z_* (x_{k+1} - x_*) + o(\|x_{k+1} - x_*\|).$$

Then, (ii) is clear from (i). If (ii) is satisfied, this estimate and the nonsingularity of G_* give

$$Z_*(x_{k+1} - x_*) = o(\|x_{k+1} - x_*\|) + o(\|x_k - x_*\|).$$

This estimate, (5.2) and the identity (3.3) show (i).

Now, let us show that in any of the situations (i), (ii) or (iii), we have

$$t_k = o(\|x_k - x_*\|). \quad (5.3)$$

This estimate is clear when (G_k^{-1}) is bounded, but we do not suppose this here. By writing $t_k = (x_{k+1} - x_*) - (y_k - x_*)$ and by using (5.1), we see that (5.3) is clearly satisfied when (i) is true and therefore when (ii) is true. When (iii) is satisfied, we have

$$G_* Z(y_k) t_k = G_k Z(y_k) t_k + o(\|x_k - x_*\|) = -g(y_k) + o(\|x_k - x_*\|).$$

Then by expanding $g(y_k)$ about x_* and by using (5.1) and the nonsingularity of G_* , we get

$$Z(y_k) t_k = o(\|x_k - x_*\|).$$

But $t_k = Z(y_k)^{-1} Z(y_k) t_k$, therefore (5.3) is still satisfied. Now, from (5.1) and (5.3), it follows that $x_{k+1} - x_* = o(\|x_k - x_*\|)$, $y_{k+1} - y_k = o(\|x_k - x_*\|)$ and, with (4.31) and (5.2),

$$r_{k+1} = o(\|x_k - x_*\|). \quad (5.4)$$

Let us now prove the equivalence (ii) \iff (iii). By expanding $g(y_{k+1})$ about y_k and by using (4.2), (5.3) and (5.4), we have

$$g(y_{k+1}) = g(y_k) + Z_*^{-T} L_* t_k + o(\|x_k - x_*\|).$$

But $g(y_k) = -G_k Z(y_k) t_k$ and $t_k = Z(y_k)^{-1} Z(y_k) t_k = Z_*^{-1} Z(y_k) t_k + o(\|x_k - x_*\|)$. So, we obtain

$$g(y_{k+1}) = -(G_k - G_*) Z(y_k) t_k + o(\|x_k - x_*\|).$$

The equivalence (ii) \iff (iii) follows. \square

In the statement (ii) of theorem 5.1, $g(y_{k+1})$ could be replaced by $g(x_{k+1})$, but the reduced gradient is not evaluated at x_{k+1} in the algorithm. The statement (iii) is equivalent to

$$(G_k^{-1} - G_*^{-1})g(y_k) = o(\|x_k - x_*\|)$$

which is based on the gap between the inverse of the Hessians. The statement (iii) can

also be replaced by many other equivalent estimates. For example,

$$(G_k - G_*) Z_*(x_k - x_*) = o(\|x_k - x_*\|).$$

The advantage of (iii) is that it does not require the boundedness of the sequences (G_k) or (G_k^{-1}) . If this boundedness is assumed, proposition 4.4 shows that the estimates can be done in relation to $\|d_k\|$ rather than $\|x_k - x_*\|$.

Condition (4.28) and condition (iii) of theorem 5.1 show the advantage of the reduced quasi-Newton methods over the SQP method with regard to the approximation of the Hessian of the Lagrangian. Indeed, a necessary and sufficient condition for the SQP method to generate q-superlinearly convergent sequences is that

$$Z_*^{-T} (L_k - L_*) (x_k - x_*) = o(\|x_k - x_*\|),$$

where L_k is the updated approximation of L_* . This famous result can be found in Boggs, Tolle and Wang (1982) and Nocedal and Overton (1985). Therefore, in the SQP method, the $(n-m) \times n$ matrix $Z_*^{-T} L_*$ has to be correctly approximated and not only the projected Hessian of the Lagrangian $Z_*^{-T} L_* Z_*^{-}$ as in reduced methods.

As a final remark, let us mention that, if we suppose that the equivalence (4.34) holds, the sufficient condition given by Byrd (1984) writes

$$(G_k - G_*) Z(y_k) t_k = o(\|y_k - x_*\|) \quad (5.5)$$

and is therefore stronger than condition (iii) of theorem 5.1. However, this estimate (5.5) is satisfied in practice with the update schemes currently proposed: Coleman and Conn (1984, theorem 3.6) proved it for their algorithm while it is proved in Gilbert (1986,a) that

$$(G_k - G_*) Z(y_k) t_k = o(\|t_k\|) \quad (5.6)$$

(which implies (5.5)) holds for both of the algorithms proposed. But those schemes are not completely satisfactory and condition (iii) of theorem 5.1 may become useful in other circumstances. We shall see in the next section that, for the globalizing technique proposed, the strongest condition (5.6) allows avoidance of the Maratos effect, i.e. allows a unit step-size parameter to be accepted asymptotically.

6. Globalization of the algorithm

In order to globalize the local algorithm (4.8)–(4.9), we introduce a step-size parameter ρ_k . For that, we consider the following *exact penalty function*:

$$\Theta_p(x) = f(x) + p \|c(x)\|_1, \quad (6.1)$$

where p is the positive *penalty parameter* and $\|\bullet\|_1$ is the l_1 -norm on \mathbb{R}^m . Other norms than the l_1 -norm can be used in (6.1): see Bonnans and Gabay (1984). If p is taken greater than $\|\lambda_*\|_\infty$ (where $\|\bullet\|_\infty$ is the sup-norm on \mathbb{R}^m), feasible minimizers of (1.1) and (6.1) are the same (see Fletcher (1981) for example). It is therefore natural to look for x_* by minimizing Θ_p . For this, we need to calculate descent directions of this non-differentiable function. On that point, a crucial observation has been made by Han (1977): the displacement d_k^{SQP} of the SQP method is a descent direction of Θ_p at x_k (if some natural hypotheses are satisfied). Therefore a better approximation x_{k+1} of the solution x_* will be obtained by taking

$$x_{k+1} = x_k + \rho_k d_k^{SQP},$$

where ρ_k gives the step-size and is obtained from some rule using Θ_p as a "merit" function.

Let us try to use the same globalizing technique for our algorithm. Is there any descent direction of Θ_p among the displacements r_k , t_k , d_k , and e_k given by (4.10)–(4.13)? The inconvenience of r_k and therefore of d_k and e_k is that this displacement is calculated by using two different points y_{k-1} and x_k that can be far from each other when x_k is far from x_* . So, it is difficult to see when those directions are descent directions for Θ_p . On the other hand, t_k uses only the point y_k in its definition and if G_k is positive definite, it is certainly a descent direction of Θ_p at y_k . Indeed, this displacement is tangent to $c^{-1}(c(y_k))$ at y_k and $f'(y_k) \cdot t_k$ is negative. Therefore at the first order, the first term of the right hand side of (6.1) will decrease while the second term will remain constant. These remarks lead us to define a *descent arc* of Θ_p at y_k , tangent to t_k :

$$y_k(\rho) = y_k + \rho t_k + \rho^a r_{k+1}, \quad a > 1. \quad (6.2)$$

Let us note that search arcs have already been proposed by Mayne and Polak (1982) to cope with the Maratos effect of the SQP method (see further) and by Gabay (1982,b), also to avoid the Maratos effect for his algorithm.

This globalizing technique based on the arc (6.2) gives priority to the minimizing step t_k ; and this is due to the asymmetry of the local method (4.8)–(4.9). This priority can be harmful in certain circumstances but it can be suppressed by adding a restoration step to the local method (see Gilbert (1986,b)).

The point y_{k+1} is then obtained from y_k by selecting a particular value ρ_k of ρ :

$$y_{k+1} := y_k(\rho_k). \quad (6.3)$$

The step-size ρ_k will be determined here so that the following Armijo-like criterion will be satisfied:

$$\beta \in]0,1[, \quad (6.4)$$

$$\rho_k := \beta^{l_k}, \quad (6.5)$$

where l_k is the smallest non-negative integer such that

$$\begin{aligned} \Theta_p(y_k(\beta^{l_k})) &\leq \Theta_p(y_k) + \beta^{l_k} \alpha f'(y_k) \cdot t_k \\ &\quad - \beta^{a l_k} \alpha (p - \|\lambda(y_k)\|_\infty) \|c(y_k)\|_1. \end{aligned} \quad (6.6)$$

In this inequality, α is a real number chosen in $]0, 1/2[$ for reasons that will be clear at the end of this section. The exponent $(a l_k)$ of β in the last term of (6.6) takes into account the curvature of the search path (6.2). The vector $\lambda(y_k)$ is the approximation at y_k of the Lagrange multiplier λ_* defined by

$$\lambda(y) := -A(y)^{-T} \nabla f(y). \quad (6.7)$$

It is just the first term of (4.14). So, usually, $(\lambda(y_k))$ will not converge superlinearly. We shall define again

$$x_{k+1} := y_k + t_k. \quad (6.8)$$

Now, we have to examine in what conditions the inequality (6.6) can be realized with a large enough l_k . This is the subject of the following lemma.

Lemma 6.1. Suppose that assumptions *A* and *B* are satisfied and that a point y_k is given in ω such that x_{k+1} and $x_{k+1} + r_{k+1}$ will also be in ω . Suppose that α is in $]0, 1[$ and that there exists positive constants \underline{p} , \bar{p} and h such that

$$\begin{aligned} \underline{p} + \|\lambda(y_k)\|_\infty &\leq p \leq \bar{p}, \\ h I &\leq G_k^{-1} \leq h^{-1} I. \end{aligned} \quad (6.9)$$

Then the rule (6.4)–(6.6) allows determination of a positive step-size ρ_k . If, moreover, M is a positive constant such that

$$\|c(y_k)\|_1 \leq M, \quad (6.10)$$

then, there exists a positive real $\underline{\rho}$ that depends only on f , c , \underline{p} , \bar{p} , h , α , β and M such that

$$\rho_k \geq \underline{\rho} > 0.$$

Proof. We shall denote by C_i ($i=1,2,\dots$) any positive constant. Using $c'(y_k) \cdot t_k = 0$, Taylor's theorem gives

$$\|c(x_{k+1}) - c(y_k)\|_1 \leq C_1 \|t_k\|^2, \quad (6.11)$$

where C_1 depends only on c . Using (6.11) and $0 < \rho \leq 1$ and applying again Taylor's

theorem, we get

$$\begin{aligned} f(y_k + \rho t_k + \rho^\alpha r_{k+1}) &\leq f(y_k) + \rho f'(y_k) \cdot t_k + \rho^\alpha (\lambda(y_k), c(y_k)) \\ &\quad + \rho^\alpha C_2 \|t_k\|^2 + C_3 \|\rho t_k + \rho^\alpha r_{k+1}\|^2, \end{aligned} \quad (6.12)$$

$$\begin{aligned} \|c(y_k + \rho t_k + \rho^\alpha r_{k+1})\|_1 &\leq (1 - \rho^\alpha) \|c(y_k)\|_1 \\ &\quad + \rho^\alpha C_1 \|t_k\|^2 + C_4 \|\rho t_k + \rho^\alpha r_{k+1}\|^2, \end{aligned} \quad (6.13)$$

where C_2 , C_3 and C_4 depend only on f and c . Supposing ρ in $]0, 1]$, we get from (6.12) and (6.13):

$$\begin{aligned} \Theta_p(y_k + \rho t_k + \rho^\alpha r_{k+1}) &\leq \Theta_p(y_k) + \rho f'(y_k) \cdot t_k \\ &\quad - \rho^\alpha (p - \|\lambda(y_k)\|_\infty) \|c(y_k)\|_1 \\ &\quad + C_5 (\rho^\alpha + \rho^2) \|t_k\|^2 + C_6 \rho^{2\alpha} \|r_{k+1}\|^2, \end{aligned} \quad (6.14)$$

where C_5 and C_6 depend only on f , c and \bar{p} . From the definition (4.10) of r_{k+1} and (6.11), we have

$$\|r_{k+1}\| \leq C_7 \|c(y_k)\|_1 + C_8 \|t_k\|^2, \quad (6.15)$$

where C_7 and C_8 depend only on c .

Now, let us suppose that (6.6) is not true for a given $\rho = \beta^l$ in $]0, 1]$. Then, with (6.9), (6.14) and (6.15), we get

$$\begin{aligned} \rho (1 - \alpha) (-f'(y_k) \cdot t_k) + \rho^\alpha (1 - \alpha) \underline{p} \|c(y_k)\|_1 \\ \leq C_5 (\rho^\alpha + \rho^2) \|t_k\|^2 + C_6 \rho^{2\alpha} \|r_{k+1}\|^2. \end{aligned}$$

But $-f'(y_k) \cdot t_k = (G_k^{-1} g(y_k), g(y_k)) \geq h \|g(y_k)\|^2$ and $\|t_k\| \leq C_9 \|g(y_k)\|$ where C_9 depends only on c and h . Then, with (6.15), the last inequality becomes

$$\begin{aligned} \rho \|g(y_k)\|^2 + \rho^\alpha \|c(y_k)\|_1 \\ \leq C_{10} (\rho^\alpha + \rho^2) \|g(y_k)\|^2 + C_{11} \rho^{2\alpha} \|c(y_k)\|_1^2, \end{aligned} \quad (6.16)$$

where C_{10} and C_{11} depend only on f , c , \underline{p} , \bar{p} , h and α . This inequality shows that ρ cannot be arbitrarily small if $\|g(y_k)\| + \|c(y_k)\|_1 \neq 0$ (if $\|g(y_k)\| + \|c(y_k)\|_1 = 0$, $\rho = \beta^l = 1$ clearly satisfies (6.6) because then $t_k = 0$, $r_{k+1} = 0$ and therefore $y_k(\rho) = y_k$). Indeed, otherwise letting ρ converged to 0 in (6.16) previously divided by ρ , we should obtain $g(y_k) = 0$ and then dividing (6.16) by ρ^α and taking the limit on ρ would give $c(y_k) = 0$. This proves the first part of the lemma.

For the second part, let us suppose that the rule (6.4)–(6.6) gives a step-size ρ_k smaller than 1. Then (6.6) is not satisfied with $\rho = \rho_k/\beta$ and the inequality (6.16) gives with (6.10):

$$\begin{aligned} & \rho_k \|g(y_k)\|^2 + \rho_k^a \|c(y_k)\|_1 \\ & \leq C_{12} \rho_k^b [\rho_k \|g(y_k)\|^2 + \rho_k^a \|c(y_k)\|_1], \end{aligned}$$

where $b := \min(1, a-1)$ and C_{12} depends only on $f, c, \underline{p}, \bar{p}, \alpha, \beta, h$ and M . Because $\rho_k \|g(y_k)\|^2 + \rho_k^a \|c(y_k)\|_1 \neq 0$ (otherwise $\rho_k = 1$) the last inequality proves the second part of the lemma with $\underline{\rho} = C_{12}^{-1/b}$. \square

The inequality (6.9) shows that the penalty parameter p has to be large enough to ensure the decrease of Θ_p along the arc (6.2) and that its lower bound depends on the current point y_k . So, sometimes it will be necessary to update the penalty parameter, which we shall denote by p_k . We shall suppose that the adapting rule of p_k will satisfy the following three conditions:

$$p_k \geq \|\lambda(y_k)\|_\infty + \underline{p}, \text{ for every } k, \quad (6.17)$$

$$\begin{aligned} & \text{there exists a subscript } K \text{ such that for every } k \text{ greater than } K, \\ & (p_{k-1} \geq \|\lambda(y_k)\|_\infty + \underline{p}) \text{ implies that } p_k = p_{k-1}, \end{aligned} \quad (6.18)$$

$$(p_k) \text{ is bounded if and only if } p_k \text{ is modified finitely often.} \quad (6.19)$$

In (6.17) and (6.18), \underline{p} is a given positive constant. The condition (6.18) means that finally (for $k \geq K$), p_k is modified only if it is necessary in order to have (6.17). So $(p_k: k \geq K)$ is an increasing sequence. An example of an adapting rule satisfying these conditions is given by Mayne and Polak (1982):

$$\begin{aligned} & \text{if } p_{k-1} \geq \|\lambda(y_k)\|_\infty + \underline{p}, \\ & \text{then } p_k := p_{k-1}, \\ & \text{else } p_k := \max(\delta p_{k-1}, \|\lambda(y_k)\|_\infty + \underline{p}), \end{aligned}$$

where δ is a given constant greater than 1 in order to satisfy (6.19).

We are now able to state the algorithm which globalizes the local method (4.8)–(4.9).

Algorithm RQN:

1. Choose a convergence tolerance $\epsilon > 0$, β in $]0,1[$, α in $]0,1/2[$ and $a > 1$.
2. Choose y_0 in ω and a symmetric positive definite matrix G_0 of order $n-m$.
3. Let $k := 0$.

4. Repeat:
 - 4.1. Linearize the constraints at y_k : choose a right inverse $A(y_k)^-$ of $\nabla c(y_k)$ and a basis $Z(y_k)^-$ of $N(\nabla c(y_k))$ according to assumption B .
 - 4.2. Evaluate $\lambda(y_k) := -A(y_k)^{-T} \nabla f(y_k)$ and $g(y_k) := Z(y_k)^{-T} \nabla f(y_k)$.
 - 4.3. If $k \geq 1$ then evaluate the symmetric positive definite matrix G_k by updating G_{k-1} .
 - 4.4. Tangent step: evaluate $t_k := -Z(y_k)^- G_k^{-1} g(y_k)$ and $x_{k+1} := y_k + t_k$.
 - 4.5. Restoration step: evaluate $c(x_{k+1})$ and $r_{k+1} := -A(y_k)^- c(x_{k+1})$.
 - 4.6. If $\|g(y_k)\| + \|c(x_{k+1})\| < \epsilon$ then stop.
 - 4.7. Adapt p_k according to (6.17)–(6.19).
 - 4.8. Search a point y_{k+1} from y_k along the arc (6.2) in order to decrease the penalty function (6.1) (with $p = p_k$) with the help of the rule (6.3)–(6.6).
 - 4.9. Next iteration: set $k := k+1$.

In the partitioned framework (see Section 3), only one linear system has to be solved at the step 4.2. Indeed, if $A(y_k) = [C(y_k) \ D(y_k)]$, $\lambda(y_k)$ is obtained by solving

$$C(y_k)^T \lambda(y_k) = -\nabla f(y_k)^{(1)},$$

where $\nabla f(y_k)^{(1)}$ is the vector formed by the first m components of $\nabla f(y_k)$. Then $g(y_k) = D(y_k)^T \lambda(y_k) + \nabla f(y_k)^{(2)}$, where $\nabla f(y_k)^{(2)}$ is the vector formed by the last $n-m$ components of $\nabla f(y_k)$.

The important question of the update of the matrices (G_k), which is mentioned at step 4.3 of algorithm RQN, has been investigated by Coleman and Conn (1984) and by Gilbert (1986,a). G_{k+1} is obtained from G_k by the BFGS formula using two vectors γ_k and σ_k in \mathbb{R}^{n-m} :

$$G_{k+1} = G_k - \frac{G_k \sigma_k \sigma_k^T G_k}{(\sigma_k, G_k \sigma_k)} + \frac{\gamma_k \gamma_k^T}{(\gamma_k, \sigma_k)}. \quad (6.20)$$

Therefore, G_{k+1} satisfies the secant equation:

$$\gamma_k = G_{k+1} \sigma_k. \quad (6.21)$$

The point now is to choose adequately the vectors γ_k and σ_k so that G_{k+1} will approximate $G_* := Z_*^{-T} L_* Z_*$. This form of G_* and the formula (4.2) suggest taking γ_k as the difference of two reduced gradients. A first possible choice consists in taking (see Coleman and Conn (1984) and Gilbert (1986,a)):

$$\gamma_k = g(x_{k+1}) - g(y_k), \quad (6.22)$$

$$\sigma_k = -G_k^{-1} g(y_k). \quad (6.23)$$

Then, if we suppose that (y_k) and (x_k) converge to x_* , Taylor's theorem gives:

$$\gamma_k = G_* \sigma_k + o(\|\sigma_k\|). \quad (6.24)$$

This relation and (6.21) show that γ_k and σ_k are correctly chosen. Unfortunately, this choice needs an additional linearization of the constraints at x_{k+1} in order to calculate the reduced gradient at this point. This may be avoided by taking (see Gilbert (1986,a)):

$$\gamma_k = g(y_{k+1}) - g(y_k), \quad (6.25)$$

$$\sigma_k = -\rho_k G_k^{-1} g(y_k). \quad (6.26)$$

But in this case, (6.24) will not be necessarily satisfied anymore which means that updating G_k by formula (6.20) would deteriorate the matrix. Therefore, an update criterion of the form

$$\|\rho_k^a r_{k+1}\| \leq \mu_k \|\rho_k t_k\|, \quad (6.27)$$

where (μ_k) is an appropriate sequence converging to zero, has to be introduced. When (6.27) is satisfied, it is not too difficult to show that the estimate (6.24) is still valid with γ_k and σ_k given by (6.25) and (6.26). The crucial point is now to choose correctly the sequence (μ_k) so that when (6.27) is *not* satisfied the superlinear rate of convergence of (x_k) can be preserved. A good choice for μ_k is

$$\mu_k = \mu \|e_{k--}\|,$$

where μ is a small enough constant and $(k--)$ is the subscript of the last but one iterate at which (6.27) was satisfied and therefore G_{k--} updated by the formula (6.20).

The update scheme at step 4.3 of algorithm RQN is expected to generate a sequence of nonsingular matrices G_k satisfying

$$h I \leq G_k^{-1} \leq h^{-1} I, \quad (6.28)$$

for some positive constant h . This property is really not easy to obtain. However, by using the same type of arguments that are used in unconstrained optimization, it can be proved either in a local framework (when (x_o, G_o) is supposed to be close to (x_*, G_*) and $\rho_k = 1$) or when it is assumed that (x_k) and (y_k) converge to x_* with

$$\sum_{k=0}^{\infty} \|x_k - x_*\| < +\infty \quad \text{and} \quad \sum_{k=0}^{\infty} \|y_k - x_*\| < +\infty.$$

See Gilbert (1986,a).

The next theorem analyses the global convergence of the algorithm RQN under the hypothesis (6.28).

Theorem 6.2. Suppose that assumptions *A* and *B* are satisfied and that f is bounded from below on ω . Let (x_k) , (y_k) and (G_k) be the sequences generated by algorithm RQN with α in $]0,1[$. Suppose that (x_k) and (y_k) are in ω and that the matrices G_k are nonsingular and satisfy (6.18) with a positive constant h independent of k . Then, either (p_k) is unbounded and $(y_k: p_k \neq p_{k-1})$ has no accumulation point in ω , or (p_k) is bounded and

$$\|g(y_k)\| + \|c(y_k)\|_1 \rightarrow 0. \quad (6.29)$$

Proof. Suppose first that (p_k) is unbounded and let \mathcal{K} be the subsequence of the subscripts $k \geq K$ (K given in (6.18)) for which $p_k \neq p_{k-1}$. By (6.18),

$$p_{k-1} < \|\lambda(y_k)\|_\infty + \underline{p}$$

for k in \mathcal{K} . Because $(p_k : k \geq K)$ is an increasing sequence, we see from this inequality that $\|\lambda(y_k)\|_\infty \rightarrow \infty$ for $k \rightarrow \infty$ in \mathcal{K} . Therefore $(y_k: p_k \neq p_{k-1})$ has no accumulation point in ω (here, we use the continuity of $y \rightarrow \lambda(y)$ and so, the surjectivity of $\nabla c(y)$ and assumption *B* are strongly invoked).

Now, let us suppose that (p_k) is bounded. From (6.19), p_k is constant when k is great enough. Let us say that $p_k = p$ for $k \geq K_1$. So, at each iteration the same penalty function Θ_p decreases. The function f being bounded from below, we get

$$p \|c(y_k)\|_1 \leq \Theta_p(y_{K_1}) - \inf f, \text{ for } k \geq K_1.$$

Therefore, $(\|c(y_k)\|_1)$ is bounded and we can apply lemma 6.1, which states the existence of a positive lower bound $\underline{\rho}$ for the sequence (ρ_k) . From (6.28), we get $(-f'(y_k) \cdot t_k) \geq h \|g(y_k)\|^2$ and then, with (6.17), (6.6) writes

$$\begin{aligned} \underline{\rho} \alpha h \|g(y_k)\|^2 + \underline{\rho}^\alpha \alpha \underline{p} \|c(y_k)\|_1 \\ \leq \Theta_p(y_k) - \Theta_p(y_{k+1}), \text{ for } k \geq K_1. \end{aligned}$$

But $(\Theta_p(y_k))$ converges (a decreasing bounded from below sequence). Therefore, taking the limit on k in this inequality shows that $\|g(y_k)\|$ and $\|c(y_k)\|_1$ converge to zero, which is the expected result. \square

A last problem to tackle concerns the question of the admissibility of the unit step-size. When $\rho_k = 1$ is accepted by (6.6), the algorithm RQN proceeds like the local method (4.8)–(4.9) and q-superlinear convergence of (x_k) will occur when the reduced Hessian G_* is correctly approached by G_k (see theorem 5.1, statement (iii)). It is known that this admissibility property is not satisfied when the SQP method is globalized with the penalty function (6.1) and the technique described at the beginning of this section. This has been called the "Maratos effect" of the SQP method (see Maratos

(1978)) and several remedies have been proposed to overcome this drawback (see Gabay (1982,b), Chamberlain, Lemaréchal, Pedersen and Powell (1982), Mayne and Polak (1982) and Bonnans (1984)). This inconvenience is not shared with our algorithm. In fact, when $c(y_k)=0$ (a favorable situation for the appearance of the Maratos effect), the total displacement $e_k = t_k + r_{k+1}$ is exactly the same as the one of the SQP method with the Mayne and Polak's correction. This may explain that fact.

Let (x_k) in ω , (y_k) in ω and (G_k) be the sequences generated by the algorithm RQN and suppose that (y_k) converges to a solution x_* of (1.1). Let \mathbb{K} be a subsequence of subscripts. We are interested in finding conditions under which ρ_k will be equal to 1 for all but finitely many subscripts k in the subsequence considered. The following four properties will be meaningful:

$$\|G_k - G_*\| \leq M \text{ for } k \text{ in } \mathbb{K}, \quad (6.30)$$

$$(G_k - G_*) Z_* t_k = o(\|t_k\|) \text{ for } k \text{ in } \mathbb{K}, \quad (6.31)$$

$$t_k = O(\|r_{k+1}\|) \text{ for } k \text{ in } \mathbb{K}, \quad (6.32)$$

$$\rho_k < 1 \text{ and } t_k = o(\|r_k\|) \text{ for } k \text{ in } \mathbb{K}. \quad (6.33)$$

The properties (6.30) and (6.31) concern the approximation of the reduced Hessian G_* by G_k . The property (6.30) is very strong when M is small and is usually not satisfied when second order derivatives are not calculated. The property (6.31) recalls condition (iii) of theorem 5.1, which writes when (G_k) and (G_k^{-1}) are bounded:

$$(G_k - G_*) Z_* t_k = o(\|d_k\|). \quad (6.34)$$

Therefore, (6.31) is usually stronger than (6.34) and, in fact, is satisfied by some subsequences of subscripts when (G_k) is updated by the BFGS formula (see Gilbert (1986,a) and the discussion following the proof of theorem 5.1). The property (6.32) concerns the comparison of the tangent step and the restoration step.

The next theorem shows that for the subsequences \mathbb{K} for which (6.30) with M small enough or (6.31) or (6.32) is satisfied, the rule (6.3)–(6.6) will give $\rho_k = 1$ for all but finitely many k in \mathbb{K} . Therefore, the unit step-size will be admissible either when G_* is correctly approximated by G_k (properties (6.30) and (6.31)) or when t_k is of the same order of magnitude as r_{k+1} (property (6.32)).

Property (6.33) is more particular. In concrete algorithms using the update scheme (6.20), (6.25) and (6.26) with the update criterion (6.27), neither of the properties (6.31)–(6.33) is satisfied for the entire sequence. These properties are satisfied only for subsequences \mathbb{K} . The result obtained in theorem 6.3 with property (6.33) is then used to prove that only properties (6.31) or (6.32) (and not (6.33)) may occur in the considered algorithm (see Gilbert (1986,a)).

Theorem 6.3. Suppose that assumptions A and B are satisfied. Let (x_k) , (y_k) and (G_k) be the sequences generated by algorithm RQN with α in $]0, 1/2[$. Suppose that (x_k) and (y_k) are in ω , that (y_k) converges to x_* and that the matrices G_k are nonsingular and satisfy (6.28) with a positive constant h independent of k . Let K be a subsequence of subscripts. Then,

- (i) there exists a positive constant \bar{M} that depends only on c , α and h such that if (6.30) is satisfied with $M < \bar{M}$ then $\rho_k = 1$ for all but finitely many k in K ,
- (ii) if (6.31) or (6.32) is satisfied then $\rho_k = 1$ for all but finitely many k in K ,
- (iii) if (6.33) is satisfied, then $r_{k+1} = o(\|r_k\| \|t_k\|)$ for k in K .

Proof. Since (y_k) converges, proposition 6.2 shows that (p_k) is bounded and by (6.19), p_k is modified finitely often. So we can suppose that $p_k = p$ for all k . By Taylor's theorem, we expand $\Theta_p(y_k + t_k + r_{k+1})$ at the second order in t_k and the first order in r_{k+1} . First, note that (y_k) converging to x_* and (G_k^{-1}) being bounded, (x_k) converges to x_* . Then, we have

$$c(x_{k+1}) = c(y_k) + \frac{1}{2} c''(x_*) \cdot t_k^2 + o(\|t_k\|^2). \quad (6.35)$$

We also have

$$\begin{aligned} f(y_k + e_k) &= f(y_k) + f'(y_k) \cdot t_k + (\lambda(y_k), c(x_{k+1})) \\ &\quad + \frac{1}{2} f''(x_*) \cdot t_k^2 + o(\|t_k\|^2) + o(\|r_{k+1}\|) \end{aligned}$$

and using the estimate (6.35), we get

$$\begin{aligned} f(y_k + e_k) &= f(y_k) + f'(y_k) \cdot t_k + (\lambda(y_k), c(y_k)) \\ &\quad + \frac{1}{2} (L_* t_k, t_k) + o(\|t_k\|^2) + o(\|r_{k+1}\|). \end{aligned} \quad (6.36)$$

On the other hand, expanding $c(y_k + e_k)$ about y_k we obtain

$$c(y_k + e_k) = c(y_k) - c(x_{k+1}) + \frac{1}{2} c''(y_k) \cdot t_k^2 + o(\|t_k\|^2) + o(\|r_{k+1}\|)$$

and using again (6.35), we get

$$c(y_k + e_k) = o(\|t_k\|^2) + o(\|r_{k+1}\|). \quad (6.37)$$

Let us define

$$\Delta_k := f'(y_k) \cdot t_k - (p - \|\lambda(y_k)\|_\infty) \|c(y_k)\|_1,$$

which is negative by (6.28) and (6.17). Finally, (6.36) and (6.37) give

$$\Theta_p(y_k + e_k) \leq \Theta_p(y_k) + \Delta_k + \frac{1}{2} (L_* t_k, t_k) + o(\|t_k\|^2) + o(\|r_{k+1}\|).$$

But $t_k = Z_*^- Z_* t_k + o(\|t_k\|)$ and the boundedness of (G_k) allows us to write $g(y_k) = -G_k Z_* t_k + o(\|t_k\|)$. Therefore, using $f'(y_k) \cdot t_k = -(G_k^{-1} g(y_k), g(y_k)) = -(G_k Z_* t_k, Z_* t_k) + o(\|t_k\|^2)$, we obtain

$$\begin{aligned} & \Theta_p(y_k + e_k) - \Theta_p(y_k) - \alpha \Delta_k \\ & \leq \left(\frac{1}{2} - \alpha\right) \Delta_k - \frac{1}{2} ((G_k - G_*) Z_* t_k, Z_* t_k) + o(\|t_k\|^2) + o(\|r_{k+1}\|). \end{aligned} \quad (6.38)$$

Using this inequality we now prove the theorem. Suppose that the step-size ρ_k is different from 1 for k in a subsequence \mathcal{K} . Then, according to Armijo's rule (6.4)–(6.6), the left hand side of (6.38) is positive and we have:

$$-\Delta_k \leq -\frac{1}{1-2\alpha} ((G_k - G_*) Z_* t_k, Z_* t_k) + o(\|t_k\|^2) + o(\|r_{k+1}\|).$$

Using the inequality $C_1 \|t_k\| \leq \|g(y_k)\|$ (where C_1 is a positive constant that depends only on c and h), the property (6.28), the inequality (6.17), the definition (4.10) of r_{k+1} and (6.35), we can obtain a lower bound for the left hand side:

$$\begin{aligned} & h C_1^2 \|t_k\|^2 + p \|c(y_k)\|_1 \\ & \leq -\frac{1}{1-2\alpha} ((G_k - G_*) Z_* t_k, Z_* t_k) + o(\|t_k\|^2) + o(\|c(y_k)\|_1) \end{aligned} \quad (6.39)$$

Now, if one of the properties (6.30) with $M < \bar{M} := (1-2\alpha)hC_1^2/\|Z_*\|^2$ or (6.31) or (6.32) is verified, this inequality leads to

$$\|t_k\|^2 + \|c(y_k)\|_1 = o(\|t_k\|^2) + o(\|c(y_k)\|_1)$$

which shows that \mathcal{K} must be finite. This proves statements (i) and (ii) of the theorem. It remains to prove (iii). With (6.33), inequality (6.39) is valid for k in \mathcal{K} and $t_k = o(\|r_k\|)$ implies for k in \mathcal{K} :

$$\|t_k\|^2 + \|c(y_k)\|_1 = o(\|r_k\| \|t_k\|) + o(\|t_k\|^2) + o(\|c(y_k)\|_1),$$

from which we deduce

$$\|t_k\|^2 + \|c(y_k)\|_1 = o(\|r_k\| \|t_k\|).$$

Then, this estimate, the definition of r_{k+1} and (6.35) give

$$r_{k+1} = o(\|r_k\| \|t_k\|) + o(\|t_k\|^2) = o(\|r_k\| \|t_k\|)$$

which is the estimate in (iii). \square

7. Conclusion

We have studied in this paper the local and global convergence of a variable metric algorithm for equality constrained optimization in which the order of the updated matrices is $n-m$. This reduced method can be seen as making a link between GRG-like methods which are feasible methods ($c(x_k) = 0$ for all k) with reduced matrices (of order $n-m$) and the SQP method which is an unfeasible method with full matrices (of order n): the studied algorithm is indeed an unfeasible method with reduced matrices. The algorithm inherits also the good properties of both methods (reduced metrics, superlinear convergence and unfeasibility) and shows, in particular, that *locally* only one restoration step is necessary to obtain the superlinear convergence of GRG-like methods when the reduced matrices are correctly approximated.

The global convergence is obtained by Han's technique to globalize the SQP method. The l_1 penalty function is used as merit function and is decreased along an arc-shaped search path. Conditions for the asymptotic admissibility of the unity step-size are given that turn out to be satisfied in practice.

An important facet of the method has not been tackled here and is reported elsewhere (Gilbert (1987)). This concerns the update of the reduced matrices G_k . This one is based on a secant equation using the change in the reduced gradient g . The fact that the gradient of g at x_* (see (4.2)) is not equal to G_* (and cannot be equal because $\nabla g(x_*)$ is an $(n-m) \times n$ matrix while G_* is of order $n-m$) leads to an alternative. Either the reduced gradient is evaluated twice per iteration, at y_k and x_{k+1} , or it is evaluated only once per iteration, at y_k . In the first case, the change $g(x_{k+1}) - g(y_k)$ is used in the secant equation and the matrices G_k are updated at each iteration but with the inconvenience of having to linearize the constraints twice per iteration: see Coleman and Conn (1984) and Gilbert (1986,a). In the second case, the change $g(y_{k+1}) - g(y_k)$ is used in the secant equation but usually the matrices G_k cannot anymore be updated at each iteration. An update criterion has to be introduced in order to decide when an update is appropriate. Despite this, the superlinear convergence can be achieved either in a local framework (see Nocedal and Overton (1985) for the algorithm (1.12)–(1.14)) or in a global framework (see Gilbert (1986,a) for the algorithm RQN of Section 6).

Acknowledgments

This work was supported in part during 1984-1985 by the "Centre d'Etudes Nucléaires", Fontenay-aux-Roses (France) under a grant from the Commission of the European Community and in part by the "Institut National de Recherche en

Informatique et en Automatique" (INRIA), Le Chesnay (France). I would like to thank those organizations and to express my gratitude to J.F. Bonnans for helpful discussions and suggestions and for his friendly and constant advice.

References

- J. Abadie, J. Carpentier (1969). Generalization of the Wolfe reduced gradient method to the case of nonlinear constraints. Optimization, R. Fletcher (ed.), Academic Press, London.
- A. Ben-Israel, T.N.E. Greville (1974). *Generalized inverses: theory and applications*. John Wiley & Sons.
- J. Blum, J.Ch. Gilbert, B. Thooris (1985). Parametric identification of the plasma current density from the magnetic measurements and the pressure profile, code IDENTC. Report of JET contract number JT3/9008. Centre d'Etudes Nucléaires (DRFC), B.P. 6, 92265 Fontenay-aux-Roses (France).
- P.T. Boggs, J.W. Tolle, P. Wang (1982). On the local convergence of quasi-Newton methods for constrained optimization. SIAM Journal on Control and Optimization 20/2, 161-171.
- J.F. Bonnans (1984). Asymptotic admissibility of the unity stepsize in exact penalty methods I: equality-constrained problems. Rapport de recherche de l'INRIA RR-273. 78153 Le Chesnay, France.
- J.F. Bonnans, D. Gabay (1984). Une extension de la programmation quadratique successive. Lecture Notes in Control and Information Sciences 63, 16-31. A. Bensoussan, J.L. Lions (eds.). Springer-Verlag.
- R.H. Byrd (1984). On the convergence of constrained optimization methods with accurate Hessian information on a subspace. Report CU-CS-270-84. Department of Computer Science, University of Colorado, Boulder, Colorado 80309.
- R.H. Byrd (1985). An example of irregular convergence in some constrained optimization methods that use the projected Hessian. Mathematical Programming 32, 232-237.
- R.H. Byrd, R.B. Schnabel (1986). Continuity of the null space basis and constrained optimization. Mathematical Programming 35, 32-41.
- R.M. Chamberlain, C. Lemaréchal, H.C. Pedersen, M.J.D. Powell (1982). The watchdog technique for forcing convergence in algorithms for constrained optimization. Mathematical Programming Study 16, 1-17.
- T.F. Coleman, A.R. Conn (1982,a). Nonlinear programming via an exact penalty function: asymptotic analysis. Mathematical Programming 24, 123-136.
- T.F. Coleman, A.R. Conn (1982,b). Nonlinear programming via an exact penalty function: global analysis. Mathematical Programming 24, 137-161.

- T.F. Coleman, A.R. Conn (1984). On the local convergence of a quasi-Newton method for the nonlinear programming problem. *SIAM Journal on Numerical Analysis* 21/4, 755-769.
- Y.M. Danilin, B.N. Pschenichny (1965). *Numerical methods in extremal problems*. MIR, Moscow. (English translation, 1978).
- J.E. Dennis, J.J. Moré (1974). A characterization of superlinear convergence and its application to quasi-Newton methods. *Mathematics of Computation* 28/126, 549-560.
- J.E. Dennis, J.J. Moré (1977). Quasi-Newton methods, motivation and theory. *SIAM Review* 19, 46-89.
- R. Fletcher (1981). *Practical methods of optimization. Vol. 2: Constrained Optimization*. Wiley.
- D. Gabay (1975). Efficient convergence of implementable gradient algorithms and step-size selection procedures for constrained minimization. *International Computing Symposium 1975*. F. Gelenbe, K.D. Potier (eds). North-Holland Publishing Company.
- D. Gabay (1982,a). Minimizing a differentiable function over a differentiable manifold. *Journal of Optimization Theory and its Application* 37/2, 171-219.
- D. Gabay (1982,b). Reduced quasi-Newton methods with feasibility improvement for nonlinearly constrained optimization. *Mathematical Programming Study* 16, 18-44.
- D. Gabay, D.G. Luenberger (1976). Efficiently converging minimization methods based on the reduced gradient. *SIAM Journal on Control and Optimization* 14/1, 42-61.
- J.Ch. Gilbert (1986,a). Une méthode à métrique variable réduite en optimisation avec contraintes d'égalité non linéaires. Rapport de recherche de l'INRIA RR-482. 78153 Le Chesnay, France.
- J.Ch. Gilbert (1986,b). Une méthode de quasi-Newton réduite en optimisation sous contraintes avec priorité à la restauration. *Lecture Notes in Control and Information Sciences* 83, 40-53. A Bensoussan, J.L. Lions (eds.). Springer-Verlag.
- J.Ch. Gilbert (1986,c). Sur quelques problèmes d'identification et d'optimisation rencontrés en physique des plasmas. Thèse de doctorat de l'université Paris VI.
- J. Ch. Gilbert (1987). Mise à jour de la métrique dans les méthodes de quasi-Newton réduites en optimisation avec contraintes d'égalité. *RAIRO Mathematical Modelling and Numerical Analysis* (to appear).
- P.E. Gill, W. Murray, M.A. Saunders, G.W. Stewart, M.H. Wright (1985). Properties of a representation of a basis for the null space. *Mathematical Programming* 33, 172-186.
- J. Goodman (1985). Newton's method for constrained optimization. *Mathematical Programming* 33, 162-171.
- S.P. Han (1976). Superlinearly convergent variable metric algorithms for general

- nonlinear programming problems. *Mathematical Programming* 11, 263-282.
- S.P. Han (1977). A globally convergent method for nonlinear programming. *Journal of Optimization Theory and its Application* 22/3, 297-309.
- N. Maratos (1978). Exact penalty function algorithms for finite dimensional and control optimization problems. Ph.D. Thesis, University of London.
- D.O. Mayne, E. Polak (1982). A superlinearly convergent algorithm for constrained optimization problems. *Mathematical Programming Study* 16, 45-61.
- H. Mukai, E. Polak (1978). On the use of approximations in algorithms for optimization problems with equality and inequality constraints. *SIAM Journal on Numerical Analysis* 15/4, 674-693.
- J. Nocedal, M.L. Overton (1985). Projected Hessian updating algorithms for nonlinearly constrained optimization. *SIAM Journal on Numerical Analysis* 22/5, 821-850.
- M.J.D. Powell (1978). The convergence of the variable metric methods for nonlinearly constrained optimization calculations. *Nonlinear Programming* 3, O.L. Mangasarian, R.R. Meyer, S.M. Robinson (eds.), Academic Press, 27-63.
- L. Schwartz (1981). *Cours d'analyse, tome I*. Hermann.
- J. Stoer (1984). Principles of sequential quadratic programming methods for solving nonlinear programs. Proceedings of the NATO ASI on Computational Mathematical Programming, Bad Windsheim, Germany.
- R.B. Wilson (1963). A simplicial algorithm for concave programming. Ph.D. thesis. Graduate School of Business Administration, Harvard Univ., Cambridge, MA.
- Y. Yuan (1985). An only 2-step Q-superlinear convergence example for some algorithms that use reduced Hessian approximations. *Mathematical Programming* 32, 224-231.